

Department of Food Hygiene and Environmental Health
Faculty of Veterinary Medicine
University of Helsinki
Helsinki

**DETERMINANTS OF
LISTERIA MONOCYTOGENES
STRESS RESPONSES**

Anna Pöntinen

ACADEMIC DISSERTATION

To be presented, with the permission of the Faculty of Veterinary Medicine of the University of Helsinki, for public examination in Auditorium 6, Metsätalo building, Unioninkatu 40, Helsinki, on 14 June 2019, at 12 noon.

Helsinki 2019

- Supervising Professor** Professor Miia Lindström, DVM, Ph.D.
Department of Food Hygiene and Environmental Health
Faculty of Veterinary Medicine
University of Helsinki
Helsinki, Finland
- Supervisors** Professor emeritus Hannu Korkeala, DVM, Ph.D., M.Soc.Sc.
Department of Food Hygiene and Environmental Health
Faculty of Veterinary Medicine
University of Helsinki
Helsinki, Finland
- Professor Miia Lindström, DVM, Ph.D.
Department of Food Hygiene and Environmental Health
Faculty of Veterinary Medicine
University of Helsinki
Helsinki, Finland
- Reviewed by** Professor Wilhelm Tham, DVM, Ph.D.
School of Hospitality, Culinary Arts and Meal Science
Örebro University
Örebro, Sweden
- Professor José Vázquez-Boland, DVM, Ph.D.
Infection Medicine
Edinburgh Medical School
University of Edinburgh
Edinburgh, United Kingdom
- Opponent** Professor emeritus Atte von Wright, M.Sc., Ph.D.
Institute of Public Health and Clinical Nutrition
University of Eastern Finland
Kuopio, Finland

ISBN 978-951-51-5292-3 (paperback)
ISBN 978-951-51-5293-0 (PDF)
<http://ethesis.helsinki.fi>

Unigrafia
Helsinki 2019

ABSTRACT

Listeria monocytogenes is a remarkable bacterium, as it is able to shift from a capable environmental saprophyte into a severe intracellular pathogen. As a strictly foodborne pathogen, *L. monocytogenes* poses a notable risk, particularly to those consumers among the risk groups for whom invasive listeriosis is potentially fatal. Furthermore, modern consumption habits and increasingly favoured ready-to-eat foods, often consumed without proper heating, increase the risk of acquiring the foodborne disease. The aim of this study was to investigate the genetic mechanisms conferring wide-ranging stress tolerance in *L. monocytogenes*.

Two-component systems, comprising a sensor histidine kinase and a cognate response regulator, aid bacteria in sensing and adapting to changes in both surrounding environmental as well as intracellular conditions. The histidine kinases, in particular, have lacked comprehensive studies on their roles in the stress tolerance of *L. monocytogenes*. Thus, histidine kinases were studied by expressional analyses under cold conditions and by mutationally disrupting each histidine kinase-encoding gene in a parental model strain, *L. monocytogenes* EGD-e. The modified strains were individually challenged at high (42.5 °C) and low (3.0 °C) temperatures, high (9.4) and low (5.6) pH levels, and high salt (6% NaCl), ethanol (3.5 vol%) and hydrogen peroxide (5 mM) concentrations. Expressional studies and growth experiments on genetically modified strains proved *lisK* and *yycG* to respectively play central roles in the acclimation and immediate growth of *L. monocytogenes* at low temperatures. The most substantial increase in gene expression under cold conditions was that of the chemotaxis gene *cheY* with 236-fold upregulation at 3 °C. The disrupted Δ *liaS* strain displayed impaired growth in response to all the other stresses, particularly at a high temperature and under osmotic stress. These studies demonstrated the prominent importance of the histidine kinase-encoding genes *yycG* and *lisK* to cold tolerance and *liaS*, with roles in the growth of *L. monocytogenes* under multiple stresses.

To shed light on the accessory genetic mechanisms that cause large strain variation in *L. monocytogenes* in withstanding heat treatments, heat resistance-conferring traits were further investigated by means of whole-genome sequencing. Comparing the complete genomes of heat-resistant *L. monocytogenes* AT3E and -sensitive AL4E strains revealed the heat-resistant strain to harbour a novel 58-kb plasmid, pLM58, which was absent in the sensitive strain. Furthermore, curing of the plasmid in AT3E produced a marked decrease in heat resistance from virtually no reduction to a 1.1 cfu/ml log₁₀ reduction at 55.0 °C. In pLM58, a 2,155-bp open reading frame annotated as an ATP-dependent ClpL protease-encoding gene was identified. Conjugation of the coding sequence and the putative promoter of the *clpL*

gene into a natively heat-sensitive *L. monocytogenes* 10403S strain, in turn, enhanced the survival of the strain from a 1.2 cfu/ml log₁₀ reduction to a 0.4 cfu/ml log₁₀ reduction in heat challenge at 55.0 °C. In this study, we presented the first evidence of plasmid-mediated heat resistance in *L. monocytogenes* and identified the protease ClpL to be a novel plasmid-borne heat-resistance mediator.

The emerging resistance of *L. monocytogenes* to benzalkonium chloride, a quaternary ammonium compound widely used as a detergent in food-processing facilities, is a significant concern for food safety and public health. The resistance of 392 *L. monocytogenes* isolates from Finland (n = 197) and Switzerland (n = 195) to benzalkonium chloride was assessed. A minimal inhibitory concentration of 20 µg/ml was defined. Altogether, 11.5% of the strains proved to be resistant to benzalkonium chloride. Serotype 1/2c harboured the highest prevalence, 32.4% (11/34), of benzalkonium chloride-resistant strains, while in total, most of the resistant strains belonged to serotype 1/2a. Altogether, 68.9% of the resistant strains harboured at least one of the efflux pump system-encoding genes, *bcrABC*, *emrE* or *qacH*, known to confer benzalkonium chloride resistance in *L. monocytogenes*. We found resistant strains with partially or completely efflux pump-dependent benzalkonium chloride resistance, with the exception of the known resistance-mediating efflux pumps, suggesting the existence of other resistance-contributing efflux pump systems. The lacking of known efflux pump system-encoding genes in addition to efflux pump-independent benzalkonium chloride resistance, in turn, indicates the contribution of completely novel benzalkonium chloride resistance mechanisms.

The aim of these studies was to shed light on the genes contributing to the versatile stress tolerance abilities and strain variation of the severe foodborne pathogen, *L. monocytogenes*. Knowledge of such traits may aid in developing targeted strategies and measures to identify and control the contamination and risks caused, in particular, by stress-tolerant *L. monocytogenes* strains.

ACKNOWLEDGEMENTS

This thesis study was conducted at the Department of Food Hygiene and Environmental Health of the Faculty of Veterinary Medicine, University of Helsinki. The Doctoral Programme in Food Chain and Health, the Finnish Ministry of Agriculture and Forestry, the Finnish Veterinary Foundation, the Finnish Foundation of Veterinary Research and the Walter Ehrström Foundation are gratefully acknowledged for their financial support.

Professor Miia Lindström and Professor emeritus Hannu Korkeala are warmly acknowledged for their insightful supervision of this thesis. Professors Wilhelm Tham and José Vázquez-Boland are gratefully appreciated for reviewing this thesis in such an expert and prompt manner and Dr Roy Siddall for carefully revising the English language.

I wish to equally acknowledge each person at the department for making our study groups a wonderful scientific community. Some of you I must salute one by one, as I am forever in your debt: Many thanks to Annukka Markkula and Mirjami Mattila, who both taught me how to survive in the lab and whose door and email were always open for my questions. My all-time officemates, Mariella Aalto-Araneda, Hanna Castro, Marita Isokallio, Noora Salin and Katja Selby, I sincerely thank for all the support and invigorating conversations. Jari Aho, Anneli Luoti, Henna Niinivirta, Erika Pitkänen, Anu Seppänen and Heimo Tasanen are appreciated for their indispensable work in the lab, and also Hanna Korpunen for the great seminars and origami. I dedicate special thanks to Kirsi Ristkari for efficiently completing the nearly inhuman pipetting sessions and to Esa Penttinen for constantly showing responsibility and initiative far beyond his duties. Johanna Laine deserves praise for around-the-clock assistance with bureaucracy.

Mariella Aalto-Araneda, Claudia Guldemann, Annukka Markkula, Anja Meier, Mikael Skurnik and Taurai Tasara are thanked for fruitful co-authoring. Mirko Rossi is acknowledged for all the help in genome analyses, Panu Somervuo for solving many of the mathematical oddities and Per Johansson for sharing his vast knowledge, regardless of the subject.

I am truly grateful to have known wonderful mates throughout these years. Mariella, not only a colleague but also a friend: thank you. My training companion Johanna, cheers for ushering me on a run when I needed it the most. My precious friend Heidi, without you I simply could not operate. Darlings Kati-Maria and Adrian, thank you for all the support and all the late nights and early mornings around endless cups of gin and the Finest English Breakfast Tea. *Le temps détruit tout – sauf l'amitié.*

Finally, I warmly thank my endearingly bizarre family for always supporting me in my obstinate choices. And Dad, for giving me a bit of your perseverance and all of your time, this is for you.

CONTENTS

ABSTRACT	3
ACKNOWLEDGEMENTS.....	5
CONTENTS.....	6
LIST OF ORIGINAL PUBLICATIONS.....	9
ABBREVIATIONS.....	10
1 INTRODUCTION	12
2 REVIEW OF THE LITERATURE	14
2.1 <i>Listeria</i> spp. and <i>Listeria monocytogenes</i>	14
2.1.1 <i>Listeria</i> spp.	14
2.1.2 Characteristics of <i>L. monocytogenes</i>	14
2.1.3 <i>L. monocytogenes</i> contamination	16
2.2 Listeriosis.....	17
2.2.1 Human listeriosis	17
2.2.2 Animal listeriosis.....	20
2.3 Plasmids in <i>Listeria</i> spp.	21
2.4 Two-component systems.....	22
2.5 Stress responses of <i>L. monocytogenes</i>	24
2.5.1 General stress responses and stress tolerance mechanisms.....	25
2.5.2 Strain diversity in stress responses	35
2.5.3 Plasmid-mediated stress tolerance in <i>L. monocytogenes</i>	36
2.5.4 Two-component systems in the stress responses of <i>L. monocytogenes</i>	37
3 AIMS OF THE STUDY	40
4 MATERIALS AND METHODS	41

4.1	<i>L. monocytogenes</i> strains, isolates and plasmids (I–IV)	41
4.2	Typing (III, IV)	41
4.2.1	Serotyping (III, IV)	41
4.2.2	Multilocus sequence typing (III, IV)	41
4.3	Growth conditions (I–IV)	43
4.4	Transcriptional analyses (I)	44
4.4.1	Total RNA extraction (I)	44
4.4.2	Real-time quantitative reverse transcription-PCR (I)	44
4.5	Genetic modifications (I–III)	49
4.5.1	Knock-out deletion (I, II)	49
4.5.2	Complementation (I, II)	50
4.5.3	Plasmid curing (III)	50
4.5.4	Plate mating (III)	51
4.5.5	Introducing <i>clpL</i> into a heat-sensitive <i>L. monocytogenes</i> strain (III)	51
4.6	Phenotypic characterisation of <i>L. monocytogenes</i> strains and isolates (I–IV)	52
4.6.1	Growth curve analyses (I–III)	52
4.6.2	Heat resistance assay (III)	52
4.6.3	Minimum and maximum growth temperatures (I–III)	53
4.6.4	Benzalkonium chloride resistance and impact of reserpine (IV)	53
4.7	Genomic analyses (III)	53
4.7.1	DNA extraction (III)	53
4.7.2	Sequencing (III)	54
4.7.3	Annotation and comparative genomic analysis (III)	54
4.8	Statistical analyses (I–IV)	54
5	RESULTS	55

5.1	Roles of two-component-system histidine kinases in the stress tolerance of <i>L. monocytogenes</i> (I, II)	55
5.1.1	Relative expression levels of histidine kinase-encoding genes at low temperatures (I)	55
5.1.2	Growth of genetically modified <i>L. monocytogenes</i> under various stresses (I, II)	57
5.1.3	Minimum and maximum growth temperatures (I, II)	59
5.2	Plasmid-mediated heat resistance in <i>L. monocytogenes</i> (III)	59
5.2.1	Novel 58-kb plasmid contributes to the heat resistance of <i>L. monocytogenes</i> AT3E (III)	59
5.2.2	Plasmid-borne <i>clpL</i> increases heat resistance in <i>L. monocytogenes</i> (III)	60
5.2.3	pLM58 is putatively nonconjugative (III)	61
5.3	<i>L. monocytogenes</i> strains display high variability under severe stresses (III, IV)	61
5.3.1	Resistance and growth of <i>L. monocytogenes</i> strains vary at high temperatures (III)	61
5.3.2	<i>L. monocytogenes</i> isolates display variable resistance to benzalkonium chloride (IV)	61
6	DISCUSSION	63
6.1	Extensive roles of two-component-system histidine kinases in the stress tolerance of <i>L. monocytogenes</i> (I–II)	63
6.2	pLM58-borne ClpL-mediated heat resistance in <i>L. monocytogenes</i> (III)	65
6.3	Characteristics of benzalkonium chloride resistance in <i>L. monocytogenes</i> (IV)	67
7	CONCLUSIONS	69
	REFERENCES	70

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following publications, referred to in the text by their Roman numerals:

- I. Pöntinen, A., Markkula, A., Lindström, M. & Korkeala, H. (2015) Two-component-system histidine kinases involved in growth of *Listeria monocytogenes* EGD-e at low temperatures. *Appl Environ Microbiol* 81: 3994-4004.
- II. Pöntinen, A., Lindström, M., Skurnik, M. & Korkeala, H. (2017) Screening of the two-component-system histidine kinases of *Listeria monocytogenes* EGD-e. LiaS is needed for growth under heat, acid, alkali, osmotic, ethanol and oxidative stresses. *Food Microbiol* 65: 36-43.
- III. Pöntinen, A., Aalto-Araneda, M., Lindström, M. & Korkeala, H. (2017) Heat resistance mediated by pLM58 plasmid-borne ClpL in *Listeria monocytogenes*. *MSphere* 2: e00364-17.
- IV. Meier, A.B., Guldemann, C., Markkula, A., Pöntinen, A., Korkeala, H. & Tasara, T. (2017) Comparative phenotypic and genotypic analysis of Swiss and Finnish *Listeria monocytogenes* isolates with respect to benzalkonium chloride resistance. *Front Microbiol* 8: 397.

These original articles have been reprinted with the kind permission of their copyright holders: the American Society for Microbiology (I, III), Elsevier (II) and Frontiers Media SA (IV).

ABBREVIATIONS

A	Adenine
ADI	Arginine deiminase
AFLP	Amplified fragment length polymorphism
Asp	Aspartate
ATP	Adenosine triphosphate
AUC	Area under the curve
a _w	Water activity
BC	Benzalkonium chloride
BC ^r	Benzalkonium chloride-resistant
BHI	Brain–heart infusion
bp	Base pair
C	Cytosine
Cap	Cold acclimation protein
CC	Clonal complex
cDNA	Complementary DNA
CDS	Coding sequence
cfu	Colony forming unit
Csp	Cold shock protein
DNA	Deoxyribonucleic acid
EC	Epidemic clone
ET	Electrophoretic type
FA	Fatty acid
G	Guanine
GABA	γ-aminobutyrate
GDA	Glutamate decarboxylase
GWAS	Genome-wide association studies
HGT	Horizontal gene transfer
His	Histidine
HK	Histidine kinase
HPt	His-containing phosphotransfer protein
Hsp	Heat shock protein
ICE	Integrative conjugative element
Inc	Incompatibility group
LB	Luria-Bertani (lysogeny broth)
LD	Linkage disequilibrium
LIPI	<i>Listeria</i> pathogenicity island
LLS	Listeriolysin S
MGE	Mobile genetic element
MH	Mueller-Hinton
MIC	Minimal inhibitory concentration
MLEE	Multilocus enzyme electrophoresis

MLST	Multilocus sequence typing
MLVA	Multiple-locus variable-number tandem repeat analysis
Mpf	Mating pair formation
mRNA	Messenger ribonucleic acid
NGS	Next-generation sequencing
OD ₆₀₀	Optical density at 600 nm
ORF	Open reading frame
PCA	Plate count agar
PCR	Polymerase chain reaction
PFGE	Pulsed-field gel electrophoresis
QAC	Quaternary ammonium compound
RAPD	Randomly amplified polymorphic DNA
REA	Restriction endonuclease analysis
RFLP	Restriction fragment length polymorphism
RNA	Ribonucleic acid
ROS	Reactive oxygen species
RR	Response regulator
rRNA	Ribosomal ribonucleic acid
RTE	Ready-to-eat
RT-qPCR	Real-time quantitative reverse transcription-PCR
Sap	Salt acclimation protein
SD	Standard deviation
SMRT	Single-molecule real-time
SNP	Single nucleotide polymorphism
SOE	Splicing-by-overlap extension
SSI	Stress survival islet
Ssp	Salt shock protein
ST	Sequence type
T	Thymine
T4SS	Type IV secretion system
TCS	Two-component signal transduction system
TS	Tryptic soy
WGS	Whole-genome sequencing

1 INTRODUCTION

The first plausible discovery of *Listeria monocytogenes* dates back to 1911 when the Swedish veterinarian Gustav Hülphers described a bacterium isolated from the necrotic liver tissue of a rabbit and, accordingly, named his discovery *Bacillus hepatis* (Hülphers, 1911). However, the isolate itself was not preserved, and it was only after the discovery by Murray *et al.* (1926) that a detailed description of the bacterium was established, often designated the first report on *L. monocytogenes*. The bacterium was, at that time, named *Bacterium monocytogenes* due to it causing a characteristic proliferation of monocytes (Murray *et al.*, 1926). In 1927, the bacterium was newly named as *Listerella hepatolytica* (Pirie, 1927), and in 1940 it finally received its present name, *L. monocytogenes* (Pirie, 1940), in honour of Lord Lister, the *primus motor* of antiseptics and hygienic surgical measures.

Although *L. monocytogenes* was isolated from a human meningitis patient as early as in 1918 (Dumont and Cotoni, 1921), the first confirmed human listeriosis cases were reported a decade later (Nyfeldt, 1929). The foodborne nature of listeriosis was only explicitly recognised after distinct outbreaks in the 1980s (Schlech *et al.*, 1983, Fleming *et al.*, 1985, Linnan *et al.*, 1988), although the association between foodstuffs and listeriosis was already conjectured in the 1950s and after (Potel, 1953, Gray, 1963). Since then, foodborne listeriosis outbreaks have unfortunately become commonplace, and as consumption habits evolve to favour ready-to-eat foods without proper heating, *L. monocytogenes* continues to be a notable food safety hazard.

Besides being a foodborne pathogen, *L. monocytogenes* dwells ubiquitously in the environment, from soil and vegetation to water supplies and cultivated land. Indeed, it is characteristically able to shift from being an environmental saprophyte to an intracellular pathogen (Chaturongakul *et al.*, 2008, Cossart, 2011). Its versatile ability to endure and overcome stressful conditions plays a fundamental role in the transmission processes of the bacterium from the environment to the vehicle foods and feed and, ultimately, into the animal or human host. Notably, *L. monocytogenes* copes well with many of the stressors, such as refrigeration and acidic and osmotic conditions, that are frequently applied to limit the number of bacteria in food-processing environments and foodstuffs (Bucur *et al.*, 2018). Thus, identifying the genetic mechanisms conferring this agile phenotypic stress tolerance is a pivotal step in understanding the survival strategies of the bacterium and, moreover, in controlling *L. monocytogenes* contamination along the food chain.

Genetic research in bacteriology has undergone marked changes in the recent years, as study methods have evolved from traditional molecular techniques to the emergence of novel whole-genome sequencing-based

approaches (Read and Massey, 2014). Next-generation sequencing methods have facilitated the study of accessory genetic mechanisms and their role in conferring stress resistance between bacterial strains and populations. Both genomic and phenotypic comparisons in the model *L. monocytogenes* strains have demonstrated how strain variability may affect the observations that we make based on the *in vitro* studies performed on these strains and lead to discrepant results (Bécavin *et al.*, 2014). High-throughput and genome-wide methods allow us to more efficiently acknowledge and take into account this strain variability and thereby avoid possible misconceptions in the interpretation of research findings due to naturally occurring differences between strains (Brul *et al.*, 2012, Bécavin *et al.*, 2014).

Expressional analyses and mutational studies have enabled the role of many genetic mechanisms to be ascertained in the stress tolerance of *L. monocytogenes*, such as stress shock proteins and membrane transport systems (Gandhi and Chikindas, 2007). Whole-genome sequencing, in turn, has provided further new tools to process hundreds to thousands of strains simultaneously and to identify even larger sets of culprit stress tolerance and virulence genes (Read and Massey, 2014, Kachroo *et al.*, 2019). However, many proteins encoded by unknown or putative genes still remain. Thus, the overall picture of the genetics underlying *L. monocytogenes* stress tolerance in foods and food-processing environments is anything but complete and warrants further investigations into the stress responses of the bacterium.

2 REVIEW OF THE LITERATURE

2.1 *Listeria* spp. and *Listeria monocytogenes*

2.1.1 *Listeria* spp.

Listeria spp. belong to the family *Listeriaceae* in the order *Bacillales* and class *Bacilli* of the phylum *Firmicutes* (McLauchlin and Rees, 2015). Currently, together with *Listeria monocytogenes* (*sensu stricto*) (Murray *et al.*, 1926, Pirie, 1940), the validated *Listeria* species (McLauchlin and Rees, 2015) include *L. grayi* (Larsen and Seeliger, 1966, Rocourt *et al.*, 1992), *L. innocua* (Seeliger, 1984), *L. ivanovii* (Seeliger *et al.*, 1984), *L. seeligeri* and *L. welshimeri* (Rocourt and Grimont, 1983). In addition, *L. marthii* (Graves *et al.*, 2010) and the phylogenetically more divergent *Listeria* species *L. rocourtaiae* (Leclercq *et al.*, 2010), *L. fleischmannii* (Bertsch *et al.*, 2013), and *L. weihenstephanensis* (Lang Halter *et al.*, 2013) have been described.

In recent years, several novel species have been isolated, including *L. aquatica* sp. nov., *L. cornellensis* sp. nov., *L. floridensis* sp. nov., *L. grandensis* sp. nov., *L. riparia* sp. nov. (den Bakker *et al.*, 2014), *L. booriae* sp. nov. and *L. newyorkensis* sp. nov. (Weller *et al.*, 2015). Additionally, subspecies *L. ivanovii* subsp. *ivanovii* and subsp. *londoniensis* have been designated within *L. ivanovii* (Boerlin *et al.*, 1992), as well as subspecies *L. fleischmannii* subsp. *fleischmannii* and subsp. *coloradonensis* within *L. fleischmannii* (den Bakker *et al.*, 2013). The formerly described *L. murrayi* (Welshimer and Meredith, 1971) has since been designated as a single species within *L. grayi* (Rocourt *et al.*, 1992).

2.1.2 Characteristics of *L. monocytogenes*

L. monocytogenes is a Gram-positive, facultative anaerobic, short rod (0.4–0.5 × 1–2 µm). It forms peritrichous flagella and manifests a characteristic tumbling motility at <30 °C. *L. monocytogenes* produces neither spores nor capsules, but it may develop filaments of ≥6 µm in length (Gray and Killinger, 1966, McLauchlin and Rees, 2015). It is β-haemolytic, catalase-positive and oxidase-negative, and typically ferments rhamnose but not xylose (Roberts *et al.*, 2006, McLauchlin and Rees, 2015). *L. monocytogenes* harbours a low average DNA G+C content (mol%) of 39% (Glaser *et al.*, 2001), varying from 36% to 42.5% (McLauchlin and Rees, 2015).

L. monocytogenes isolates form a structured population, currently divided into four evolutionary lineages (Orsi *et al.*, 2011, Tsai *et al.*, 2011) (Table 1). The majority of *L. monocytogenes* strains belong to lineages I and II (Doumith *et al.*, 2004), first described by Piffaretti *et al.* (1989). These

lineages are clearly distinguishable in regard to common sources and genetic characteristics, while lineages III and IV are comprised of more diverse strains.

A clonal genetic structure has been proposed for *L. monocytogenes*, supported by virulence gene allelic analysis, ribotyping and extensive linkage disequilibrium (LD) within the virulence genes and certain lineage I and II electrophoretic types (ETs) (Piffaretti *et al.*, 1989, Wiedmann *et al.*, 1997). However, based on the diversity of housekeeping, virulence and stress response genes, Nightingale *et al.* (2005b) later reported that while lineage I is indeed highly clonal, isolates of lineage II display greater genetic diversity. Furthermore, through genome-wide analyses, recombination has been determined to occur more commonly in lineage II than lineage I isolates (Orsi *et al.*, 2008). A high recombination rate could explain the diverse isolation sources of lineage II when compared to the seemingly more host-adapted lineage I (Orsi *et al.*, 2008). Besides atypical 4b serotypes (Roberts *et al.*, 2006, Orsi *et al.*, 2011), rhamnose-negative serotype 4c and 4a strains have been reported in lineages III and IV, respectively (Liu *et al.*, 2006).

Table 1. Summary of *L. monocytogenes* lineages and serotypes.

Lineage	Serotypes	Genetic characteristics	Sources	References
I	1/2b, 3b, 3c, 4b, 4d, 4e	Clonal	Various sources, human isolates overrepresented	(Piffaretti <i>et al.</i> , 1989, Bibb <i>et al.</i> , 1990, Nadon <i>et al.</i> , 2001, Ward <i>et al.</i> , 2004)
II	1/2a, 1/2c, 3a, 3c	Recombinant	Various sources, food-related and environmental isolates, sporadic human isolates	(Piffaretti <i>et al.</i> , 1989, Bibb <i>et al.</i> , 1990, Ward <i>et al.</i> , 2004)
III	4a, 4b (atypical), 4c	Diverse isolates, moderate recombination rates	Predominantly animal origin, rare human and food isolates	(Rasmussen <i>et al.</i> , 1995, Nadon <i>et al.</i> , 2001, Ward <i>et al.</i> , 2004, Nightingale <i>et al.</i> , 2005b, Roberts <i>et al.</i> , 2006)
IV	4a, 4b (atypical), 4c, 7	Few isolates to date	Few animal, human, and food isolates	(Roberts <i>et al.</i> , 2006) (IIIB); (Liu <i>et al.</i> , 2006, Ward <i>et al.</i> , 2008)

Typing

L. monocytogenes strains are phenotypically categorised into 13 serotypes, when also including serotype 4ab, based on cell surface heat-stable somatic (O) and heat-labile flagellar (H) antigens (Seeliger and Höhne, 1979). In addition to serotyping, other conventional phenotypic methods, such as multilocus enzyme electrophoresis (MLEE) (Bibb *et al.*, 1989, Piffaretti *et al.*,

1989) and phage typing (Audurier *et al.*, 1979, Rocourt *et al.*, 1985), can be applied to type *L. monocytogenes* isolates. More novel, genotypic subtyping methods include amplified fragment length polymorphism (AFLP) (Vos *et al.*, 1995, Keto-Timonen *et al.*, 2003), pulsed-field gel electrophoresis (PFGE) (Ojeniyi *et al.*, 1991, Ojeniyi *et al.*, 1996, Katzav *et al.*, 2006), PCR-based randomly amplified polymorphic DNA (RAPD) (Mazurier and Wernars, 1992, Lawrence *et al.*, 1993), restriction endonuclease analysis (REA) (Nocera *et al.*, 1990, Ericsson *et al.*, 1993), PCR-restriction fragment length polymorphism (RFLP) (Vines *et al.*, 1992), ribotyping (Graves *et al.*, 1991, Graves *et al.*, 1994) and DNA sequencing-based subtyping methods, such as multilocus sequence typing (MLST) (Salcedo *et al.*, 2003, Haase *et al.*, 2014, Moura *et al.*, 2016), multiple-locus variable-number tandem repeat analysis (MLVA) (Keim *et al.*, 2000, Murphy *et al.*, 2007, Lindstedt *et al.*, 2008) and single nucleotide polymorphism (SNP) typing (Nelson *et al.*, 2004). Epidemic clones (ECs) consist of phenotypically and genotypically closely related epidemic-associated *L. monocytogenes* isolates, even though isolates of the same epidemic clone might cause outbreaks geographically or temporally distant from each other (Piffaretti *et al.*, 1989, Kathariou, 2002). The first sequenced *L. monocytogenes* genome was that of the EGD-e strain (Glaser *et al.*, 2001). Subsequently, whole-genome sequencing (WGS)-based approaches have become increasingly common typing methods in bacteriology (Salipante *et al.*, 2015) and have facilitated large epidemiological outbreak investigations and surveillance of infectious diseases (Kwong *et al.*, 2016, Revez *et al.*, 2017, Schjørring *et al.*, 2017, Allam *et al.*, 2018).

2.1.3 *L. monocytogenes* contamination

L. monocytogenes is a ubiquitous saprophyte in the environment (Weis and Seeliger, 1975). It can be found in soil and vegetation, in water supplies and sewage, on feeding grounds, in animal and human faeces and in poorly prepared fodder (Weis and Seeliger, 1975, Husu *et al.*, 1990a, Renterghem *et al.*, 1991, MacGowan *et al.*, 1994, Unnerstad *et al.*, 2000, Pauly and Tham, 2003, Lyautey *et al.*, 2007a, Lyautey *et al.*, 2007b).

Besides being a saprophyte dwelling in the environment, *L. monocytogenes* has been isolated from numerous raw foods, with the highest prevalences reported in poultry (Miettinen *et al.*, 2001, Rørvik *et al.*, 2003) and a somewhat more seldom occurrence in other raw meats and fish (Autio *et al.*, 1999, Autio *et al.*, 2000, Markkula *et al.*, 2005). Although the total number of bacteria and *Listeria* spp. is generally fairly low in Finnish bulk tank milk (Ruusunen *et al.*, 2013), *L. monocytogenes* has also been isolated in raw milk samples (Husu, 1990, Ruusunen *et al.*, 2013, Castro *et al.*, 2017). Epidemiological studies have implied faecal contamination to be a putative cause of the frequent dissemination of *L. monocytogenes* in tank milk and raw meat products (Skovgaard and Nørrung, 1989, Husu, 1990,

Husu *et al.*, 1990b). A longitudinal study employing PFGE further indicated that *L. monocytogenes* contamination in tank milk may derive from udders or milking equipment (Castro *et al.*, 2018). *L. monocytogenes* has also been isolated in food-processing plants, where certain subtypes have persisted over long periods of months to several years (Miettinen *et al.*, 1999a, Vogel *et al.*, 2001, Hoffman *et al.*, 2003, Lundén *et al.*, 2003b, Keto-Timonen *et al.*, 2007, Bērziņš *et al.*, 2010, Almeida *et al.*, 2013). *L. monocytogenes* is thus potentially a highly challenging contaminant in food-processing environments.

The initial occurrence of *L. monocytogenes* in slaughterhouses and other food-processing premises in the production chain has frequently been indicated to derive from raw materials (Autio *et al.*, 2000, Berrang *et al.*, 2002, Autio *et al.*, 2004, Markkula *et al.*, 2005). However, tracing *L. monocytogenes* strains with phenotypic or genetic molecular typing methods has revealed contamination of the final products originating from sources other than raw material (Rørvik *et al.*, 1995, Autio *et al.*, 1999) or not exclusively incriminating the raw material as a contamination source (Norton *et al.*, 2001, Vogel *et al.*, 2001). Continuous listerial contamination in a food-processing facility may result from persistent strains dwelling in the processing environment rather than repeatedly deriving from incoming raw materials (Hoffman *et al.*, 2003). Recontamination of food products in the processing line after heating is a frequent source for the occurrence of *L. monocytogenes* in processed foods (Tompkin, 2002, Reij *et al.*, 2004). This is of importance particularly concerning ready-to-eat (RTE) foods, which are often consumed without heating.

2.2 Listeriosis

Of *Listeria* spp., *L. monocytogenes* and *L. ivanovii* are considered pathogenic (Nyfeldt, 1929, Ivanov, 1962, Vazquez-Boland *et al.*, 2001). *L. monocytogenes* is the predominant cause of listeriosis in both humans and animals, while *L. ivanovii* mainly leads to illness in ruminants (Ramage *et al.*, 1999, Vazquez-Boland *et al.*, 2001). Rare sporadic human cases caused by *L. ivanovii*, *L. innocua* and *L. grayi* have been reported, with predisposing factors and symptoms similar to those associated with *L. monocytogenes* (Cummins *et al.*, 1994, Lessing *et al.*, 1994, Perrin *et al.*, 2003, Snapir *et al.*, 2006, Rapose *et al.*, 2008, Guillet *et al.*, 2010).

2.2.1 Human listeriosis

The incidence of reported human listeriosis cases in EU/EEA countries is approximately 0.6 and in the USA 0.3 sporadic cases per 100 000 population annually (Silk *et al.*, 2012, CDC, 2013, ECDC, 2016). Even though invasive listeriosis is a fairly rare condition, it is a severe illness with an average case-

fatality rate of 20% to 30% (Fleming *et al.*, 1985, Mylonakis *et al.*, 1998, Mead *et al.*, 1999, de Valk *et al.*, 2005, CDC, 2013, Crim *et al.*, 2014). In patients suffering from listerial meningitis, mortality of up to 36% has been reported (Koopmans *et al.*, 2013).

Listeriosis is likely to manifest as sporadic cases, but outbreaks have also been reported (Fleming *et al.*, 1985, Linnan *et al.*, 1988, McLauchlin, 1996, Lyytikäinen *et al.*, 2000, Swaminathan and Gerner-Smidt, 2007, Angelo *et al.*, 2017, Schjørring *et al.*, 2017, Allam *et al.*, 2018). Most (99%) of the listeriosis cases are foodborne (Schlech *et al.*, 1983, Pinner *et al.*, 1992, Mead *et al.*, 1999, Swaminathan and Gerner-Smidt, 2007). However, aberrant transmission from animals to humans has been reported, mostly as localised infections among farmers and veterinarians (McLauchlin and Low, 1994, Regan *et al.*, 2005). As most listeriosis cases are sporadic and the incidence is low, and due to the long incubation period of invasive listeriosis and the subsequent delay in diagnosis, the vehicle foods are often challenging to trace. However, a high risk of acquiring listeriosis is frequently attributable to RTE foods with prolonged shelf lives and consumed without proper heating (Schwartz *et al.*, 1988, Pinner *et al.*, 1992, Cartwright *et al.*, 2013, Lopez-Valladares *et al.*, 2018). In recent years, outbreaks have been linked to a varied range of such foods, including caramel apples (Angelo *et al.*, 2017), ice cream products (Chen *et al.*, 2017) and cold-smoked salmon (Schjørring *et al.*, 2017).

L. monocytogenes is a facultative intracellular pathogen (Cossart, 2011, Radoshevich and Cossart, 2018). The virulence genes essential for the intracellular lifecycle of the bacterium are clustered in the chromosomal *Listeria* pathogenicity islands (LPIs) (Vázquez-Boland *et al.*, 2001). *L. monocytogenes* enters the host via the gastrointestinal tract and, while in the intestine, may secrete the bacteriocin compound listeriolysin S (LLS) to restrict the growth of competing commensal bacteria in the host (Cotter *et al.*, 2008, Quereda *et al.*, 2016). *L. monocytogenes* is able to invade the intestinal barrier through junctions between mucus-producing goblet cells and enterocytes, via epithelial M cells in the Peyer's patches, or through the tip of the intestinal villi where apoptotic epithelial cells extrude into the intestinal lumen (Jensen *et al.*, 1998, Nikitas *et al.*, 2011, Ribet and Cossart, 2015, Radoshevich and Cossart, 2018). *L. monocytogenes* is able to infect both macrophages and nonphagocytic cells and, once intracellular, can spread while evading the hostile extracellular environment (Berche *et al.*, 1988, Cossart, 2011, Pizarro-Cerdá *et al.*, 2012, Jones *et al.*, 2015). After penetrating the intestinal barrier, *L. monocytogenes* transfers to the spleen and liver, where it is eliminated by the T-cell-mediated immune system and Kupffer cells (Ebe *et al.*, 1999, Gregory and Liu, 2000, Pope *et al.*, 2001, Shi *et al.*, 2010, Blériot *et al.*, 2015). However, in hosts with depressed cell-mediated immunity, *L. monocytogenes* may spread into the circulatory system and cause bacteraemia. Furthermore, it may cross the blood–brain barrier or during pregnancy the placenta and, consequently, cause the

symptoms of invasive listeriosis (Mylonakis *et al.*, 1998, Mylonakis *et al.*, 2002, Bakardjiev *et al.*, 2006, Disson and Lecuit, 2012).

High-risk groups for invasive listeriosis include neonates, the elderly and individuals immunocompromised due to comorbid malignant diseases, autoimmune disorders, organ transplantation, immunosuppressive medication or pregnancy (Goulet and Marchetti, 1996, Mylonakis *et al.*, 1998, Goulet *et al.*, 2012, Silk *et al.*, 2012). Invasive listeriosis manifests most commonly as bacteraemia or septicaemia, meningitis and meningoencephalitis. Less frequently, pneumonia, peritonitis and other focal infections may occur (Ericsson *et al.*, 1997, López-Prieto *et al.*, 2000, Schlech, 2000, Swaminathan and Gerner-Smidt, 2007, Aubin *et al.*, 2016). During pregnancy, infection may lead to abortion or neonatal listeriosis, while pregnant women frequently display non-specific symptoms resembling influenza (Evans *et al.*, 1985, Mylonakis *et al.*, 2002, Vázquez-Boland *et al.*, 2017). Early-onset neonatal listeriosis derives from intrauterine infection. It can result in stillbirth or septicaemia and pneumonia in the infant within a few days after delivery. The rarer late-onset form is considered to derive from passage through the birth canal during labour. Meningitis, as the most common clinical presentation, occurs during a few weeks following delivery (Evans *et al.*, 1985, Mylonakis *et al.*, 2002, Sapuan *et al.*, 2017, Vázquez-Boland *et al.*, 2017).

In healthy individuals, *L. monocytogenes* may cause a noninvasive infection presenting as febrile gastroenteritis (Salamina *et al.*, 1996, Dalton *et al.*, 1997, Miettinen *et al.*, 1999b, Aureli *et al.*, 2000, Frye *et al.*, 2002, Ooi and Lorber, 2005, Jacks *et al.*, 2016). Sporadic cases of localised eye and cutaneous infections have also been described, which are frequently, but not exclusively, acquired through direct exposure of the site of infection instead of foodborne transmission and haematogenous dissemination (Betriu *et al.*, 2001, Regan *et al.*, 2005, Tay *et al.*, 2008, Godshall *et al.*, 2013).

Invasive listeriosis usually presents a prolonged incubation period of approximately 20 to 30 days, but this may vary widely from one up to 70 days (Linnan *et al.*, 1988, Goulet *et al.*, 2013). The minimum infectious dose of *L. monocytogenes* for either healthy or immunocompromised individuals is unclear, but the development of clinical illness appears to be dose-dependent (Ooi and Lorber, 2005). Prolonged daily intake may further increase the risk (Maijala *et al.*, 2001). In risk groups, *L. monocytogenes* levels of $>10^2$ cfu/g have caused invasive listeriosis (Ericsson *et al.*, 1997, Maijala *et al.*, 2001). In noninvasive listeriosis, the onset of symptoms is more rapid and usually takes place within a day or two after the ingestion of contaminated food, with a median of 18 to 31 hours (Salamina *et al.*, 1996, Dalton *et al.*, 1997, Miettinen *et al.*, 1999b, Aureli *et al.*, 2000, Frye *et al.*, 2002, Sim *et al.*, 2002, Carrique-Mas *et al.*, 2003). The infectious dose is usually markedly higher than that in invasive cases, approximately 10^5 to 10^9 cfu/g, although inocula down to 3×10^1 cfu/g have been detected in a potential vehicle food of noninvasive gastroenteritis (Dalton *et al.*, 1997,

Miettinen *et al.*, 1999b, Aureli *et al.*, 2000, Frye *et al.*, 2002, Sim *et al.*, 2002, Carrique-Mas *et al.*, 2003).

While *L. monocytogenes* strains isolated from food samples are most commonly serogroup 1/2 strains (Gilot *et al.*, 1996, Gianfranceschi *et al.*, 2003), the most common serotypes causing listeriosis are 4b, 1/2a and 1/2b (McLauchlin, 1990, Gilot *et al.*, 1996, Gianfranceschi *et al.*, 2003, Pontello *et al.*, 2012, Koopmans *et al.*, 2013, Prieto *et al.*, 2016): serotype 1/2a has been overrepresented in clinical cases in the Nordic countries (Lukinmaa *et al.*, 2003, Parihar *et al.*, 2008, Lopez-Valladares *et al.*, 2018), while 4b has been associated with high rates of hospitalisations and deaths particularly in North America (Pinner *et al.*, 1992, Cartwright *et al.*, 2013). Moreover, serotype 1/2a may gradually be outnumbering 4b as the most common serotype causing listeriosis worldwide, possibly due to its recurrent occurrence in increasingly favoured RTE foods (Lopez-Valladares *et al.*, 2018). However, based on putative associations between virulence and serotypes, McLauchlin (1990) suggested 4b to be more virulent than other serotypes. Ward *et al.* (2008) found by multilocus genotyping that mutations in *inlA*, leading to truncated InlA and decreased virulence, were exclusively harboured by serogroup 1/2 isolates, which may have explained the past overrepresentation of 4b isolates in listeriosis cases (Ward *et al.*, 2008). Notably, the bacteriocin LLS, beneficial for the invasion process inside host intestines, has only been recognised among a subset of lineage I strains (Cotter *et al.*, 2008, Quereda *et al.*, 2016), possibly further explaining the differences in virulence between lineages and serotypes. In the WGS era, putative virulence loci, such as the *Listeria* pathogenicity island LIPI-4, have been recognised in particularly virulent clinical clonal complexes (CC) among lineage I, which could explain their tropism for the placenta and central nervous system (Maury *et al.*, 2016, Radoshevich and Cossart, 2018).

2.2.2 Animal listeriosis

Among domestic animals, listeriosis mainly occurs in small ruminants and to a lesser extent in cattle, although nearly every domestic species is potentially susceptible to listeriosis (Nightingale *et al.*, 2004, Lecuit, 2007). Rarely, pathogenic *Listeria* spp. has been isolated in livestock and wildlife species such as swine, horses, fish and birds (Lecuit, 2007). Besides *L. monocytogenes*, *L. ivanovii* is also pathogenic in animals (Gouin *et al.*, 1994, Chand and Sadana, 1999). As in humans, a number of outbreaks along with sporadic cases have been reported in animals (Wilesmith and Gitter, 1986, Wiedmann *et al.*, 1996, Chand and Sadana, 1999, Lecuit, 2007).

The course of infection in animals is predominantly similar to that described in human listeriosis (Farber *et al.*, 1991, Lecuit, 2007). Clinical listeriosis in bovine, ovine and caprine livestock most often manifests as septicaemia, meningoencephalitis or, in gravid individuals, abortion (Wilesmith and Gitter, 1986, Campero *et al.*, 2002, Lecuit, 2007).

Septicaemic manifestation is usually seen in neonates. Rarely, ocular infections, mastitis and gastroenteritis have been reported (Lecuit, 2007). Clinically healthy animals may carry and shed the bacteria, and cattle in particular have been considered to transmit *L. monocytogenes* in the farm environment (Husu, 1990, Renterghem *et al.*, 1991, Nightingale *et al.*, 2004).

Predisposing factors for animal listeriosis include contaminated feed, deficient housing and management practices, and poor animal health. Similarly to human infections, the most common source of animal listeriosis is poorly prepared feed, with spoiled silage most often indicated as a culprit of transmission in farm animals (Wilesmith and Gitter, 1986, Husu *et al.*, 1990a, Vázquez-Boland *et al.*, 1992, Wiedmann *et al.*, 1996, Nightingale *et al.*, 2005a) Ruminants have also acquired infection through abrasions in the buccal mucosa followed by unilateral ascending neuritis along the trigeminal nerve (Braun *et al.*, 2002).

2.3 Plasmids in *Listeria* spp.

Plasmids, like other mobile genetic elements (MGEs), are capable of mediating genetic material between cells through horizontal gene transfer (HGT) (Frost *et al.*, 2005, Rankin *et al.*, 2011). HGT mechanisms include the uptake of free DNA from the environment via transformation, transduction conducted by bacteriophages, and conjugation, in which conjugative plasmids or integrative conjugative elements (ICEs) are transferred via a mating pair formation (Mpf) complex between two adjacent cells (Frost *et al.*, 2005, Thomas and Nielsen, 2005, Goessweiner-Mohr *et al.*, 2014). Natural competence, i.e. ability to uptake extracellular DNA via transformation, has not been detected in *L. monocytogenes* (Borezee *et al.*, 2000), while conjugation and transduction do occur.

The plasmid DNA transport machinery is encoded by *tra* operons harboured by the plasmid itself. For a complete conjugative apparatus of self-transmissible plasmids, the origin of transfer (*oriT*), relaxase, type IV coupling protein and the type IV secretion system (T4SS) are required. Mobilisable plasmids, in turn, utilise the T4SS of other co-resident conjugative plasmids in order to transmit via conjugation (Schröder and Lanka, 2005, Abajy *et al.*, 2007, Smillie *et al.*, 2010, Goessweiner-Mohr *et al.*, 2014).

Plasmids are most often circular, double-stranded DNA molecules, while some bacteria may harbour linear plasmids (Hayakawa *et al.*, 1979, Hinnebusch and Tilly, 1993). Plasmids control their copy number independently. They are self-replicating entities, with circular plasmids applying theta or rolling-circle replication (del Solar *et al.*, 1998, Slater *et al.*, 2008). Plasmids are classified into incompatibility groups (Inc) by replication control mechanisms and partitioning functions. Plasmids within

one Inc group cannot coexist in the same cell (Thomas and Smith, 1987, Frost *et al.*, 2005).

The presence of native plasmids in *Listeria* spp. was first described by Pérez-Díaz *et al.* (1982). Since then, a little over one-third of *L. monocytogenes* strains on average have been found to harbour plasmids, with isolation rates varying from 0% to 87% (Pérez-Díaz *et al.*, 1982, Flamm *et al.*, 1984, Fistrovici and Collins-Thompson, 1990, Lebrun *et al.*, 1992, Peterkin *et al.*, 1992, Dykes *et al.*, 1994, McLauchlin *et al.*, 1997, Margolles and de los Reyes-Gavilán, 1998, Fox *et al.*, 2016, Hingston *et al.*, 2017b). Infrequently, two nonidentical plasmids of similar or more divergent sizes have been found to coexist in a single *Listeria* spp. isolate (Kolstad *et al.*, 1991, Lebrun *et al.*, 1992, Dykes *et al.*, 1994, McLauchlin *et al.*, 1997, Hingston *et al.*, 2017b).

L. monocytogenes plasmids are more often associated with lineage II than lineage I strains (Lebrun *et al.*, 1992, McLauchlin *et al.*, 1997, Margolles and de los Reyes-Gavilán, 1998, Orsi *et al.*, 2011, Hingston *et al.*, 2017b). Within serotype 1/2a, plasmids have more often been isolated from food-related strains compared to clinical ones, while in serotype 1/2c, plasmids seem to be more abundant in clinical isolates (Lebrun *et al.*, 1992, McLauchlin *et al.*, 1997).

L. monocytogenes plasmids are mosaics but share many highly homologous fragments (Kolstad *et al.*, 1991, Canchaya *et al.*, 2010, Gilmour *et al.*, 2010). While a common replicon type has been recognised, suggesting a common ancestral origin, Kuenne *et al.* (2010) found that *Listeria* spp. plasmids can be divided into two distinct phylogenetic groups based on their *repA*-encoded replication initiation proteins. Of these, group 2 plasmids are larger and display more heterogeneity than group 1 (Kuenne *et al.*, 2010, Hingston *et al.*, 2017b). Comparative genetic analysis has revealed multiple phage defence systems to be found only among group 2 plasmids, suggesting a role for these plasmids in resisting phage infections (Kuenne *et al.*, 2010).

2.4 Two-component systems

Two-component signal transduction systems (TCSs) allow bacteria to rapidly sense and adapt to a varied range of stressors. The prototypical modular TCS consists of a periplasmic sensor histidine kinase (HK) and a cognate cytoplasmic response regulator (RR) (Fig. 1). HK protein harbours a variable N-terminal sensor (input) domain and a conserved C-terminal cytoplasmic kinase (transmitter) domain, linked by a transmembrane region with a variable number of segments spanning the cell membrane. Unlike membrane-bound HK sensors, some HKs are soluble cytoplasmic proteins, able to detect changes in the intracellular environment. The RR, in turn, is comprised of a conserved N-terminal regulatory (receiver) domain and a variable C-terminal effector (output) domain (Chang and Stewart, 1998,

Stock *et al.*, 2000). The variable domains within HK and RR respectively reflect the need to detect a vast number of specific stimuli and to generate an appropriate response by the cell (Gao and Stock, 2009).

The operation of TCSs is based on protein phosphorylation. In the direct phosphotransfer system, the sensor domain of the HK detects a stimulus, leading to ATP-dependent autophosphorylation at a specific histidine (His) residue, catalysed by the kinase domain. The cognate RR then catalyses the transfer of the phosphoryl group to its own aspartate (Asp) residue in the regulatory domain. Phosphorylation of the regulatory domain of the RR leads to activation of the effector domain and, finally, to an appropriate response to a particular stimulus. Predominantly, RRs function as DNA-binding transcription factors (West and Stock, 2001, Casino *et al.*, 2010). More complex phosphorelays may employ multiple HK and RR proteins, hybrid HKs with both His and Asp residues, and His-containing phosphotransfer proteins (HPT) that act as an intermediary in phosphoryl group transmission (Appleby *et al.*, 1996, West and Stock, 2001).




Figure 1 Schematic representation of a prototypical two-component system (A) and a phosphorelay with a hybrid histidine kinase and a His-containing phosphotransfer (B). Modified from (Mitrophanov and Groisman, 2008).

The RR phosphorylation level defines the extent of the output response. Dephosphorylation of RR may be catalysed by its own autophosphatase activity or by bifunctional HKs, limiting the active state (Lois *et al.*, 1993, Weiss *et al.*, 2002). HK autophosphorylation and phosphotransfer activity are also potential signal transfer control steps (Casino *et al.*, 2010).

Besides signalling between cognate HK and RR, signal transduction may take place between noncognate partners. In branched signalling pathways, one HK may phosphorylate multiple RRs (one-to-many) or several HKs one RR (many-to-one) (Laub and Goulian, 2007, Goulian, 2010). An example of

such a branched pathway is the chemotaxis machinery with HK CheA phosphorylating the RRs CheB and CheY (Kirby, 2009). In *Escherichia coli*, cross-talk between HK PmrB and its cognate RR PmrA, as well as noncognate QseB, augments polymyxin B resistance (Guckes *et al.*, 2017).

TCSs are abundant in eubacteria and have also been identified, although less frequently, in archaea and eucaryotes, while being absent in animals and humans (West and Stock, 2001, Wuichet *et al.*, 2010). An estimated 1% of the proteins encoded in eubacteria are comprised of TCSs (West and Stock 2001). However, the number of TCSs varies greatly between bacterial species. In *Synechocystis* sp., 80 open reading frames (ORFs) encode TCS proteins (Mizuno *et al.*, 1996), while the *Mycoplasma genitalium* genome harbours none (Mizuno, 1998). In the sequenced genome of *L. monocytogenes* EGD-e, 16 TCSs have been identified (Glaser *et al.*, 2001). Of these, 15 are complete signalling systems, while one, DegU, is an orphan RR lacking a cognate genetically linked HK (Williams *et al.*, 2005a).

2.5 Stress responses of *L. monocytogenes*

L. monocytogenes surmounts many diverse and harsh stress conditions both in the environment and inside host organisms. While it displays optimal growth at temperatures from 30 °C to 37 °C (McLauchlin and Rees, 2015), it is able to both survive and grow over a wide temperature range from <0 °C to 45 °C (Junttila *et al.*, 1988, Walker *et al.*, 1990, Hudson *et al.*, 1994, McLauchlin and Rees, 2015). In a vacuum, growth has been detected down to -1.5 °C (Hudson *et al.*, 1994). However, *L. monocytogenes* does not survive pasteurisation treatment at 60 °C for 30 minutes (McLauchlin and Rees, 2015) nor at 72 °C for 15 seconds (Bradshaw *et al.*, 1985). *L. monocytogenes* grows in a water activity down to 0.9 (Nolan *et al.*, 1992) and in salt concentrations of up to 10% (w/v) (Simon, 1956, McLauchlin and Rees, 2015). Combined with low temperature (5 °C), survival has been detected in 18% NaCl (Cole *et al.*, 1990). *L. monocytogenes* grows at a pH as high as 9.6 (Simon, 1956, Gray and Killinger, 1966). Under acidic conditions, growth has been reported at pH 4.4 (George *et al.*, 1988) and, combined with low temperature (5 °C), survival at a pH down to 4.2 has been observed (Cole *et al.*, 1990). *L. monocytogenes* is a facultative anaerobic bacterium and thus able to grow under both aerobic and anaerobic conditions (Jydegaard-Axelsen *et al.*, 2005, McLauchlin and Rees, 2015). Consequently, refrigeration, vacuum packaging and preservation measures relying on a high salt concentration or extreme pH values are not sufficient to prevent the survival and growth of *L. monocytogenes*. If not killed by extreme stress, *L. monocytogenes* may either adjust to the stress conditions or, by means of flagellation, escape the harmful exposure by moving towards a more benign environment (Galsworthy *et al.*, 1990, Wright *et al.*, 2014).

2.5.1 General stress responses and stress tolerance mechanisms

Low temperature

L. monocytogenes is a psychrotrophic organism able to resist a cold milieu. Environmental bacteria may encounter low temperatures in their natural habitat, while foodborne pathogens also experience cold stress during the storage and refrigeration of vehicle foods. Temperature downshift has multiple direct and indirect impacts on bacteria, such as a loss of membrane fluidity and molecule solubility, slow nutrient transport and diffusion, increased RNA and DNA secondary structure stability, a decelerated lag phase and growth rate, and a restricted function of enzymes and other proteins, as well as direct protein damage (Beales, 2004, Tasara and Stephan, 2006, Chan and Wiedmann, 2009, Soni *et al.*, 2011).

L. monocytogenes is particularly capable of modulating its membrane lipids, which consist of an unusually high proportion of odd-numbered, iso and anteiso, branched-chain fatty acids (FAs) (Raines *et al.*, 1968, Annous *et al.*, 1997). In response to a decrease in temperature, multiple alterations in membrane lipid structure take place to allow the cell to maintain membrane fluidity and, consequently, to uphold vital functions in regard to enzyme activity and cross-membrane transportation (Beales, 2004, Tasara and Stephan, 2006, Chan and Wiedmann, 2009, Soni *et al.*, 2011). One of the key changes is the increased relative proportion of C_{15:0} FAs and, in turn, a decrease in C_{17:0} (Püttmann *et al.*, 1993, Russell *et al.*, 1995, Annous *et al.*, 1997, Mastronicolis *et al.*, 1998). As C_{15:0} outnumbers C_{17:0}, the proportion of shorter FAs rises, which decreases the interaction between carbon chains, lowers their melting temperature, and thereby retains membrane fluidity (Russell *et al.*, 1995, Beales, 2004). In addition, unsaturation due to a slight increase in C_{18:1} supports the fluid structure (Russell *et al.*, 1995). Annous *et al.* (1997) stated the shift from iso to anteiso branching, particularly to anteiso-C_{15:0}, to be of importance in the growth of *L. monocytogenes* at cold temperatures.

In addition to physical changes within the cell membrane, *L. monocytogenes* is able to actively transport and accumulate compatible solutes from its environment into the cell and harness them as cryoprotectants without inhibiting other vital cellular functions (Wilkins *et al.*, 1972, Tasara and Stephan, 2006, Chan and Wiedmann, 2009). Upon cold stress, the expression of the transporter-encoding genes and thereby the uptake of carnitine and glycine betaine is induced (Angelidis and Smith, 2003, Wemekamp-Kamphuis *et al.*, 2004a, Miladi *et al.*, 2017). Carnitine is mainly transported into the cell by the OpuC transporter and glycine betaine by Gbu, and to a lesser extent by BetL (Angelidis and Smith, 2003). These compatible solutes markedly stimulate the growth of cold-stressed *L. monocytogenes* cells (Beumer *et al.*, 1994, Ko *et al.*, 1994, Wemekamp-Kamphuis *et al.*, 2004a).

One of the key responses of psychrotrophic bacteria to low temperatures is the recruitment of cold shock proteins (Csps), synthesized upon an abrupt

decrease in temperature, and cold acclimation proteins (Caps), produced in balanced growth at cold temperatures (Hébraud and Potier, 1999). Bayles *et al.* (1996) recognised twelve Csps whose synthesis was induced in *L. monocytogenes* 10403S following a temperature downshift from 37 °C to 5 °C. Four of these were also identified as Caps (Bayles *et al.*, 1996). At 4 °C, the production of 32 proteins designated as Csps was induced ≥ 3 -fold in *L. monocytogenes* EGD (Phan-Thanh and Gormon, 1995). Cold shock protein CspA, CspB and CspD-encoding genes were upregulated at 4 °C and, of these proteins, CspA proved to be required for the growth of *L. monocytogenes* at both 4 °C and 10 °C (Schmid *et al.*, 2009). While a core set of 22 Cap-encoding genes was induced from the early lag to the late stationary phase, the cold-stress regulon of *L. monocytogenes* has been shown to be highly variable across different growth phases, implicating a growth phase-dependent response to low temperatures (Hingston *et al.*, 2017a).

Besides Csps and Caps, induction of the expression and production of multiple genes and proteins, respectively, has been reported in *L. monocytogenes* upon temperature downshift, indicating diverse mechanisms contributing to cold tolerance (Liu *et al.*, 2002, Chan *et al.*, 2007b, Cacace *et al.*, 2010). Liu *et al.* (2002) detected 11 genes whose expression was induced at 10 °C. Among these, ferritin-like cold-adaptive response protein, encoded by *flp*, has been named both as Csp and Cap (Bayles *et al.*, 1996, Hébraud and Guzzo, 2000). Other induced genes encode the general stress response proteins ClpB and ClpP and proteins involved in regulatory adaptive responses, alterations in the cell surface, and amino acid and degradative metabolism (Liu *et al.*, 2002). Chan *et al.* (2007b) reported the upregulation of genes that encode lipid and carbohydrate metabolism, motility, transcription and translation-related proteins at 4 °C, further illustrating the versatility of cold stress responses. Altogether 105 and 170 genes were ≥ 2 -fold transcribed in log- and stationary-phase cells, respectively (Chan *et al.*, 2007b). Furthermore, several TCSs (reviewed in chapter 2.5.4) play an important role in sensing low temperature and mediating the cold stress response (Chan *et al.*, 2007b, Chan *et al.*, 2008). DEAD-box RNA helicases have a pivotal function in RNA metabolism and translation initiation (Chan and Wiedmann, 2009, Linder and Jankowsky, 2011). The DEAD-box proteins Lmo0866, Lmo1450 and Lmo1722 play a role in the growth and motility of *L. monocytogenes* at low temperature (Chan *et al.*, 2007b, Markkula *et al.*, 2012b). Regarding motility and flagellation, flagellin-encoding *flaA* presented elevated transcript levels at 10 °C (Liu *et al.*, 2002). The *flhA* and *motA* genes encoding the temperature-sensitive flagella synthesis proteins FlhA and MotA, respectively, contribute to the growth and motility of *L. monocytogenes* at 3 °C (Mattila *et al.*, 2011).

Alternative stress sigma factor σ^B , encoded by *sigB*, coordinates many stress responses in *L. monocytogenes* (Becker *et al.*, 1998, Becker *et al.*, 2000, Kazmierczak *et al.*, 2003, Raengradub *et al.*, 2008) and directly and

positively regulates over 140 genes (Raengpradub *et al.*, 2008). The activity of σ^B is induced in stationary-phase cells upon temperature downshift (Becker *et al.*, 2000). Deletion of *sigB* renders the cell unable to accumulate betaine and carnitine (Becker *et al.*, 2000). Also, carnitine transporter-encoding *opuCA* is at least partly σ^B -dependent (Chan *et al.*, 2007a), further suggesting a role for σ^B in harnessing compatible solutes. On the other hand, the transcription of *opuCA* as well as of cold stress genes *fri* and *oppA* is independent of σ^B during cold stress at 4 °C (Chan *et al.*, 2007a). Although supporting the cold stress response, it was further demonstrated that σ^B is not unequivocally essential for the growth of *L. monocytogenes* at 4 °C (Chan *et al.*, 2007a).

High temperature

Although a psychrotrophic organism, *L. monocytogenes* tolerates fairly high temperatures, particularly as compared to other nonspore-forming foodborne pathogens, and may survive mild heat treatments (Doyle *et al.*, 1987, Doyle *et al.*, 2001, van der Veen *et al.*, 2007). Foodborne pathogens frequently encounter high temperatures during food processing. When subjected to heat reaching lethal temperatures, the DNA structure, ribosomes and ribosomal RNA degrade, the cytoplasmic membrane is damaged leading to the leakage of cellular components, and protein denaturation is initiated, resulting in metabolic malfunctions and reduced enzyme activity (Russell, 2003, Soni *et al.*, 2011). Thus, to maintain cellular mechanisms and to survive high temperatures, cells need to preserve nucleic acid functions, repair membranes and produce new proteins while removing damaged ones (van der Veen *et al.*, 2009, Soni *et al.*, 2011).

To overcome heat shock, *L. monocytogenes* harnesses two specific class I and III heat shock response mechanisms and a class II general stress response (van der Veen *et al.*, 2007). Class I contains an operon of classical chaperon-encoding *hrcA*, *grpE*, *dnaK*, *dnaJ*, *groEL* and *groES* genes (Hanawa *et al.*, 2000). The operon is negatively regulated by HrcA, which binds at a conserved inverted repeat element (Zuber and Schumann, 1994, Hanawa *et al.*, 2000, Hu *et al.*, 2007a). Class III heat shock proteins (Hsps) are comprised of Clp ATPases that function as ATP-dependent proteases, breaking down damaged and misfolded proteins, and as chaperons, aiding in protein folding (Wawrzynow *et al.*, 1996). Class III genes are negatively regulated by the transcriptional repressor CtsR (Derré *et al.*, 1999, Nair *et al.*, 2000, Hu *et al.*, 2007b). Besides stress tolerance, the chromosomally encoded ClpC, ClpP and ClpE class III Hsps are involved in the virulence of *L. monocytogenes* (Nair *et al.*, 2000, Rouquette *et al.*, 1996). Class II, positively regulated by σ^B , is comprised of diverse genes including listeriolysin positive regulatory protein-encoding *prfA* and general stress protein-encoding *ctc* (Benson and Haldenwang, 1993, van der Veen *et al.*, 2007). Despite the positive regulation of class II by σ^B , heat stress resistance

in *L. monocytogenes* seems to be at least partly independent of σ^B alone (Ferreira *et al.*, 2001).

Marked overlap has been detected within HrcA, CtsR and σ^B regulons, demonstrating a large regulatory network contributing to heat stress tolerance in *L. monocytogenes* (Hu *et al.*, 2007a, Hu *et al.*, 2007b, Raengpradub *et al.*, 2008, Chaturongakul *et al.*, 2011). Five genes were identified to be coregulated by all three regulators (Hu *et al.*, 2007a), and σ^B was detected to coregulate 25 and 21 genes with CtsR and HrcA, respectively (Chaturongakul *et al.*, 2011). In addition, σ^B contributes to the positive regulation of HrcA-encoding *hrcA* (Hu *et al.*, 2007a).

pH stress

L. monocytogenes encounters low pH throughout the food chain and within hosts: in silage, in fermented and other low-pH foods, during gastric passage in a host and inside macrophages upon phagocytosis (Beauregard *et al.*, 1997, Ryser *et al.*, 1997, Cotter *et al.*, 2001a, Cotter *et al.*, 2001b, Cotter and Hill, 2003). Alkaline conditions are commonly encountered in contact with detergents, but also upon pancreatic secretion in the host small intestine (Giotis *et al.*, 2007a, Soni *et al.*, 2011).

Acid stress reduces the growth rate and yield (Cheroutre-Vialette *et al.*, 1998, Bowman *et al.*, 2012). Weak organic acids are more harmful to bacterial cells compared to strong acids. They are membrane-permeable in their undissociated form and, once intracellular, weak acids dissociate, release a proton and directly acidify the intracellular space. Strong acids nevertheless lower the external pH and damage enzymes and, as proton permeability is induced due to a high gradient, strong acids may also ultimately acidify the cytoplasm, albeit less drastically than weak acids (Phan-Thanh *et al.*, 2000, Cotter and Hill, 2003, Beales, 2004). Alkaline conditions, in turn, disturb cell division and growth, elongate listerial cells and damage cell membranes, leading to the leakage of intracellular components (Zilberstein *et al.*, 1984, Sampathkumar *et al.*, 2003, Giotis *et al.*, 2007a).

Upon pH stress, bacterial cells need to maintain pH homeostasis: proton extrusion under acidic and inward transport of protons under alkaline conditions (Zilberstein *et al.*, 1984, Shabala *et al.*, 2002, Krulwich *et al.*, 2011). Transient failure of pH control may precede the reestablishment of the intracellular pH (Zilberstein *et al.*, 1984). Besides alterations in transport and metabolic functions, surmounting both high and low pH involves structural changes in cell composition (Krulwich *et al.*, 2011). In *L. monocytogenes*, the proportion of anteiso FAs increases at an alkaline pH, while the opposite takes place under acidic conditions (Giotis *et al.*, 2007b). The proportion of iso and anteiso forms is putatively of more importance in pH adaptation than changes in the saturation of FAs (Giotis *et al.*, 2007b).

Membrane transport functions are pivotal in maintaining intracellular pH homeostasis in *L. monocytogenes*, particularly under acid stress (Cotter and

Hill, 2003). As a facultative anaerobic bacterium, *L. monocytogenes* may utilise both the active transport of protons, through a H⁺-linked electron transport system in the aerobic respiratory chain, and transport via H⁺-ATPase molecules, including F₀F₁-ATPase, which use energy anaerobically derived from ATP hydrolysis (Cotter *et al.*, 2000, Shabala *et al.*, 2002). To maintain the intracellular pH under acidic conditions, *L. monocytogenes* employs the glutamate decarboxylase system (GAD), encoded by *gadA*, *gadB* and *gadC* (Cotter *et al.*, 2001a, Cotter *et al.*, 2001b). Another GAD and an additional antiporter is encoded by *gadD* and *gadE*, respectively (Wemekamp-Kamphuis *et al.*, 2004b). GAD consumes intracytoplasmic protons by producing γ -aminobutyrate (GABA) from extracellular glutamate, hence elevating the cytoplasmic pH (Small and Waterman, 1998). GABA is then released, and the substitution of glutamate with GABA outside the cell will also slightly alkalise the extracellular environment (Small and Waterman, 1998). In addition to GAD, *L. monocytogenes* harbours a functional arginine deiminase system (ADI), encoded by *arcA*, *arcB*, *arcC* and *arcD*, also playing a role in the acid tolerance of *L. monocytogenes* (Ryan *et al.*, 2009). The σ^B mutant displays impaired resistance of *L. monocytogenes* to acid stress, indicating the importance of σ^B in surmounting acidic conditions (Wiedmann *et al.*, 1998, Ferreira *et al.*, 2001, Wemekamp-Kamphuis *et al.*, 2004b). Furthermore, the expression of the *gadCB* operon and ADI system is regulated by σ^B (Kazmierczak *et al.*, 2003, Wemekamp-Kamphuis *et al.*, 2004b, Ryan *et al.*, 2009). During acid adaptation at pH 5.5 and acid stress at pH 3.5, the synthesis of 37 and 47 proteins, respectively, was induced ≥ 2 -fold as compared to a neutral pH of 7.2 (Phan-Thanh and Mahouin, 1999). The number of proteins may reflect the increasing effort to maintain intracytoplasmic pH homeostasis under harshening acid stress (Phan-Thanh and Mahouin, 1999).

While the exact mechanisms allowing *L. monocytogenes* to tolerate high pH levels remain fairly obscure, upon alkali stress, *L. monocytogenes* appears to strive at diminishing excess alkalinisation and energy outlay whilst mobilising carbon sources (Giotis *et al.*, 2008b). Transcriptional profiles were found to differ between abrupt alkali shock and alkali adaptation (Giotis *et al.*, 2010). Giotis *et al.* (2008b) found the synthesis of 45 proteins to be induced ≥ 1.5 -fold upon alkali adaptation, including DnaK and GroEL (Giotis *et al.*, 2008b). Alkali stress seems to mimic phosphate starvation in *L. monocytogenes*, and phosphate uptake and utilisation appears to be revised upon alkali exposure (Giotis *et al.*, 2008b).

Osmotic stress

Foodborne pathogens frequently encounter osmotic stress due to high salt concentrations, as sodium chloride is commonly used as a food preservative and flavour additive. Desiccation and accumulating osmotically active substances leads to a decrease in water activity (a_w), the water quantity available to bacterial cells (Gutierrez *et al.*, 1995, Beales, 2004).

Consequently, the cytoplasmic volume decreases and cells begin to lose their intracellular turgor pressure, cellular functions deteriorate, and bacterial growth decelerates (Csonka, 1989, Cheroutre-Vialette *et al.*, 1998, Beales, 2004). Bacteria resist these changes through osmoadaptation (Galinski, 1995, Gutierrez *et al.*, 1995).

Under osmotic stress, listerial cells try to maintain the proper lipid bilayer of their cell membrane by increasing the proportion of anionic lipids, particularly diphosphatidylglycerol (Russell *et al.*, 1995). Furthermore, cells restore their internal turgor pressure by accumulating from their environment the same compatible solutes, glycine betaine and carnitine, as under cold conditions (Csonka, 1989, Patchett *et al.*, 1992, Gutierrez *et al.*, 1995, Bayles and Wilkinson, 2000). The compatible solutes stabilise intracellular enzymes (Russell *et al.*, 1995) and potentially hydrate the protein surfaces, stabilising proteins and preventing their misfolding (Kempf and Bremer, 1998). They also induce the growth of *L. monocytogenes* upon osmotic stress and a_w decrease (Beumer *et al.*, 1994, Bayles and Wilkinson, 2000). In 4% NaCl, a 200-fold increase in the transportation rate of glycine betaine has been reported (Ko *et al.*, 1994), and associated loci are upregulated even under milder osmotic stress (1.2% NaCl) (Bae *et al.*, 2012). Proline also serves as an osmolyte in *L. monocytogenes* (Sleator *et al.*, 2001). Besides free amino acids, peptides aid in maintaining the cell turgor and thereby contribute to osmotolerance (Amezaga *et al.*, 1995). While proline only mediates osmotolerance at high concentrations (10 mM) (Beumer *et al.*, 1994), as a peptide, it confers osmoprotection even as a mild (1 mM) solution (Amezaga *et al.*, 1995).

Upon osmotic stress, Duché *et al.* (2002) detected six rapidly induced (<30 min) salt shock proteins (Ssps) and eleven salt acclimation proteins (Saps) that are more slowly induced (60–90 min) but maintain the induced level for a prolonged period (Duché *et al.*, 2002a). The Ssps included heat shock protein DnaK, general stress response protein Ctc, and proteins involved in general metabolism (Duché *et al.*, 2002a, Duché *et al.*, 2002b). Glycine betaine transporter Gbu was induced later during the adaptational phase (Duché *et al.*, 2002a). Ctc is putatively involved in osmotolerance, regardless of osmoprotectants, as glycine betaine is able to ameliorate the detrimental osmotic effects similarly in both the wild type and *ctc* mutant (Gardan *et al.*, 2003).

σ^B also regulates multiple stress response genes in the osmotolerance of *L. monocytogenes* (Becker *et al.*, 1998, Kazmierczak *et al.*, 2003, Abram *et al.*, 2008, Ribeiro *et al.*, 2014). The osmotic induction of carnitine transporter-encoding *opuC* is σ^B -dependent (Fraser *et al.*, 2003), and the absence of σ^B also diminishes the uptake of betaine, suggesting σ^B to play a role in osmolyte transportation upon osmotic stress in *L. monocytogenes* (Becker *et al.*, 1998). In response to salt stress, σ^B represses the transcription of flagellar protein-encoding genes, indicating an indirect impact on motility and chemotaxis (Raengpradub *et al.*, 2008). Although chemotaxis allows

bacterial movement to more favourable surroundings, flagellar synthesis is an energy-consuming process, and downregulation of flagellar genes may be a means for bacterial cells to allot energy to more crucial mechanisms needed to surmount osmotic stress (Hingston *et al.*, 2015). Additionally, genes encoding RNA helicases (Markkula *et al.*, 2012a) and TCSs (Kallipolitis and Ingmer, 2001) are involved in the growth of *L. monocytogenes* under osmotic stress.

Ethanol stress

Foodborne bacteria frequently encounter high ethanol concentrations during sanitation and preservation processes and in some food products. While Gram-negative bacteria seem to be fairly susceptible to ethanol, Gram-positives and particularly lactic acid bacteria survive markedly high ethanol concentrations (Ingram, 1976, Gold *et al.*, 1992, Silveira *et al.*, 2004, van Bokhorst-van de Veen *et al.*, 2011). *Lactobacilli* can grow in 10% ethanol and intermediate growth of some strains has even been observed in 16% (Gold *et al.*, 1992). In *L. monocytogenes*, a 5% ethanol concentration seems to inhibit growth (Oh and Marshall, 1993). Ethanol may also augment the lethal effects of concomitantly administered acids and salt on *L. monocytogenes*, possibly associated with cell membrane damage (Barker and Park, 2001).

Ethanol increases membranous permeability in bacterial cells, both by altering the aqueous environment and by impairing the hydrophobic membrane core barrier. Being freely permeable through the cell membrane, ethanol affects not only enzymes on cell surfaces and within membranes but also in the cytoplasm (Ingram, 1990). Ethanol hampers protein folding and macromolecular interactions, consequently inhibiting membrane protein synthesis and decelerating cell growth (Ingram, 1990, Seydlová *et al.*, 2012). Growth inhibition also emanates from the fluidising effect of ethanol on lipid bilayers, which may further lead to a compensatory increase in membrane rigidity (Dombek and Ingram, 1984, Huffer *et al.*, 2011, Seydlová *et al.*, 2012). Commonly, an increase in saturated and unbranched FAs within the cell membrane occurs upon ethanol stress (Huffer *et al.*, 2011). In *Bacillus subtilis*, ethanol exposure and adaptation leads to the recruitment of molecular chaperones in order to restore phospholipid synthesis and membrane fluidity (Seydlová *et al.*, 2012).

The impacts of ethanol on microbial cells are thought to derive from mechanisms affecting hydrophobic associations and from the colligative impact rather than via specific receptors (Ingram, 1976, Ingram, 1990). However, it has been shown that ethanol induces the termination of Rho-dependent transcription in *E. coli* and inhibits the mRNA synthesis and translation of nonstart AUG codons, indicating a significant impact on the transcriptional and translational functions upon ethanol stress (Haft *et al.*, 2014). Haft *et al.* (2014) further suggested that alterations in *E. coli* membranes upon ethanol exposure might partly be due to ethanol directly affecting ribosomes and RNA polymerases.

In addition to genes encoding proteins directly applied to coping with high ethanol concentrations, such as alcohol and aldehyde dehydrogenases YqhD and AldB, respectively (Soufi *et al.*, 2015), many general stress response genes are induced upon ethanol stress. In *B. subtilis*, a promoter of extracytoplasmic function sigma factor σ^M was induced under ethanol stress (Thackray and Moir, 2003). In *L. monocytogenes*, RNA helicase Lmo0866 and RNA-binding protein Hfq have been shown to play roles in ethanol tolerance (Christiansen *et al.*, 2004, Markkula *et al.*, 2012a). The expression of *hfq* was, furthermore, found to be σ^B -dependent upon ethanol stress (Christiansen *et al.*, 2004).

Oxidative stress

L. monocytogenes cells experience oxidative stress both in the external environment and during intracellular host infection (Beckerman *et al.*, 1993, Gorski *et al.*, 2008, Soni *et al.*, 2011). Reactive oxygen species (ROS), such as hydrogen peroxide (H_2O_2), are metabolic by-products but also derive from external chemical processes or are released by competing microorganisms as a defence mechanism (Mishra and Imlay, 2012, Imlay, 2015a, Imlay, 2015b). As very reactive agents, ROS damage enzymes, other proteins and nucleic acids, and are harmful to cell lipid membranes. Consequently, ROS sensing and detoxification and the induction of cellular repair mechanisms are pivotal in surmounting oxidative stress (Cabiscol *et al.*, 2000, Imlay, 2002, Lee and Helmann, 2006, Mishra and Imlay, 2012, Imlay, 2015a).

The predominant regulator of the oxidative stress response in Gram-positive bacteria is PerR, a repressor of the oxidative response gene regulon (Chen *et al.*, 1995, Imlay, 2015b). In *B. subtilis*, PerR regulates the expression of genes for catalase KatA, alkyl hydroperoxide reductase AhpCF, the heme synthesis operon HemAXCDBL, the zinc uptake system ZosA, the ferric uptake repressor Fur, and protective DNA-binding Dps encoded by *mrgA* (Chen *et al.*, 1995, Helmann *et al.*, 2003). PerR and Dps-like Fri also facilitate the oxidative stress response in *L. monocytogenes* (Dussurget *et al.*, 2005, Olsen *et al.*, 2005, Rea *et al.*, 2005). The PerR regulon in *L. monocytogenes* includes *fri*, *fur*, *hemA*, *kat*, *trxB*, *lmo0641* and *lmo1604* (Rea *et al.*, 2005). Of these, *lmo0641* is the closest homolog of *zosA* in *B. subtilis* PerR-regulated genes, while *lmo1604* displays 32% identity with *ahpC*, and *fri* 35% identity with *mrgA* (Rea *et al.*, 2005), indicating a similar regulon to that described in *B. subtilis*. PerR senses peroxides by metal-catalysed oxidation (Lee and Helmann, 2006). It responds to low levels of H_2O_2 when the metal-binding site of PerR is occupied by Fe(II) while being less sensitive to H_2O_2 when occupied by Mn(II) (Herbig and Helmann, 2001). Although PerR is a direct sensor of H_2O_2 , *perR* undergoes little induction by H_2O_2 (Fuangthong *et al.*, 2002). However, apart from *perR* and *fur*, other PerR regulon genes are induced by H_2O_2 (Fuangthong *et al.*, 2002).

Mutational analyses have revealed roles for superoxide dismutase, σ^B , and RNA helicase Lmo1450 in the viability of *L. monocytogenes* upon oxidative

stress (Ferreira *et al.*, 2001, Archambaud *et al.*, 2006, Markkula *et al.*, 2012a). Under cold conditions, elevated levels of transcripts and the induction of proteins involved in protection against oxidative stress, such as thioredoxin reductase-encoding *trxB*, ferritin-like *flp*, superoxide dismutase and catalase, have been detected, suggesting *L. monocytogenes* to undergo concomitant oxidative stress at low temperatures (Liu *et al.*, 2002, Cacace *et al.*, 2010).

Benzalkonium chloride

Benzalkonium chloride (BC) belongs to the group of quaternary ammonium compounds (QAC), cationic surfactants widely used as detergents for disinfection on both organic and abiotic surfaces (McDonnell and Russell, 1999). The antimicrobial attributes of QACs are proposed to be based on the generalised disruption of cytoplasmic cell membranes, followed by a loss of structural integrity, degradation of proteins and nucleic acids and leakage of intracellular matrix, as well as cell wall lysis (Denyer, 1995, McDonnell and Russell, 1999). BC has also been shown to inhibit biofilm formation (Houari and Di Martino, 2007).

Repeated exposure to sublethal concentrations of detergents and disinfectants may result in emerging resistance of bacterial strains to these compounds (Lundén *et al.*, 2003a, Houari and Di Martino, 2007). Concerningly, BC-resistant (BC^r) strains of *L. monocytogenes* have been isolated from both food-processing environments (Mullapudi *et al.*, 2008) and foodstuffs (Soumet *et al.*, 2005), as well as clinical cases (Elhanafi *et al.*, 2010). Approximately 10% to 46% of *L. monocytogenes* isolates from foodstuffs and processing environments have been reported as BC-resistant (Aase *et al.*, 2000, Mullapudi *et al.*, 2008). However, it should be noted that the definition of BC resistance varies between studies. For instance, Aase *et al.* (2000) and Soumet *et al.* (2005) considered the strains with a minimal inhibitory concentration (MIC) of BC of more than 4 µg/ml and 7.5 µg/ml, respectively, to be BC^r, while others have estimated a BC resistance threshold of ≥10 µg/ml (Mullapudi *et al.*, 2008, Dutta *et al.*, 2013, Jiang *et al.*, 2016).

Genes known to confer BC resistance in *L. monocytogenes* include the efflux pump systems-encoding *bcrABC* (Elhanafi *et al.*, 2010), *emrE* (Gilmour *et al.*, 2010, Kovacevic *et al.*, 2016), and *qacH* on the transposon Tn6188 (Müller *et al.*, 2013). While *L. monocytogenes* strains harbour BC resistance-mediating genes, such as *bcrABC*, on the chromosome (Dutta *et al.*, 2013), plasmid-borne *bcrABC* has also been detected (Elhanafi *et al.*, 2010, Dutta *et al.*, 2013, Jiang *et al.*, 2016), indicating that BC resistance may be efficiently acquired and spread via horizontal gene transfer. The efflux pump MdrL-encoding *mdrL*, found to be both chromosomal and plasmid-borne (Romanova *et al.*, 2002), has been proposed at least with a partial role in the adaption of *L. monocytogenes* to BC (Mereghetti *et al.*, 2000, Romanova *et al.*, 2006). Previously, the chromosomal MdrL, negatively regulated by LadR, was shown to play a role in the survival of

L. monocytogenes at both lethal and sublethal BC concentrations (Jiang *et al.*, 2019).

Cross-adaptation

Exposure to certain stress conditions may enhance the tolerance of *L. monocytogenes* to subsequent exposure to the same or different stressors (Lou and Yousef, 1997, Lundén *et al.*, 2003a, Skandamis *et al.*, 2008, Soni *et al.*, 2011). Cross-protection has potentially grave implications for food safety, as multiple stress conditions are often concurrently used in food processing to prevent bacterial growth. Under many stresses, preadaptation to sublethal stress increases resistance to more severe conditions of the same media. Acid adaptation has been widely shown to augment resistance in *L. monocytogenes* to subsequent exposure to a lethal acidic pH (Kroll and Patchett, 1992, Hill *et al.*, 1995, O'Driscoll *et al.*, 1996, Lou and Yousef, 1997, Phan-Thanh *et al.*, 2000). Alkali-adapted cells, in turn, are more resistant to harsher alkaline stress (Giotis *et al.*, 2008a). Adaptational augmentation in resistance has also been demonstrated under ethanol, oxidative (Lou and Yousef, 1997, Ferreira *et al.*, 2001) and heat stresses (Shen *et al.*, 2014).

Reciprocal cross-protection under different stress conditions has been described in *L. monocytogenes*, as osmoadaptation has been found to increase acid resistance (Faleiro *et al.*, 2003), while acid tolerance, in turn, provides cross-protection against osmotic shock (Phan-Thanh *et al.*, 2000, Faleiro *et al.*, 2003). Similarly, heat-adapted cells are more resistant to acid shock (Phan-Thanh *et al.*, 2000), while acid adaptation leads to increased heat resistance (Farber and Pagotto, 1992, Phan-Thanh *et al.*, 2000).

Lou and Yousef (1997) revealed a complicated net of adaptational responses and cross-protection in *L. monocytogenes*. Resistance to lethal oxidative stress was enhanced by sublethal exposure to ethanol, acid, osmotic and heat stress. Additionally, survival under ethanol stress was increased by preceding acid and heat exposure, acid resistance improved following ethanol adaptation, and resistance to osmotic stress increased following preadaptation to ethanol and heat (Lou and Yousef, 1997). In addition to low temperature treatment providing cross-protection against mild osmotic stress, a prolonged duration of cold preadaptation was found to progressively enhance the survival of *L. monocytogenes* under subsequent osmotic stress (Pittman *et al.*, 2014). Furthermore, acid- as well as alkali-adapted cells display enhanced resistance to heat, osmotic and alcohol stresses (Phan-Thanh *et al.*, 2000, Taormina and Beuchat, 2001, Giotis *et al.*, 2008a).

Extensive cross-protection under various stresses may be explained by the recruitment of many of the same response mechanisms and induction of the same proteins under different conditions. *L. monocytogenes* employs versatile genetic mechanisms and molecular determinants to comprehensively contend with stress conditions and to persist in and adapt to niche-specific environments. Particular stress survival islets (SSI) have been recognised to play roles under multiple suboptimal conditions: a self-

regulating five-gene (*lmo0444–lmo0448*) stress survival islet, SSI-1, contributes to the growth of *L. monocytogenes* under acid and osmotic stresses (Ryan *et al.*, 2010), while SSI-2, mainly harboured by sequence type 121 (ST121) *L. monocytogenes* strains, has a role under alkaline and oxidative stresses (Harter *et al.*, 2017). Survival under harsh stresses is energy-consuming, and proteins involved in the ATP synthase chain and oxidative phosphorylation are thus commonly induced upon stress adaptation (Pittman *et al.*, 2014). Bergholz *et al.* (2012) found 888 genes with temperature-dependent alterations in transcript levels during osmotic adaptation. Based on proteomic analysis, Pittman *et al.* (2014) identified 299 differentially expressed proteins associated with the osmotic response of cold-adapted cells, pinpointing proteins putatively involved in cross-protection. Amongst others, these included cell envelope proteins and proteins involved in metabolic pathways (Pittman *et al.*, 2014). Regarding cellular processes, compatible solutes are needed for viability under both cold and osmotic stress (Bayles and Wilkinson, 2000). The alternative stress sigma factor σ^B and σ^B -regulated genes also play roles under multiple stresses (Bergholz *et al.*, 2012). The expression of SSI-1 genes was shown to be regulated by σ^B (Ryan *et al.*, 2010). Besides cold tolerance, cold shock proteins CspA, CspB and CspD contribute to some extent to osmotic and oxidative stress tolerance (Schmid *et al.*, 2009, Loepfe *et al.*, 2010). Many heat stress proteins mediate tolerance to other stresses, such as HtrA to osmotic, acid and oxidative stress (Wonderling *et al.*, 2004, Stack *et al.*, 2005, Wilson *et al.*, 2006) and DnaK and GroEL to alkaline stress (Giotis *et al.*, 2008b). Phan-Thanh and Gormon (1995) also found six proteins to be induced by both heat and cold shock.

2.5.2 Strain diversity in stress responses

L. monocytogenes strains do not form a homogeneous population but vary in their phenotypic characteristics (Abee *et al.*, 2016, Grad and Fortune, 2016). Certain lineages and serotypes display particular resistance or sensitivity to stresses frequently encountered along the food chain. Salt stress tolerance phenotypes vary according to lineage, as lineage I strains and III grow faster under osmotic stress than those of lineage II (Bergholz *et al.*, 2010, Ribeiro and Destro, 2014). Higher transcript levels of σ^B -dependent *gadD* and CtsR-dependent *clpB* were found in lineage I strains as compared to lineage II strains post-saline shock (Ringus *et al.*, 2012). Serotype 4b isolates have proven to be more heat-sensitive than other serotypes (Lianou *et al.*, 2006), while presenting a higher tolerance to low pH and high salt concentrations (van der Veen *et al.*, 2008). However, under heat and acid stresses, death rates have also been shown to vary markedly between strains (Mackey *et al.*, 1990, Lianou *et al.*, 2006). In the next-generation sequencing (NGS) era, serotypes 1/2a and 1/2b have presented on average more tolerant phenotypes under cold conditions than 4b and 1/2c (Hingston *et al.*, 2017b). Strain-

dependent behaviour is also seen in biofilm formation and in the growth of *L. monocytogenes* at low temperatures, low a_w and in vacuum packaging (Barbosa *et al.*, 1994, Barbosa *et al.*, 1995, Begot *et al.*, 1997, Nilsson *et al.*, 2011, Cordero *et al.*, 2016). Differences in stress tolerance have also been associated with the isolation sources of *L. monocytogenes* isolates. Clinical strains seem to display greater resistance to acid and osmotic stresses than environmental and food-related strains (Dykes and Moorhead, 2000, Vialette *et al.*, 2003).

Besides strain variability, population heterogeneity, i.e. variability between single cells of a strain or in a population, describes the diverse stress tolerance of pathogenic bacteria and may have further implications for food safety (Lianou and Koutsoumanis, 2013, Abee *et al.*, 2016, den Besten *et al.*, 2017). Heterogeneity may indeed benefit cell survival and thus be of advantage to *L. monocytogenes* under stressful conditions (Metselaar *et al.*, 2013). Furthermore, strain variability as well as population heterogeneity seem to increase along with harshening stress conditions (Francois *et al.*, 2006, Lianou and Koutsoumanis, 2013). Thus, in addition to unravelling specific traits rendering single strains tolerant, it is increasingly pivotal to comprehensively address strain variability in food hygiene research.

2.5.3 Plasmid-mediated stress tolerance in *L. monocytogenes*

Plasmids, like other MGEs, are profoundly self-seeking entities and are thought to essentially operate irrespective of the needs of their hosts (Rankin *et al.*, 2011). Carrying a plasmid may be costly for the host cell due to the synthesis of proteins needed in plasmid replication and transmission. However, plasmids frequently harbour genes that may engender fitness in the host and may even benefit the neighbouring cells by degrading harmful compounds (Kado, 1998, Löfmark *et al.*, 2008, Slater *et al.*, 2008, Rankin *et al.*, 2011). Plasmids may also provide their hosts with adaptive traits conferring enhanced antibiotic resistance or virulence (Poirel *et al.*, 2007, Herrero *et al.*, 2008, Périchon *et al.*, 2008, Carattoli *et al.*, 2010, Venturini *et al.*, 2010). Intriguingly, conjugative natural antibiotic resistance plasmids have also been isolated in commensal organisms, suggesting a resistance gene reservoir that may not be affected by a decrease in antibiotic usage and the following alleviation of selective pressure (Yates *et al.*, 2006). In *L. monocytogenes*, identical plasmids have been isolated from different geographic locations over significantly long time periods (Flamm *et al.*, 1984, Hingston *et al.*, 2017b), suggesting that these plasmids benefit their parental strains by improved resistance and survival: A conserved plasmid was found in four different strains isolated over a period of 12 years (Fox *et al.*, 2016).

Plasmids have been shown to confer tolerance of *L. monocytogenes* to stressors such as the disinfectant benzalkonium chloride (Elhanafi *et al.*, 2010, Rakic-Martinez *et al.*, 2011, Katharios-Lanwermeier *et al.*, 2012, Jiang *et al.*, 2016, Kremer *et al.*, 2017, Zhang *et al.*, 2018) and heavy metals

(Lebrun *et al.*, 1994, McLauchlin *et al.*, 1997, Rakic-Martinez *et al.*, 2011, Katharios-Lanweremeyer *et al.*, 2012). Hingston *et al.* (2017b) discovered the presence of plasmids to associate with enhanced acid tolerance and cold sensitivity of *L. monocytogenes*, and Zhang *et al.* (2018) reported pLMSZo8 plasmid curing resulting in decreased salt stress tolerance. However, knowledge of plasmids mediating environmental stress tolerance is rather limited and should be broadened, given the grave implications it has for food safety.

Plasmid-mediated antibiotic resistance, particularly to last-resort antibiotics, poses an emerging worldwide threat to both human and animal health (Liu *et al.*, 2016, Bachiri *et al.*, 2017). Concerningly, plasmids conferring resistance to antibiotics such as quinolones and the last-resort antibiotic colistin have been found in *E. coli* isolated from environmental samples (Jørgensen *et al.*, 2017, Ranjbar and Farahani, 2017). In *L. monocytogenes*, naturally occurring plasmids were already associated with enhanced resistance to antibiotics such as chloramphenicol, erythromycin and other macrolides, lincosamides, streptomycin and tetracycline in the early 1990s (Poyart-Salmeron *et al.*, 1990, Hadorn *et al.*, 1993). Thus, besides enhanced environmental fitness within bacteria, potential cotransfer of plasmid-borne stress resistance genes with genes conferring virulence and antibiotic resistance is of great concern (Herrero *et al.*, 2008, Bojer *et al.*, 2010, Gullberg *et al.*, 2014, Fox *et al.*, 2016). Plasmids may augment the ability of pathogenic hosts to occupy auxiliary niches, which creates further opportunities to infect new hosts. Conjugational transfer has been demonstrated not only between *Listeria* spp. strains (Katharios-Lanweremeyer *et al.*, 2012, Korsak *et al.*, 2019) but also to and from other species, such as *Lactococcus* spp. (Guglielmetti *et al.*, 2009), *Enterococcus* spp., *Staphylococcus* spp. (Poyart-Salmeron *et al.*, 1990) and *Streptococcus* spp. (Pérez-Díaz *et al.*, 1982, Flamm *et al.*, 1984, Poyart-Salmeron *et al.*, 1990), indicating naturally occurring inter-species horizontal gene transfer.

2.5.4 Two-component systems in the stress responses of *L. monocytogenes*

Signalling through TCSs is a fundamental means for bacteria to respond to and cope with versatile stresses. TCSs have been appointed a role among others in the cold tolerance and adaptation of *Clostridium botulinum* (Lindström *et al.*, 2012, Derman *et al.*, 2013, Dahlsten *et al.*, 2014, Mascher *et al.*, 2014) and *B. subtilis* (Aguilar *et al.*, 2001), as well as *B. subtilis* heat tolerance (Takada *et al.*, 2018), in the growth and motility of *Yersinia pseudotuberculosis* under cold conditions (Palonen *et al.*, 2011), and in the osmotolerance of *E. coli* (Nakashima *et al.*, 1992, Cai and Inouye, 2002). In addition to environmental stresses, TCSs are of importance in antibiotic resistance and the survival of pathogenic bacteria during host infection

(Beier and Gross, 2006, Fernández *et al.*, 2010, Altamirano-Silva *et al.*, 2018, Cao *et al.*, 2018). The exact physiological signal triggering the function of a particular TCS often remains obscure, but signal molecules such as nitrate, O₂, K⁺ and Mg²⁺ have been recognised (Krell *et al.*, 2010, Zschiedrich *et al.*, 2016). Specifically, changes in membrane fluidity and unsaturated FAs were recognised as the signals through which the TCS DesKR responds to cold stress in *B. subtilis* (Aguilar *et al.*, 2001).

TCSs with phenotypically confirmed roles in augmenting the tolerance of *L. monocytogenes* to environmental stress conditions closely related to food processing and hygiene are presented in Table 2. Besides heat tolerance, the orphan RR DegU (Lmo2515) is a pleiotropic temperature-responsive regulator needed for flagellar synthesis and motility and is also involved in biofilm formation and virulence (Knudsen *et al.*, 2004, Williams *et al.*, 2005b, Gueriri *et al.*, 2008). Intriguingly, DegU controls flagellar genes in an unphosphorylated state within the receiver domain (Mauder *et al.*, 2008). CheYA (Lmo0691/Lmo0692) is needed for the characteristic swarming motility of *L. monocytogenes* (Flanary *et al.*, 1999, Dons *et al.*, 2004, Williams *et al.*, 2005a). The *agr* quorum-sensing system of *L. monocytogenes*, including the TCS AgrCA (Lmo0050/Lmo0051), affects adherence on abiotic surfaces and therefore the early stages of biofilm formation (Rieu *et al.*, 2007). LiaSR (Lmo1021/Lmo1022), LisRK (Lmo1377/Lmo1378) and CesRK (Lmo2422/Lmo2421) are involved in cell-envelope stress and, accordingly, in the resistance of *L. monocytogenes* to cell wall-acting antibiotics, such as cephalosporins (Cotter *et al.*, 2002, Kallipolitis *et al.*, 2003, Gottschalk *et al.*, 2008, Collins *et al.*, 2012, Nielsen *et al.*, 2012). For instance, LiaSR responds to cell-envelope stress by upregulating membrane-associated genes and remodelling the composition of the cytoplasmic membrane (Fritsch *et al.*, 2011). Furthermore, LisRK controls the transcription of virulence gene *htrA* (Sleator and Hill, 2005). VirR (Lmo1745) regulates the modification of cell surface components and enhances cell invasion and virulence in *L. monocytogenes* (Mandin *et al.*, 2005). ResD (Lmo1948) confers resistance to pediocin, affects carbon source utilisation and controls respiration in *L. monocytogenes* (Larsen *et al.*, 2006).

Table 2. Two-component systems with phenotypically confirmed roles in enhancing the tolerance of *L. monocytogenes* to high and low temperatures, acid and alkali pH, and osmotic, ethanol and oxidative stresses.

Two-component system (HK/ RR) ^a	Stress condition	Phenotypic evidence ^b	Reference
AgrA; Lmo0051 (RR)	Osmotic	Reduced growth of deletion mutant in 3% and 5% NaCl (L)	(Garmyn <i>et al.</i> , 2012)
YycF; Lmo0287 (RR)	Cold	Upregulated transcription at 4 °C (L, S)	(Chan <i>et al.</i> , 2007b)
Lmo1060 (RR)	Cold	Reduced growth of deletion mutant at 4 °C (L)	(Chan <i>et al.</i> , 2008)
Lmo1172 (RR)	Cold	Reduced growth of deletion mutant at 4 °C (L)	(Chan <i>et al.</i> , 2008)
LisK; Lmo1378 (HK)	Acid	Reduced growth of deletion mutant at pH 3.5 (L)	(Cotter <i>et al.</i> , 1999)
	Osmotic	Reduced growth of deletion mutant in 8% NaCl (L)	(Sleator and Hill, 2005)
LisR; Lmo1377 (RR)	Acid	Reduced growth of insertion mutant at pH 5.25 (lag-phase inoculum)	(Kallipolitis and Ingmer, 2001)
	Cold	Reduced growth of deletion mutant at 4 °C (L)	(Chan <i>et al.</i> , 2008)
	Heat	Reduced growth of insertion mutant at 43.5 °C (L, S)	(Kallipolitis and Ingmer, 2001)
	Oxidative	Reduced growth of insertion mutant in 0.025% H ₂ O ₂ (L)	(Kallipolitis and Ingmer, 2001)
VirR; Lmo1745 (RR)	Ethanol	Reduced growth of deletion mutant in 5% ethanol (L)	(Williams <i>et al.</i> , 2005a)
ResD; Lmo1948 (RR)	Ethanol	Reduced growth of deletion mutant in 5% ethanol (L)	(Williams <i>et al.</i> , 2005a)
CesR; Lmo2422 (RR)	Heat	Reduced growth of insertion mutant at 43.5 °C (L, S)	(Kallipolitis and Ingmer, 2001)
	Osmotic	Reduced growth of insertion mutant in 9% NaCl (L, S)	(Kallipolitis and Ingmer, 2001)
PhoP; Lmo2501 (RR)	Ethanol	Reduced growth of deletion mutant in 5% ethanol (L)	(Williams <i>et al.</i> , 2005a)
DegU; Lmo2515 (RR)	Heat	Reduced growth of deletion mutant at 44 °C (L)	(Gueriri <i>et al.</i> , 2008)
	Ethanol	Reduced growth of deletion mutant in 5% ethanol (L)	(Williams <i>et al.</i> , 2005a)
KdpE; Lmo2678 (RR)	Heat	Reduced growth of insertion mutant at 43.5 °C (L, S)	(Kallipolitis and Ingmer, 2001)
	Osmotic	Reduced growth of insertion mutant in 2% and 9% NaCl (L, S)	(Kallipolitis and Ingmer, 2001, Brøndsted <i>et al.</i> , 2003)

^a HK, histidine kinase; RR, response regulator

^b L, logarithmic growth phase; S, stationary growth phase

3 AIMS OF THE STUDY

The aims of this study were to investigate the genetic mechanisms conferring stress tolerance in *L. monocytogenes* and to decipher strain diversity in *L. monocytogenes* stress resistance.

The specific aims of this study were:

1. To assess the role of two-component-system histidine kinases in the growth and stress tolerance of *L. monocytogenes* EGD-e at high and low temperatures, under acid, alkali and oxidative conditions, and in high salt and ethanol concentrations (I, II);
2. To identify accessory genetic mechanisms and plasmid-borne genes conferring heat resistance in *L. monocytogenes* (III);
3. To investigate strain variability and the occurrence of resistance-conferring genes with respect to benzalkonium chloride resistance within 392 *L. monocytogenes* isolates of Finnish and Swiss origin (IV).

4 MATERIALS AND METHODS

4.1 *L. monocytogenes* strains, isolates and plasmids (I–IV)

The sequenced *L. monocytogenes* EGD-e (Glaser *et al.*, 2001) was used for genetic modifications of HKs (I, II), *L. monocytogenes* isolates AL4E and AT3E (Lundén *et al.*, 2008) for plasmid-mediated heat resistance studies (III) and *L. monocytogenes* 10403S (Bishop and Hinrichs, 1987) for horizontal gene transfer experiments (III). The strains and plasmids used in studies I–III are presented in Table 3. A total of 392 *L. monocytogenes* isolates were selected for study IV from the culture collections of the Department of Food Hygiene and Environmental Health of the University of Helsinki, Finland (n = 197) and the Swiss National Reference Centre for Enteropathogenic Bacteria and Listeria, Switzerland (n = 195). The strains were of serotype 1/2a (n = 239), 1/2b (n = 44), 1/2c (n = 34), 3a (n = 5) and 4b (n = 70) and isolated between the years 1975 and 2013 (IV).

4.2 Typing (III, IV)

4.2.1 Serotyping (III, IV)

L. monocytogenes AL4E and AT3E and the 392 *L. monocytogenes* isolates were serotyped using the *Listeria* Antisera Set (Denka Seiken, Tokyo, Japan) including O- and H-factor antisera according to the manufacturer's instructions (III, IV) and by using the Pasteur typing tool (http://bigsd.b.pasteur.fr/perl/bigsd/bigsdb.pl?db=pubmlst_listeria_seqdef_public accessed 13 July 2017) (III).

4.2.2 Multilocus sequence typing (III, IV)

MLSTs of *L. monocytogenes* AL4E and AT3E were obtained from whole-genome sequences by using the PasteurMLST typing tool (Haase *et al.*, 2014, Moura *et al.*, 2016) (III). MLSTs of BC^r *L. monocytogenes* strains were defined by sequencing the PCR products of the seven housekeeping genes appointed to MLST typing (Ragon *et al.*, 2008) (IV).

Table 3. Bacterial strains and plasmids used in studies I–III.

Strain or plasmid	Genotype or characteristic ^a	Reference or source ^b
<i>Listeria monocytogenes</i>		
EGD-e	Wild-type strain, serotype 1/2a	(Glaser <i>et al.</i> , 2001)
AL4E	Wild-type strain, serotype 1/2c	(Lundén <i>et al.</i> , 2008)
AT3E	Wild-type strain, serotype 1/2c	(Lundén <i>et al.</i> , 2008)
10403S	Wild-type strain, serotype 1/2a, streptomycin-resistant	(Bishop and Hinrichs, 1987)
$\Delta lmo0050$	In-frame deletion of the <i>lmo0050</i> gene (1296 bp) with 1101 nucleotides	I, II
$\Delta lmo0288$ ($\Delta yycG$)	Disruption of the wild type with 1581 bp insertion including out-of-frame deletion of the <i>yycG</i> gene (1833 bp) with 1682 nucleotides	I, II
$\Delta lmo0692$ ($\Delta cheA$)	In-frame deletion of the <i>cheA</i> gene (1857 bp) with 1752 nucleotides	I, II
$\Delta lmo1021$	In-frame deletion of the <i>lmo1021</i> gene (1059 bp) with 930 nucleotides	I, II
$\Delta lmo1061$	Out-of-frame deletion of the <i>lmo1061</i> gene (1446 bp) with 1249 nucleotides	I, II
$\Delta lmo1173$	Out-of-frame deletion of the <i>lmo1173</i> gene (1458 bp) with 1078 nucleotides	I, II
$\Delta lmo1378$ ($\Delta lisK$)	In-frame deletion of the <i>lisK</i> gene (1452 bp) with 1116 nucleotides	I, II
$\Delta lmo1508$	In-frame deletion of the <i>lmo1508</i> gene (1440 bp) with 1230 nucleotides	I, II
$\Delta lmo1741$	In-frame deletion of the <i>lmo1741</i> gene (1041 bp) with 786 nucleotides	I, II
$\Delta lmo1947$ ($\Delta resE$)	In-frame deletion of the <i>resE</i> gene (1791 bp) with 1542 nucleotides	I, II
$\Delta lmo2011$	In-frame deletion of the <i>lmo2011</i> gene (1740 bp) with 1437 nucleotides	I, II
$\Delta lmo2421$ ($\Delta cesK$)	In-frame deletion of the <i>cesK</i> gene (1143 bp) with 1020 nucleotides	I, II
$\Delta lmo2500$ ($\Delta phoR$)	In-frame deletion of the <i>phoR</i> gene (1776 bp) with 1641 nucleotides	I, II
$\Delta lmo2582$	Out-of-frame deletion of the <i>lmo2582</i> gene (1380 bp) with 1339 nucleotides	I, II
$\Delta lmo2679$ ($\Delta kdpD$)	In-frame deletion of the <i>kdpD</i> gene (2691 bp) with 2652 nucleotides	I, II
$\Delta yycGc$	$\Delta yycG$, tRNA ^{Arg} ::pyycGc, complemented strain	I
$\Delta lisKc$	$\Delta lisK$, tRNA ^{Arg} ::plisKc, complemented strain	I
$\Delta agrCc$	Complemented strain, wild-type-like	II
$\Delta liaSc$	Complemented strain, wild-type-like	II
$\Delta virSc$	Complemented strain, wild-type-like	II
EGD-epPL2	EGD-e, tRNA ^{Arg} ::pPL2	(Markkula <i>et al.</i> , 2012b)

Table 3. Continued

Strain or plasmid	Genotype or characteristic ^a	Reference or source ^b
$\Delta yycGpPL2$	$\Delta yycG$, tRNA ^{Arg} ::pPL2	I
$\Delta lisKpPL2$	$\Delta lisK$, tRNA ^{Arg} ::pPL2	I
AT3Epc	AT3E strain, plasmid-cured	III
10403Sp $cplL$	10403S, tRNA ^{Arg} :: $cplL$	III
10403SpPL2	10403S, tRNA ^{Arg} ::pPL2	III
<i>Escherichia coli</i>		
DH5 α pMAD	DH5 α strain containing the shuttle vector plasmid pMAD	(Mattila <i>et al.</i> , 2011)
TOP10	Electrocompetent strain	Invitrogen
NEB5 α	Electrocompetent strain	New England Biolabs
NEB5 α	Chemically competent strain	New England Biolabs
HB101	Conjugation donor containing the helper plasmid pRK24	CRBIP
Plasmid		
pMAD	Cloning shuttle integration vector plasmid	(Arnaud <i>et al.</i> , 2004)
pMAD- $\Delta yycG$	pMAD containing homologous region up- and downstream of EGD-e <i>yycG</i>	I
pMAD- $\Delta lisK$	pMAD containing homologous region up- and downstream of EGD-e <i>lisK</i>	I
pMAD- $\Delta agrCc$	pMAD containing coding sequence of EGD-e <i>agrC</i>	II
pMAD- $\Delta liaSc$	pMAD containing coding sequence of EGD-e <i>liaS</i>	II
pMAD- $\Delta virSc$	pMAD containing coding sequence of EGD-e <i>virS</i>	II
pPL2	Site-specific integration vector	(Lauer <i>et al.</i> , 2002)
pyycGc	pPL2 containing 613 bp upstream nucleotides and the coding sequence of <i>yycG</i>	I
plisKc	pPL2 containing 1443 bp upstream nucleotides, the coding sequence and downstream terminator sequence of <i>lisK</i>	I
pLM58	Plasmid, AT3E strain	III
$cplL$	pPL2 containing 423 bp of upstream nucleotides and the coding sequence of ATP-dependent protease <i>cplL</i>	III

^a bp, base pairs; cds, coding sequence

^b CRBIP, Biological Resource Centre of the Institut Pasteur

4.3 Growth conditions (I–IV)

All the strains were preserved in vials with beads (I–III) or in brain–heart infusion (BHI) broth (Oxoid, Cheshire, England) supplemented with 20% glycerol (Sigma-Aldrich, St. Louis, MO, USA) (IV) at -80°C . *L. monocytogenes* strains and isolates were routinely grown at 37°C on blood, on tryptic soy (TS) agar or in TS broth (Oxoid), on BHI agar or in BHI broth (BD, Franklin Lakes, NJ, USA and Oxoid), on Mueller-Hinton (MH) agar or in MH broth (Oxoid) or on ALOA (Harlequin™ *Listeria* Chromogenic) agar (Labema, Helsinki, Finland). *E. coli* strains were grown on Luria-Bertani (LB) agar or in LB broth (BD and Oxoid). Appropriate antibiotics

(Sigma-Aldrich), cadmium sulphate ($3 \text{ CdSO}_4 \cdot 8 \text{ H}_2\text{O}$) (Merck, Darmstadt, Germany) and reserpine (Sigma-Aldrich) were added when needed.

4.4 Transcriptional analyses (I)

The transcription levels of each of the 15 HK-encoding genes in *L. monocytogenes* EGD-e were measured from total RNA extracted in the logarithmic growth phase (OD_{600} , 0.7) at 3 °C and at 30 min, 3 h and 7 h post-cold shock at 5 °C. Each experiment was repeated three times. Prior to the RNA extraction, the cells were lysed with 25 mg/ml lysozyme and 250 IU/ml mutanolysin (Sigma-Aldrich) in Tris-EDTA (Fluka Biochemica, Buchs, Switzerland) buffer in a 37 °C water bath for 30 min.

4.4.1 Total RNA extraction (I)

Total RNA was extracted by using the RNeasy Midi-kit (Qiagen, Venlo, the Netherlands). DNase treatment was performed by using the RNase-Free DNase set (Qiagen) and an additional DNase treatment by using the Ambion DNA-free kit (Ambion, Austin, TX, USA). The quantity of RNA was measured with a Nanodrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA), and RNA integrity was verified by using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA).

4.4.2 Real-time quantitative reverse transcription-PCR (I)

Reverse transcription from 800 ng of each total-RNA sample into complementary DNA (cDNA) was performed using the DyNAmo cDNA Synthesis Kit (Thermo Fischer Scientific, Inc., Waltham, MA, USA) in duplicate reactions. An additional predenaturation at 65 °C for 5 min and an elongated cDNA synthesis step at 40 °C for 40 min were used. A control reaction lacking the reverse transcriptase enzyme (no-RT control) was performed for each RNA sample. For real-time quantitative PCR (qPCR), each cDNA sample was diluted 1:1000. qPCR reactions were performed in duplicate in a RotorGene qPCR 3000 device (Corbett Research, Sydney, Australia) with the DyNAmo Flash SYBR Green qPCR Kit (Thermo Fischer Scientific, Inc.). The relative expression levels of the target HK-encoding genes in the logarithmic growth phase at 3 °C and at 30 min, 3 h and 7 h post-cold shock at 5 °C were calibrated to their expression levels in the logarithmic growth phase at 37 °C and to t_0 pre-shock, respectively, according to the Pfaffl method (Pfaffl, 2001). This method intrinsically includes the assay efficiencies, which for the target genes ranged from 0.88 to 1.00 and for the reference was 0.82 (Table 4). The 16S rRNA-encoding gene *rrn* was used as an internal normalization reference (Tasara and Stephan, 2007).

Table 4. Primers used in studies I–IV.

Application	Primer	Sequence (5' to 3') ^{a, b, c}	Reference
Real-time reverse transcriptase quantitative PCR			
	0050 F	AATCGCAGGTTTTGATGGA	I
	0050 R	AGTTCGTTGGTTGCCGTATT	I
	0288 F	AGCCCAAGCAATGACTGAAG	I
	0288 R	CGTGTCCCCGATGTCTAAAA	I
	0692 F	CGTCGTGAAAAATGCCAAG	I
	0692 R	CTAAAATCGTTGCCCCAGAA	I
	1021 F	ATGGCTGTTTGCTCAGGTTTA	I
	1021 R	GAAATGGCAGTAAGCGGAAC	I
	1061 F3	TCGTAAAAGCAGGCGAAGC	I
	1061 R3	TCGTGCCGTCTACAACAGTC	I
	1173 F	AAGAGGACGAGCAGGAATG	I
	1173 R	CGCAATAAGGAAACAACAGA	I
	1378 F	TGATGGGCAGAAGATGATGAC	I
	1378 R	GGAAGTGAGCGGATTTACCA	I
	1508 F	CTGCGGATGATAAGAAAAAG	I
	1508 R	AGCACGATAGAACCGACA	I
	1741 F	TCTCGTGCTGGGCTATTTCT	I
	1741 R	GTTTCTTCGCCATTAGTTGGT	I
	1947 F	GAACGGTGAAACTGGTGTTG	I
	1947 R	ATGCGAGAAGAAAAGACGAAAG	I
	2011 F2	CAAATGGGTTATCTGCTCCTTTAC	I
	2011 R2	TGTTTCGTTGGATGGTTGG	I
	2421 F2	CCCAGCAAGCCAGAAATG	I
	2421 R2	AAAATGGATGTAAGTGGTGTCC	I
	2500 F	TGGTAGAGGTGGACGAAG	I
	2500 R	AATAGTAACGGATGGGATTG	I
	2582 F	CAACTACTCACACTCGCTTCACTT	I
	2582 R	CCAACTCCACTCGGTCATTT	I
	2679 F	GCTAATGGGAGCGTTTGGT	I
	2679 R	TGTGTGAGCGGACCTTCTTC	I
	16S rRNA	GATGCATAGCCGACCTGAGA	(Mattila <i>et al.</i> , 2011)
	Forward		
	16S rRNA	CTCCGTCAGACTTTTCGTCCA	(Mattila <i>et al.</i> , 2011)
	Reverse		
Mutant construction			
	0050-1	NNNNNN <u>GGATCC</u> TTATGGCTCAGACGGTATTCT	I, II
	0050-2	<i>TCATCCCCATTCTTCTACATGGCACTTACAAAAACAATCA</i>	I, II
	0050-3	<i>TGATTGTTTTGTAAAGTGCCATGTAGAAGATGGGGATGAA</i>	I, II
	0050-4	NNNNNN <u>ACGCGT</u> CCACACCTTTTGTCGTATCT	I, II
	0288-1 MluI	NNNNNN <u>ACGCGT</u> CTTGTAGTAGATGACGAAAAACC	I, II
	0288-2	<i>CTAACCTGTTCCTCCCATTGTAAGCGAAAAATACCCAAAA</i>	I, II

Table 4. *Continued*

Application	Primer	Sequence (5' to 3') ^{a, b, c}	Reference
	0288-3	TTTTGGGTATTTTCGCTTACAATGGGAGGAACAGGGTTAGG	I, II
	0288-4	NNNNNNACGCGTGCCTTAGGAAAAGATTGCGTTTG	I, II
	0692-1	NNNNNNGGATCCCGTGAACAATCAATCCATCG	I, II
	0692-2	CTGAAAAAGCCGCCAAGTGTCATGCTATCCACCTCCATTTCT	I, II
	0692-3	AAATGGAGGTGGATAGCATGACACTTGGCGGCTTTTTCAG	I, II
	0692-4	NNNNNNACGCGTGCCTGTTTCCCTTTTTCCTC	I, II
	1021-1 MluI	NNNNNNACGCGTCGTAAGGAGGTGTGTCAGC	I, II
	1021-2	TTATTTTGATGGTGCCGGAGCAAACAGCCATCATCAG	I, II
	1021-3	CTGATGATGGCTGTTTGCTCCGGCACCATCAAATAA	I, II
	1021-4	NNNNNNACGCGTCACCAAATCGCCCAAGAC	I, II
	1061-1 MluI	NNNNNNACGCGTAAAAAGAACAGTAGGAGGCAAA	I, II
	1061-2	ATAATAAACAAACCTAAACCAGAGCCGTAAAAAGTAAATGA	I, II
	1061-3	ATTTTACTTTTACGGCTCTGGTTTAGGGTTGTTTATTATTGC	I, II
	1061-4	NNNNNNACGCGTCGCTGTATTTTCTTTGTCATCTG	I, II
	1173-1 MluI	NNNNNNACGCGTCGTGACAGGAATGAATGGAAG	I, II
	1173-2	ATGACCGTTGTCCGTTACTTTTTGGTTGTTTTGGTGGTG	I, II
	1173-3	CACCACCAAAAACAACCAAAAAGTAACGGACAACGGTCAT	I, II
	1173-4	NNNNNNACGCGTATCAAACAGGGATAAGTTCTG	I, II
	1378-1 MluI	NNNNNNACGCGTAGCGGTTGCTAATGATGGAC	I, II
	1378-2	TCTGGCTTTATCTACACGGTGCAGTCTTTTACTTCTGTTTCTCT	I, II
	1378-3	ACCAGAAGTAAAGAAGTGCACCGTGTAGATAAAGCCAGAAGC	I, II
	1378-4	NNNNNNACGCGTCGAAAGAAGAACAAGCAACTATTC	I, II
	1508-1 MluI	NNNNNNACGCGTAATGTAAGAGGAGTGAGAACAGAA	I, II
	1508-2	CCAATACCAGATTCGCCACAAGCAATGAGCAAAATGATAAA	I, II
	1508-3	ATCATTTTGCTCATTGCTTGTGGCGAATCTGGTATTGG	I, II
	1508-4	NNNNNNACGCGTACAGGGCGTCGTAAAGAAGT	I, II
	1741-1 MluI	NNNNNNACGCGTGGTCAAAAACAACCCCGATA	I, II
	1741-2	CTCGTGCTGGGCTATTTCTGTGGCGTAGGAACAGAAGTGA	I, II
	1741-3	CACTTCTGTTCTACGCCACAGAAATAGCCCAGCACGAGA	I, II
	1741-4	NNNNNNACGCGTAAGGCGGTAAGTTTGTTCCTCA	I, II
	1947-1 MluI	NNNNNNACGCGTGCCCCGAAGATGAACGAATA	I, II
	1947-2	CAAAACAGCCAGTATCATATGGAAGCACACAATGGAAAA	I, II
	1947-3	TTTTCCATTGTGTGCTTCCATGATGATACTGGCTGTTTTGTCT	I, II
	1947-4	NNNNNNACGCGTTTTTGTCTGTAAGTGAAGGAGT	I, II
	2011-1 MluI	NNNNNNACGCGTGAAGTTTGTTCGGTTATTCCAA	I, II
	2011-2	CAATCTCCAAAGAGCGGAAGATACGAGCCACATCGGTCT	I, II
	2011-3	AGACCGATGTGGCTCGTATCTTCCGCTCTTTGGAGATTG	I, II
	2011-4	NNNNNNACGCGTGTGGGATTGTGTTTGGCTTT	I, II
	2421-1	NNNNNNGGATCCGAAATCTCGCCTTGTTCA	I, II
	2421-2	GTTTCTTGGGTTTTGGCGTTATCGTCAAAGTCCACTAAT	I, II
	2421-3	TAGTGGCAGTTTGACGATAAACGCCAAAACCAAGAAAC	I, II
	2421-4	NNNNNNGAATTCGACTGGAAAGGACAATGAGTATGA	I, II
	2500-1 MluI	NNNNNNACGCGTCATTCCGACAACCGCTACTC	I, II

Table 4. *Continued*

Application	Primer	Sequence (5' to 3') ^{a, b, c}	Reference
	2500-2	TATGGCTGAAAATCGGGTTTTCCATCGTCAAACATCTCG	I, II
	2500-3	CGAGATGTTTGACGATGGAACCCGATTTTCAGCCATA	I, II
	2500-4	NNNNNNACGCGTGCTTCTATTGTTACCTTGCTACA	I, II
	2582-1	NNNNNNACGCGTTCAGCCAACGAAAATAAC	I, II
	2582-2	AGTTGTCACGATGCTATCCGATTCACATTCGGTTT	I, II
	2582-3	AAACGGAATGTGAATCGACAAGCATCGTGACAACT	I, II
	2582-4	NNNNNNACGCGTGAGGCAAGTGATGGGGTAGA	I, II
	2679-1	NNNNNNACGCGTTTCTTCTGCCATCCGGTATC	I, II
	2679-2	ATGGAACGAATCGTCCAAGCGGAGAGATGGAAATGAA	I, II
	2679-3	TTCATTTTCCATCTCCTCCGCTTGACGATTCGTTTCCAT	I, II
	2679-4	NNNNNNACGCGTGCGTGCTTTATCCAGGTGTT	I, II
Complementation and conjugation			
	Komp 0288F1	NNNNGGATCCGGTGCGGATGACTATGTAACG	I
	Komp oper	NNNNTCTAGACCATTTTCTTCCACTCAAACC	I
	0288R		
	Komp oper	NNNNGGATCCTTTGCTCGTTACATTTCTGC	I
	1378F2		
	Komp oper	NNNNTCTAGATTATTCGTTTCCTTCACAGC	I
	1378R2		
	BamHI <i>clpL</i> FN	NNNNGGATCCAGTTTCAAAGGTCGTTCTGGC	III
	BamHI <i>clpL</i> RN	NNNNGGATCCTCTATCAAGCAATCTCCTTCCC	III
	NC16	GTCAAAACATACGCTCTTATC	(Lauer <i>et al.</i> , 2002)
	PL95	ACATAATCAGTCCAAAGTAGATGC	(Lauer <i>et al.</i> , 2002)
Small-scale sequencing			
	Seq 0050 1F	ATGACAAGAATCGGACATTT	I
	Seq 0050 1R	TTTTTACATACCTTTTGCCTGA	I
	Seq 0050 2F	CGGAAAACACTAAAAAGAAAAGC	I
	Seq 0050 2R	CCGTTCTTCTGCTTGCTTC	I
	Seq 0288 1F	TTATGAAGGCAGGCCAGAAT	I
	Seq 0288 1R	GGAATAGTGCGGACGTGAAT	I
	Seq 0288 2F	CAGCCGAGGAAGAAGAAAAC	I
	Seq 0288 2R	GTGAAATAAACCGAATGTAGCC	I
	Seq 0692 1F	TTATCCGTTGCTGATGCTGA	I
	Seq 0692 1R	TGCGTTTTCTCCCTTTTAC	I
	Seq 0692 2F	CGAGGGGCTTTTCTTTTGAT	I
	Seq 0692 2R	GACCTTTTCCGTTTGGATAG	I
	Seq 1021 1F	GGAAGAAACATCCGACTAAAG	I
	Seq 1021 1R	CGCATTTTCATCCGTTG	I
	Seq 1021 2F	AAAGTTGGGCTTGTTGTTGC	I
	Seq 1021 2R	TTTTTCTGGTTTAGCGGTTAGC	I
	Seq 1061 1F	CGTTCGCTACTTGATGATG	I

Table 4. *Continued*

Application	Primer	Sequence (5' to 3') ^{a, b, c}	Reference
	Seq 1061 1R	CCAGGGAGCATCATTTG	
	Seq 1061 2F	ATGCCGAGTATGGACGGGTA	
	Seq 1061 2R	ACCTTTTGATGCGGAACCTT	
	Seq 1173 1F	GCTCTTTTACTCTTTCGCTCAA	
	Seq 1173 1R	TGATTTTCTTGTTTCACGGTCTAC	
	Seq 1173 2F	ACACGGAGAAAGCAAAAGGA	
	Seq 1173 2R	ACAACAAAATCCCCAGCAAA	
	Seq 1378 1F	TAACGAGTGGGTGCCTTTTT	
	Seq 1378 1R	TACAGATGGGTTTTTCAGCA	
	Seq 1378 2F	GCTAAACAAACAACGCTACAG	
	Seq 1378 2R	CGCCCGAACAAAAGAAA	
	Seq 1508 1F	TGGGGAGTTATGAAGAAAGTATGAA	
	Seq 1508 1R	TCCAAGTTTTAGCACCAATGT	
	Seq 1508 2F	TCCGTGAAGTAAGCGATGTG	
	Seq 1508 2R	TGTGATTATGCCGATTGTCC	
	Seq 1741 1F	AAGCAATAAACACGGCATCC	
	Seq 1741 1R	TTGAAAAATGGAGAAGTAGTCGT	
	Seq 1741 2F	AGTGTGTGTTTTGCCCATC	
	Seq 1741 2R	TTGTTGCTGTTGGCGTTAGT	
	Seq 1947 1F	AACGAAAAGGTCTGCCAACA	
	Seq 1947 1R	TGATTTGATGATGCCTGGTAA	
	Seq 1947 2F	TTTACCATCCGCACCCAAT	
	Seq 1947 2R	AGAATCAGCAGGAGGCACAC	
	Seq 2011 1F	CGTTTTTGTTAGGCGATTAGAC	
	Seq 2011 1R	CTTAGTATGGCTTTTCCTCGTATT	
	Seq 2011 2F	GACGCTCTTCCCATTCTTCTT	
	Seq 2011 2R	ACGGTGTTTGTCCACTTCCA	
	Seq 2421 1F	ACAGTGCGGGACCATAATAAA	
	Seq 2421 1R	GATGACGATTGAAAGTGCTCA	
	Seq 2421 2F	GCATTTCCCTCCACCACTCT	
	Seq 2421 2R	CAAGGCTGAAGACAACGACA	
	Seq 2500 1F	TTTTCGTCTTTACCGCCAAC	
	Seq 2500 1R	AACAGCACTTTATGAGAATGAATC	
	Seq 2500 2F	TTTGTGCTTTAGGGTTAGTGTGTTG	
	Seq 2500 2R	CCAAGAGAAGTAGTGGCGAGA	
	Seq 2582 1F	CATCCGAAACAGACAGCGTA	
	Seq 2582 1R	TACGCTGTCTGTTTCGGATG	
	Seq 2582 2F	AAAATTGGCAACCAAGAACG	
	Seq 2582 2R	TCATCTGTGCCCACTTCAAA	
	Seq 2679 1F	ATCAAGCGCTTCGAGTCAAT	
	Seq 2679 1R	ACACGAATATCCCGAAGCAC	
	Seq 2679 2F	TGCTGCGGCATATCAAGTAG	
	Seq 2679 2R	AAACGTTTGGCCAATTTTCAAG	

Table 4. *Continued*

Application	Primer	Sequence (5' to 3') ^{a, b, c}	Reference
Confirmation of plasmid curing			
	<i>oriV</i> -F	GAACAAGCGATCCGTCATGC	III
	<i>oriV</i> -R	TCGTTGCTAGGACTTGTCTGG	III
	<i>clpL</i> -F	ACAGGCTCGTGATGGCTTAC	III
	<i>clpL</i> -R	ACCGCGATATTGAGTTCCCG	III
Plate mating			
	ESAT-6 F	GCAATCAGTGCGGAAGGACTG	III
	ESAT-6 R	ATCCATCGCTTGTTTTCTCG	III
	<i>secA</i> -F	ACTACTGCCAAAACATCGAAGC	III
	<i>secA</i> -R	AAGACGCACTGGATTCCCTC	III
Screening efflux pump systems			
	p1	CATTAGAAGCAGTCGCAAAGCA	(Elhanafi <i>et al.</i> , 2010)
	p2	GTTTTCGTGTCAGCAGATCTTTGA	(Elhanafi <i>et al.</i> , 2010)
	radC fwd	CTTGCCAATGATAATATCATC	(Müller <i>et al.</i> , 2013)
	radC rev	GTGGTCTGAATGCTCCATCG	(Müller <i>et al.</i> , 2013)
	EmrE fw	GACCAACACCACCTAAGT	IV
	EmrE rv	GTCTGATGGACTTACAAAGCT	IV

^a N, any of the bases, i.e., adenine (A), cytosine (C), guanine (G) or thymine (T).

^b Restriction sites are underlined.

^c Overlapping 5'-end extensions of splicing-by-overlap extension PCR are marked in italics.

4.5 Genetic modifications (I–III)

In order to identify the role of HKs in the growth of *L. monocytogenes* EGD-e under different stresses, HK-encoding genes were deleted and restored (I, II). To assess the role of plasmid-borne ClpL protease in the heat resistance of *L. monocytogenes*, the plasmid-carrying AT3E strain was cured and the ClpL protease-encoding *clpL* gene introduced into heat-sensitive *L. monocytogenes* 10403S (III).

4.5.1 Knock-out deletion (I, II)

HK-encoding genes were individually deleted in *L. monocytogenes* EGD-e by allelic replacement using a pMAD shuttle vector (Arnaud *et al.*, 2004), resulting in mutants without an associated antibiotic resistance gene. Target gene-deficient inserts with 700 to 800 bp up- and downstream regions were built by using splicing-by-overlap extension (SOE) PCR (Warrens *et al.*, 1997). To construct the vector pMADΔ*target_gene*, the inserts were

individually ligated between the *Mlu*I and *Bam*HI or *Eco*RI sites of pMAD using T4 ligase (Thermo Fischer Scientific, Inc.). Propagated vectors were then separately electroporated (25 μ F, 200 Ω , 2.3 kV) into *L. monocytogenes* EGD-e. Single-crossover mutants were selected at 39 °C with erythromycin (5 μ g/ml) and double-crossovers without antibiotic. Deletions were confirmed by site-specific PCR and by sequencing the target regions (Institute of Biotechnology, University of Helsinki, Finland). Considering the putatively essential *yycG* gene, the mutational approach resulted in the disruption of the wild-type sequence of the *lmo0288–lmo0290* operon with the target gene-deficient insert, while harbouring the *yycG* coding sequence upstream of the insert. The primers used in mutant construction, PCR and small-scale sequencing are presented in Table 4.

4.5.2 Complementation (I, II)

To verify stress-sensitive mutant phenotypes, the coding sequences of the target gene wild-type copy with the related upstream regions, including the putative promoters (I, II), and related genes in putative operons (Toledo-Arana *et al.*, 2009) (I) were transformed into the respective phenotypically affected mutant strains. Complementation was performed by using the PSA prophage site-specific integration vector pPL2, provided by Prof. Martin Loessner (Swiss Federal Institute of Technology, Zürich, Switzerland) according to Lauer *et al.* (2002) (I), or by allelic replacement without an integrated plasmid using the pMAD vector (Arnaud *et al.*, 2004) (II). The amplified inserts were restricted with *Xba*I and *Bam*HI and ligated into the compatible *Spe*I and *Bam*HI sites of pPL2. The resultant plasmids were then conjugated into recipient *L. monocytogenes* strains according to Ma *et al.* (2011), and the strains carrying the pPL2 constructs were selected by chloramphenicol (25 μ g/ml) on ALOA agar. Integration of pPL2 was verified by PCR using primers NC16 and PL95 (Lauer *et al.*, 2002) and the presence of the insert by using gene-specific primers (Table 4) (I). When complementation was performed by allelic replacement, inserts were amplified using the outermost primers for mutant strain construction (Table 4), ligated into the *Mlu*I site of pMAD, and transformed into recipient mutant strains according to Markkula *et al.* (2012b). Successful complementation was confirmed by PCR using gene-specific and outermost mutant construction primers (Table 4) (II).

4.5.3 Plasmid curing (III)

pLM58 plasmid curing in the heat-resistant *L. monocytogenes* AT3E was conducted using ten consecutive cultures at 40 °C for 24 h per incubation in TSB supplemented with 0.2 μ g/ml novobiocin (Margolles and de los Reyes-Gavilán, 1998). Inoculations (1:100) into fresh broth were conducted between each incubation and final cultures were plated on TSA containing

novobiocin. Removal of the plasmid was confirmed from purified colonies by PCR with the pLM58-specific primers *oriV*-F and *oriV*-R and *clpL*-F and *clpL*-R (Table 4).

4.5.4 Plate mating (III)

To examine whether pLM58, harboured by *L. monocytogenes* AT3E, is self-transmissible, standard plate mating was conducted between *L. monocytogenes* wild-type AT3E and 10403S. The plasmid-borne *cadAC* genes, also found in pLM58, confer cadmium resistance (McLauchlin *et al.*, 1997, Lebrun *et al.*, 1994), which enables the exclusion of recipient cells that have not received pLM58. The innate streptomycin resistance of 10403S allows the selection of transconjugants from the donor (den Bakker *et al.*, 2012). Streptomycin-resistant 10403S, received from Prof. Martin Wiedmann, Cornell University, Ithaca, NY, USA, was routinely grown in BHI supplemented with 200 µg/ml streptomycin. Equal volumes (100 µl) of donor and recipient cells, grown to the logarithmic growth phase (OD₆₀₀, 0.5), were spotted on top of each other on BHI agar and incubated at room temperature for 1 h followed by incubation at 37 °C for 24 h. Transconjugants were screened on BHI containing 200 µg/ml streptomycin and 65 µg/ml or 130 µg/ml cadmium sulphate (CdSO₄). Colonies were screened by PCR using pLM58-specific *oriV* primers, and genomic AT3E ESAT-6 and 10403S *secA* primers (Table 4). To serve as both positive and negative controls, AT3E and 10403S were separately plated on BHI agar containing 130 µg/ml CdSO₄ or 200 µg/ml streptomycin.

4.5.5 Introducing *clpL* into a heat-sensitive *L. monocytogenes* strain (III)

To verify that *clpL* plays a role in the heat resistance of *L. monocytogenes*, the 423-bp upstream region including the putative promoter, and the coding sequence of *clpL* were amplified using primers BamHI *clpL* FN and BamHI *clpL* RN (Table 4) and conjugated into the heat-sensitive *L. monocytogenes* 10403S by filter mating using the integration vector pPL2 as described above (Lauer *et al.*, 2002, Ma *et al.*, 2011). Control pPL2 lacking the insert was separately conjugated into the recipient 10403S. Integration of the plasmids was confirmed by using NC16 and PL95 (Lauer *et al.*, 2002) and the presence of the insert by using *clpL* gene-specific primers (Table 4).

4.6 Phenotypic characterisation of *L. monocytogenes* strains and isolates (I–IV)

4.6.1 Growth curve analyses (I–III)

To analyse the growth of the *L. monocytogenes* wild-type and genetically modified strains exposed to cold, heat, osmotic, acid, alkali, ethanol or oxidative conditions, three to five separate colonies per strain were inoculated into 10 ml of BHI broth and incubated overnight at 37 °C (I–III). The cultures were diluted (1:100) in BHI broth (I–III) or BHI supplemented with 6% NaCl, 3.5 vol% ethanol or 5 mM H₂O₂, or BHI adjusted to pH 5.6 with 37% HCl or pH 9.4 with 5 M NaOH (II). Each dilution (300–350 µl) was transferred into separate wells of a 100-well honeycomb plate in duplicate technical replicates. The strains were grown in a Bioscreen C Microbiology Reader (Growth Curves Ltd., Helsinki, Finland) at 3 °C for 23 to 40 days (cold), at 42.5 °C or 42 °C for 10 to 24 h (heat), and at 37 °C for 10 to 48 h (other stresses). OD₆₀₀ values were measured at 15-min or 1-h intervals. Control growths were performed at 37 °C for 24 h (I, II). The mean growth parameters, including the maximum growth rate, onset time of growth (lag phase) and maximum optical density, were obtained by using the DMFit web edition software (Computational Microbiology Research Group, Institute of Food Research, Colney, Norwich, UK) with the complete Baranyi and Roberts model (Baranyi and Roberts, 1994) (I, II), or by using the ‘grofit’ package (Kahm *et al.*, 2010) in R versions 3.2.2 and 3.4.0 (R Core Team, Vienna, Austria) applying the logistic model with the default settings (III). The area under the curve (AUC) was determined in ‘grofit’ (III) or by using a designed script in Matlab R2014a (MathWorks, Natick, MA, USA) (II). The correspondence between the OD₆₀₀ values and the viable cell numbers of the wild-type and genetically modified strains was confirmed by plate counts in early and late logarithmic and early stationary growth phases (I, II), or mid-logarithmic and stationary growth phases (III).

4.6.2 Heat resistance assay (III)

L. monocytogenes strains 10403S, AL4E and AT3E, plasmid-cured AT3Epc, and conjugation strains 10403SpPL2 and 10403SpclpL were tested for heat survival according to Lundén *et al.* (2008), except that three overnight cultures of each strain were separately inoculated into TSB and challenged at 55 °C for 40 min. The decrease in the cell concentration (log₁₀ reduction) was determined by colony counts on plate count (PCA) agar (Oxoid). Strains with a reduction of <1.0 log₁₀ were considered heat resistant.

4.6.3 Minimum and maximum growth temperatures (I–III)

The differences in minimum and maximum growth temperatures were examined using the Gradiplate W10 incubator (BCDE Group, Helsinki, Finland) (Hinderink *et al.*, 2009). Dilutions of two (I, II) and three (III) separate overnight cultures of each strain were plated in duplicate by the stamping technique as parallel lines onto TSA (25 g agar/l). Minimum temperatures were measured for 21 days with a temperature gradient set to range from 1 °C to 7 °C (I). Differences in maximum growth temperatures were measured with a temperature gradient of 39 °C to 44 °C for 48 h (II) or 39.2 °C to 45.7 °C for 24 h (III). Growth boundaries were determined by using a stereomicroscope (Olympus SZ61, Nikon, Tokyo, Japan). The growth temperature threshold was determined as the boundary where dense bacterial growth discontinued (I–III).

4.6.4 Benzalkonium chloride resistance and impact of reserpine (IV)

The susceptibility of *L. monocytogenes* isolates to BC was tested using the agar dilution method (Elhanafi *et al.*, 2010). Single colonies of each strain were suspended into 100 µl of MH broth (Oxoid), of which 5 µl was spotted in technical duplicates onto MH agar (Oxoid) supplemented with 0 to 30 µg/ml BC. MICs were measured after 48 h at 37 °C. Strains displaying confluent growth in ≥ 20 µg/ml were considered BC resistant (BC^r).

Efflux pump activity in BC^r *L. monocytogenes* isolates was assessed by measuring the BC MICs on MH agar (Oxoid) supplemented with 0 µg/ml to 30 µg/ml BC and an additional 20 µg/ml of the efflux pump inhibitor reserpine (Sigma-Aldrich). The BC resistance of the *L. monocytogenes* isolates was categorised as fully efflux pump-dependent when reserpine caused a ≥ 10 µg/ml decrease in the BC MIC and partially dependent when the resultant decrease was < 10 µg/ml.

4.7 Genomic analyses (III)

4.7.1 DNA extraction (III)

After growth to the stationary growth phase at 37 °C, the cells of the *L. monocytogenes* AT3E and AL4E strains were lysed using lysozyme and mutanolysin at 37 °C for 2 h. Genomic DNA was extracted by using guanidium thiocyanate (Pitcher *et al.*, 1989). DNA integrity was verified by gel electrophoresis and the DNA yield was measured by using the Qubit dsDNA BR Assay Kit (Life Technologies, Carlsbad, CA, USA).

4.7.2 Sequencing (III)

WGS of *L. monocytogenes* AL4E and AT3E was conducted by the Institute of Biotechnology (Helsinki, Finland) using single-molecule real-time (SMRT) sequencing in the PacBio RS platform with coverages of 199x (AL4E), 252x (AT3E) and 422x (pLM58) (III). AL4E and AT3E genomes were *de novo* assembled according to the RS_HGAP_Assembly 3 protocol (Pacific Biosciences of California, Inc., Menlo Park, CA, USA).

4.7.3 Annotation and comparative genomic analysis (III)

L. monocytogenes AL4E and AT3E genomes were annotated by using RAST 2.0 (Aziz *et al.*, 2008) and genome comparison was conducted in SEED Viewer 2.0 (Overbeek *et al.*, 2005). Sequence-based operon prediction of pLM58 was performed by using FGENESB (Solovyev and Salamov, 2011) and putative prophages of pLM58 were predicted by using PHASTER (Zhou *et al.*, 2011, Arndt *et al.*, 2016). Circular visualisation and comparison of AL4E and AT3E chromosomes was conducted by using BRIG (BLAST Ring Image Generator) (Alikhan *et al.*, 2011) and pLM58 was visualized by using SnapGene Viewer 3.3.4 (GSL Biotech LLC, Chicago, IL). The coding sequences of the replication initiation protein and the ATP-dependent protease ClpL harboured by pLM58 were compared using BLASTN 2.2.26 (Altschul *et al.*, 1997) to *L. monocytogenes* plasmid sequences deposited in GenBank at NCBI by 13 July 2017.

4.8 Statistical analyses (I–IV)

The statistical analyses were performed in Excel 2010 (Microsoft, Redmond, WA, USA) (I), in IBM SPSS Statistics 23 and 24 (IBM, Armonk, NY, USA) (II, III) and using the JMP program v. 11.0.0 (SAS Institute, Inc., NC, USA) (IV). Differences in the expression levels of the HK-encoding genes were tested by the paired *t*-test (I). Differences in the growth parameters (I–III), the minimum and maximum growth temperatures (I–III) and the log₁₀ reductions (III) were tested using the independent-samples two-tailed *t*-test. Fisher's exact test was used in pairwise comparisons of BC resistance and sensitivity within serotypes and isolate origin (IV).

5 RESULTS

5.1 Roles of two-component-system histidine kinases in the stress tolerance of *L. monocytogenes* (I, II)

5.1.1 Relative expression levels of histidine kinase-encoding genes at low temperatures (I)

At 3 °C, *cheA* (*lmo0692*) was upregulated 236-fold ($P < 0.001$) as compared to the expression level at 37 °C, while *liaS* (*lmo1021*), *lmo1173*, *lisK* (*lmo1378*), *lmo1508*, *lmo1741*, *resE* (*lmo1947*) and *cesK* (*lmo2421*) displayed 1.6- to 3-fold ($P < 0.05$) upregulation (Fig. 2). At each time point post-cold shock from 37 °C to 5 °C, *lmo0050*, *yycG* (*lmo0288*), *lmo1061*, *lmo1741* and *kdpD* (*lmo2679*) were upregulated 1.9- to 3.4-fold ($P < 0.05$) as compared to *t₀* (Figs 3A–C). *liaS* and *phoR* (*lmo2500*) were upregulated 1.5- to 2.1-fold ($P < 0.05$) both at 30 min and 3 h (Figs 3A and B). Cold shock induced the expression of *cheA* 7.8-fold ($P < 0.01$) at 7 h (Fig. 3C), and that of *lmo2011* 1.6-fold ($P < 0.05$) at 30 min (Fig. 3A). The expression of other HK-encoding genes was unaffected under cold stress.




Figure 2 Relative gene expression levels of the HK-encoding genes in *L. monocytogenes* EGD-e during the logarithmic growth phase at 3 °C in relation to the expression level in logarithmic growth at 37 °C. Gene expression was normalized to 16S *rrn*. Error bars represent the minimum and maximum ratios between three replicate cultures. Significantly increased relative expression levels ($P < 0.05$, < 0.01 or < 0.001 , paired *t*-test) are indicated by asterisks *, ** and ***, respectively (I).




Figure 3 Relative expression levels of the HK-encoding genes in *L. monocytogenes* EGD-e 30 min (A), 3 h (B) and 7 h (C) post-cold shock from 37 °C to 5 °C, each calibrated to t_0 at 37 °C. Gene expression was normalized to 16S *rrm*. Error bars represent the minimum and maximum ratios between three replicate cultures. Significantly increased relative expression levels ($P < 0.05$, < 0.01 or < 0.001 , paired *t*-test) are indicated by asterisks *, ** and ***, respectively (I).

5.1.2 Growth of genetically modified *L. monocytogenes* under various stresses (I, II)

At 3 °C, the growth of the $\Delta lisK$ deletion mutant strain was nearly completely restricted, with the growth rate 86% ($P < 0.001$) lower and the maximum optical density 73% ($P < 0.001$) lower than those of the wild-type EGD-e strain. $\Delta yycG$ displayed a prolonged lag phase, a 31% ($P < 0.001$) lower growth rate and an 8% ($P < 0.01$) lower maximum optical density compared to the wild type. $\Delta resE$ displayed a later onset of growth and a 29% ($P < 0.001$) lower growth rate than those of the wild type (I). Complementation of $\Delta yycG$ restored the cold-sensitive phenotype to the wild-type level, while complementation of $\Delta lisK$ did not fully restore growth at low temperature, although a slight increase in the maximum optical density was observed (I).

The growth of $\Delta liaS$ was impaired under each stress condition other than low temperature when compared to EGD-e, with marked changes at high temperature (42.5 °C) and under osmotic stress (6% NaCl), moderate changes under pH stresses (5.6 and 9.4) and with minor changes in 3.5% ethanol and 5mM H₂O₂. AUC values of $\Delta liaS$ were 31% and 30% lower ($P < 0.01$) under osmotic and heat stresses, respectively, 21% and 15% lower ($P < 0.001$) under acid and alkali stresses, respectively, and 11% lower ($P < 0.01$) under both ethanol and oxidative stresses compared to AUC values of EGD-e under corresponding stress conditions. The maximum growth rates of $\Delta liaS$ decreased by 67% ($P < 0.001$) in 6% NaCl and 15% to 46% ($P < 0.01$) under other stresses. $\Delta virS$ displayed almost completely restricted growth at 42.5 °C and moderately impaired growth in ethanol. The AUC values of $\Delta virS$ were 58% and 19% lower ($P < 0.01$) under heat and ethanol stresses, respectively. The maximum growth rate of $\Delta virS$ in ethanol was 33% lower ($P < 0.01$), while its growth rate at high temperature was negligible. The $\Delta agrC$ strain displayed notably impaired growth under osmotic stress, with a 33% lower ($P < 0.01$) maximum growth rate. $\Delta agrC$ also displayed slight growth deficiency and a 14% lower ($P < 0.01$) AUC value under oxidative stress (II). Complementation restored the wild-type phenotypes to the $\Delta agrC$, $\Delta liaS$ and $\Delta virS$ mutants (II). At 37 °C, the growth of none of the mutant strains deviated from that of the wild-type EGD-e (I, II).

Table 5. Mean maximum growth rates of *L. monocytogenes* wild-type EGD-e and HK mutant strains. Modified from I and II.

Strain	Maximum growth rate \pm SD (OD ₆₀₀ units/h) ^a						
	4 °C	42.5 °C	pH 5.6	pH 9.4	6% NaCl	3.5% ethanol	5 mM H ₂ O ₂
EGD-e	0.0035 \pm 0.0001	0.059 \pm 0.016	0.068 \pm 0.001	0.13 \pm 0.004	0.006 \pm 0.0000	0.06 \pm 0.006	0.22 \pm 0.007
Δ Imo0050 (Δ agrC)	0.0057 \pm 0.0005	0.068 \pm 0.030	0.072 \pm 0.001	0.12 \pm 0.003*	0.004 \pm 0.0000*	0.05 \pm 0.005	0.18 \pm 0.004
Δ Imo0288 (Δ ycyG)	0.0024 \pm 0.0003**	0.065 \pm 0.013	0.062 \pm 0.001*	0.20 \pm 0.011	0.007 \pm 0.0000	0.04 \pm 0.003	0.23 \pm 0.008
Δ Imo0692 (Δ cheA)	0.0033 \pm 0.0001	0.056 \pm 0.015	0.063 \pm 0.001**	0.12 \pm 0.003	0.006 \pm 0.0000	0.05 \pm 0.004	0.21 \pm 0.006
Δ Imo1021 (Δ liaS)	0.0055 \pm 0.0007	0.050 \pm 0.015	0.043 \pm 0.001**	0.07 \pm 0.002**	0.002 \pm 0.0000**	0.04 \pm 0.003*	0.17 \pm 0.004**
Δ Imo1061	0.0050 \pm 0.0004	0.060 \pm 0.014	0.062 \pm 0.001**	0.13 \pm 0.004	0.006 \pm 0.0000	0.05 \pm 0.004	0.22 \pm 0.007
Δ Imo1173	0.0044 \pm 0.0003	0.058 \pm 0.014	0.062 \pm 0.001*	0.13 \pm 0.004	0.008 \pm 0.0001	0.05 \pm 0.005	0.22 \pm 0.007
Δ Imo1378 (Δ lisk)	0.0005 \pm 0.0003**	0.058 \pm 0.023	0.065 \pm 0.001	0.16 \pm 0.005	0.009 \pm 0.0001	0.07 \pm 0.008	0.22 \pm 0.007
Δ Imo1508	0.0056 \pm 0.0005	0.062 \pm 0.014	0.063 \pm 0.001*	0.24 \pm 0.009	0.008 \pm 0.0001	0.06 \pm 0.005	0.22 \pm 0.008
Δ Imo1741 (Δ virS)	0.0044 \pm 0.0002	0.000 \pm 0.003*	0.066 \pm 0.002*	0.16 \pm 0.006	0.007 \pm 0.0000	0.04 \pm 0.003*	0.24 \pm 0.008
Δ Imo1947 (Δ resE)	0.0025 \pm 0.0001**	0.051 \pm 0.014	0.061 \pm 0.001*	0.13 \pm 0.004	0.008 \pm 0.0001	0.05 \pm 0.004	0.21 \pm 0.007
Δ Imo2011	0.0036 \pm 0.0001	0.051 \pm 0.010	0.065 \pm 0.001*	0.13 \pm 0.004	0.009 \pm 0.0001	0.05 \pm 0.005	0.22 \pm 0.008
Δ Imo2421 (Δ cesK)	0.0036 \pm 0.0002	0.054 \pm 0.015	0.065 \pm 0.001	0.13 \pm 0.005	0.009 \pm 0.0001	0.06 \pm 0.009	0.21 \pm 0.006
Δ Imo2500 (Δ phoR)	0.0032 \pm 0.0002	0.050 \pm 0.013	0.064 \pm 0.001	0.21 \pm 0.007	0.009 \pm 0.0002	0.05 \pm 0.005	0.18 \pm 0.004
Δ Imo2582	0.0040 \pm 0.0002	0.073 \pm 0.026	0.069 \pm 0.001	0.12 \pm 0.003*	0.007 \pm 0.0001	0.05 \pm 0.005	0.21 \pm 0.006
Δ Imo2679 (Δ kdpD)	0.0079 \pm 0.0007	0.064 \pm 0.025	0.068 \pm 0.001	0.13 \pm 0.004	0.006 \pm 0.0000	0.05 \pm 0.004	0.22 \pm 0.007

^a Significantly decreased values ($P < 0.01$, $P < 0.001$, 2-tailed t -test) compared to the corresponding value of the wild type are indicated by asterisks * and **, respectively.

5.1.3 Minimum and maximum growth temperatures (I, II)

The minimum growth temperatures of $\Delta yycG$ and $\Delta lisK$ were 3.7 °C and 2.7 °C higher ($P < 0.001$), respectively, when compared to EGD-e after a 21-day incubation, while $\Delta cesK$ and $\Delta kdpD$ presented 2.3 °C and 0.5 °C ($P < 0.05$) higher minimum growth temperatures, respectively (I). The maximum growth temperature of $\Delta liaS$ was 0.9 °C lower ($P < 0.001$) and that of $\Delta virS$ 0.7 °C lower ($P < 0.001$) than EGD-e (Fig. 4) (II).




Figure 4 Maximum growth temperatures of *L. monocytogenes* EGD-e, the histidine kinase deletion mutant strains $\Delta liaS$ and $\Delta virS$ and the complemented strains $\Delta liaSc$ and $\Delta virSc$ within 48 h in the Gradiplate W10 incubator in a temperature gradient of 39 °C to 44 °C (II).

5.2 Plasmid-mediated heat resistance in *L. monocytogenes* (III)

5.2.1 Novel 58-kb plasmid contributes to the heat resistance of *L. monocytogenes* AT3E (III)

Genome analysis revealed the heat-resistant AT3E to harbour a novel plasmid pLM58 (Fig. 5). The 58-kb plasmid, with a GC content of 36.6%, contains 70 predicted ORFs and 19 predicted operons. Plasmid curing resulted in significantly impaired heat resistance of the derivative AT3Epc compared to the AT3E parent, with a log₁₀ reduction of 1.1 cfu/ml ($P < 0.001$) at 55 °C (Fig. 6).




Figure 5 Genetic map of *L. monocytogenes* plasmid pLM58. Initiation of replication protein-encoding *ori*, ATP-dependent protease-encoding *cplA*, cadmium resistance genes *cadA* and *cadC*, transposon Tn5422 and unique restriction sites are indicated (III).

5.2.2 Plasmid-borne *cplA* increases heat resistance in *L. monocytogenes* (III)

Annotation of pLM58 revealed an ORF putatively encoding an ATP-dependent ClpL protease (Fig. 5). Introduction of *cplA* augmented the heat resistance of natively heat-sensitive *L. monocytogenes* 10403S, observed by a significant decrease in \log_{10} reduction from 1.2 cfu/ml to 0.4 cfu/ml ($P < 0.01$) at 55 °C (Fig. 6).




Figure 6 pLM58 plasmid-borne ATP-dependent protease ClpL mediates heat resistance in *L. monocytogenes*. Error bars represent the standard deviations of three replicate cultures. Statistically significant \log_{10} reduction values ($P < 0.01$, $P < 0.001$, independent samples 2-tailed *t*-test) are indicated by asterisks * and **, respectively. NS, not significant. (III)

5.2.3 pLM58 is putatively nonconjugative (III)

Upon standard plate mating between AT3E and 10403S, no colonies were detected on selective plates containing both 130 µg/ml CdSO₄ and 200 µg/ml streptomycin, indicating no conjugative transmission of pLM58. pLM58 also lacked the known T4SS-encoding genes required for the conjugation of self-transmissible plasmids.

5.3 *L. monocytogenes* strains display high variability under severe stresses (III, IV)

5.3.1 Resistance and growth of *L. monocytogenes* strains vary at high temperatures (III)

We first tested the heat resistance at 55 °C, growth at 42 °C and maximum growth temperature of *L. monocytogenes* AL4E and AT3E in order to elucidate the differences between their thermoresistance and growth at high temperature. With a 0.0 cfu/ml log₁₀ reduction, *L. monocytogenes* AT3E proved to be more heat resistant than AL4E (1.4 cfu/ml log₁₀ reduction; $P < 0.01$) at 55 °C. At 42°C, the differences between their growth were negligible. Strain AL4E exhibited a 0.5 °C higher maximum growth temperature than the heat-resistant AT3E strain ($P < 0.01$).

5.3.2 *L. monocytogenes* isolates display variable resistance to benzalkonium chloride (IV)

The BC resistance profiles for the Finnish and Swiss *L. monocytogenes* isolates were determined. BC MICs ranged from 7.5 µg/ml to over 30 µg/ml. Isolates with BC MIC of ≥ 20 µg/ml were considered BC^r. Altogether, 11.5% (45/392) of the strains proved to be resistant to BC. Considering the isolation country, 10.7% (21/107) and 12.3% (24/195) of isolates of Finnish and Swiss origin strains, respectively, were BC^r.

Overall, most of the BC^r strains belonged to lineage II (39/45; 86.7%). Considering serotypes, the highest prevalence of BC^r strains was found among serotype 1/2c (11/34; 32.4%), while, in total, most of the resistant strains belonged to the serotype 1/2a (28/45, 62.2%). Genotypically, the 45 BC^r strains belonged to 14 MLSTs, of which two were the newly described ST25 and ST28. Most of the BC^r strains were of ST121 (14/45; 31.1%) and ST9 (11/45; 24.4%). Furthermore, most of the BC^r strains were associated with clonal complexes (CC) CC121 (16/45; 35.6%) and CC9 (11/45; 24.4%).

PCR-based analysis revealed 68.9% (31/45) of the BC^r strains to harbour at least one of the known BC resistance-conferring genes *bcrABC*, *emrE* or *qacH*. *bcrABC* was found in Swiss isolates alone, while only one BC^r isolate, that of Finnish origin, was found to harbour *emrE*. A screen with reserpine

showed that 40% (18/45) of the BC^r strains were at least partially efflux pump-dependent. Of the 14 strains lacking the known BC resistance-conferring genes, the BC resistance of three strains proved to be independent of efflux pump systems.

6 DISCUSSION

6.1 Extensive roles of two-component-system histidine kinases in the stress tolerance of *L. monocytogenes* (I–II)

In these studies we assessed the roles of HK-encoding genes in the stress tolerance of *L. monocytogenes* to ambient stress conditions, low and high temperatures, low and high pH levels, and high salt, ethanol and hydrogen peroxide concentrations, widely used to control bacterial contamination in food processing and preservation (NicAogáin and O'Byrne, 2016, Bucur *et al.*, 2018). While RRs of *L. monocytogenes* have been more extensively studied, information has been lacking on the role of HKs. However, phosphorylation of the HK is a key step in the phosphotransfer chain that further governs the phosphorylation of the cognate RR and the final output (Stock *et al.*, 2000, Jacob-Dubuisson *et al.*, 2018), and the functions of HKs are thus of special interest.

Of the HKs, YycG and LisK proved to be essential to the cold tolerance of *L. monocytogenes*. Based on expressional analyses of HK-encoding genes, YycG seems to be particularly needed under abrupt cold stress, while the role of LisK is accentuated during cold acclimation. In addition, ResE appears to play a minor role in the early growth phase of *L. monocytogenes* under cold conditions. The contribution of all of these TCSs to cold tolerance might be explained by potential cell envelope-related stress responses (Mohedano *et al.*, 2005, Williams *et al.*, 2005a, Gottschalk *et al.*, 2008, Nielsen *et al.*, 2012). The YycFG system (synonymous for WalkR) is regarded as essential and is highly conserved among low G+C-content Gram-positive bacteria such as *B. subtilis* (Fabret and Hoch, 1998, Szurmant *et al.*, 2007, Takada *et al.*, 2018) and *S. aureus* (Martin *et al.*, 1999, Poupel *et al.*, 2016). Due its essential and conserved nature, YycFG has attracted long-lasting research interest as a potential antimicrobial target (Fabret and Hoch, 1998, Dubrac *et al.*, 2008, Tiwari *et al.*, 2017).

We concluded HK kinase LiaS to have a universal impact on the stress tolerance of *L. monocytogenes* EGD-e. Mutational studies revealed impaired growth of $\Delta liaS$ under each stress condition studied except cold: under heat, acid, alkali, osmotic, ethanol, and oxidative conditions. The most prominent roles were noted upon heat and osmotic stresses at 42.5 °C and in 6% NaCl, respectively. Since the upkeep of the cell wall structure and membrane functions have a key role in environmental stress adaptation (Soni *et al.*, 2011, NicAogáin and O'Byrne, 2016), the comprehensive role of the TCS LiaSR may be attributable to its actions of remodelling the cytoplasmic membrane composition in response to cell-envelope disturbances (Fritsch *et al.*, 2011).

Besides LiaSR-mediated stress responses, our studies, combined with previous findings on RRs YycF (Chan *et al.*, 2007b), LisR (Chan *et al.*, 2008), VirR (Williams *et al.*, 2005a) and AgrA (Garmyn *et al.*, 2012), point out that the complete TCSs YycFG and LisRK, AgrCA and VirRS play roles under cold, osmotic and ethanol stresses, respectively. The potential versatility in stress tolerance of the Agr quorum sensing system, which includes AgrCA, is illustrated by the advantage that it provides *L. monocytogenes* in survival in natural soil, the most complex growth environment (Vivant *et al.*, 2014, Vivant *et al.*, 2015). VirRS and the Agr quorum sensing system have also been associated with the virulence of *L. monocytogenes* (Autret *et al.*, 2003, Mandin *et al.*, 2005, Riedel *et al.*, 2009, de las Heras *et al.*, 2011, Grubaugh *et al.*, 2018), with VirR playing a key role as a virulence regulator and the *virR* gene itself being upregulated *in vivo* during infection in a mouse model (Camejo *et al.*, 2009). VirR potentially participates in virulence by gene regulation of cell-surface modulation (Mandin *et al.*, 2005, Camejo *et al.*, 2009). Alterations in cell-envelope structures were, in turn, suggested to result in enhanced ethanol tolerance (Williams *et al.*, 2005a). Furthermore, exposure to osmotic stress similar to conditions in the human intestine increases virulence gene expression in *L. monocytogenes* (Sue *et al.*, 2004). Due to these potential interfaces between virulence and stress tolerance mechanisms, it may not be surprising that both TCSs AgrCA and VirRS would be involved not only in virulence but also in stress tolerance in *L. monocytogenes*. However, it should be further noticed that, instead of the toxin production-controlling VirRS system, extensively described in *Clostridium perfringens* (Lyrstis *et al.*, 1994, Ohtani *et al.*, 2010) in which the regulatory function occurs via RNA molecules (Shimizu *et al.*, 2002), VirRS of *L. monocytogenes* is a member of the BceRS-like TCSs (Dintner *et al.*, 2011).

Intriguingly, cognate TCS partners do not always seem to feature together in the same stress tolerance responses. Besides the obvious methodological differences, as well as potential variation within parental strains between studies (Bécavin *et al.*, 2014), cross-phosphorylation (cross-talk or cross-regulation) may take place between noncognate partners (Laub and Goulian, 2007, Siryaporn and Goulian, 2010). Indeed, while we found LiaS to play a significant role in heat stress, the same is evidently not the case with its cognate RR LiaR (Kallipolitis and Ingmer, 2001, Williams *et al.*, 2005a). LiaS is a potential phosphorylator of the orphan RR DegU (Williams *et al.*, 2005a), also assigned a role in contributing to the heat tolerance of *L. monocytogenes* (Gueriri *et al.*, 2008). Mandin *et al.* (2005) also suggested that, inside a host, VirR activation might be unrelated to the cognate HK VirS but occur via cross-talking or molecular compound phosphorylation instead. Verifying cross-phosphorylation between noncognate partners would require further investigations into protein–protein interactions between particular HKs and RRs (Bielecki *et al.*, 2015).

For those HK-encoding genes upregulated at low temperatures while presenting no significant changes in mutational studies, the apparent discrepancies may be due to putative posttranscriptional regulation. For instance, Csp-encoding genes are mainly regulated at the posttranscriptional level (Narberhaus *et al.*, 2006). Overall, we found that, apart from the chemotaxis-encoding *cheA*, the relative expression levels of the HK-encoding genes at low temperatures only moderately differed from that of the expression levels at 37 °C. Although a recent comparative transcriptome analysis demonstrated fairly high transcript levels (8.9-fold change) for HK-encoding *kdpD* upon sodium lactate adaptation (Suo *et al.*, 2018), as such, transcriptional studies may not be the optimum single method of choice to unambiguously ascertain the role of a certain HK in the stress tolerance of *L. monocytogenes*.

Although the presence and the output domain types of the TCSs of *L. monocytogenes* have been described (Glaser *et al.*, 2001, Williams *et al.*, 2005a), much remains to be discovered concerning the detailed structures and the triggering signals of the TCS signalling structures of *L. monocytogenes* (Zetmann *et al.*, 2016, Pinheiro *et al.*, 2018). Furthermore, by collecting phenotypic data with a considerably larger number of strains than in the present studies, the discrepancies between the putative roles of certain TCSs reported in previous studies could possibly be elucidated. Strain and population heterogeneity might influence the results when using only one or few model strains, particularly when using laboratory strains that do not always optimally represent the naturally occurring *L. monocytogenes* strains (Bécavin *et al.*, 2014).

6.2 pLM58-borne ClpL-mediated heat resistance in *L. monocytogenes* (III)

This study is very plausibly the first description of plasmid-mediated heat resistance in *L. monocytogenes*, conferred via a novel plasmid, pLM58. Genes potentially related to heat tolerance have been annotated in *L. monocytogenes* plasmids (Fox *et al.*, 2016). Canchaya *et al.* (2010) also identified a *clpL* gene that was suggested to originate from lactic acid bacteria, in which *clpL* was upregulated during heat shock (Suokko *et al.*, 2005). However, conclusive evidence of plasmid-borne heat resistance-mediating genes in *L. monocytogenes* has been lacking to date. Our results indicate that plasmid-borne ClpL is a potential predictor of augmented heat resistance in *L. monocytogenes*. As heat treatment is a central means to control *L. monocytogenes* contamination during food processing (Bucur *et al.*, 2018), the possibility of a heat resistance-conferring trait harboured by a MGE in a foodborne pathogen raises serious concerns with regard to food hygiene and safety. Of other pathogenic bacteria, a heat resistance-mediating plasmid has previously been described at least in *Klebsiella pneumoniae*

(Bojer *et al.*, 2010, Bojer *et al.*, 2012). Regarding plasmid-mediated traits within bacteria, the focus of the past and present research has been on the rapidly emerging antibiotic resistance (Poyart-Salmeron *et al.*, 1990, Liu *et al.*, 2016, Liu *et al.*, 2019, Schweizer *et al.*, 2019, Slettemeås *et al.*, 2019), while much less is known about the plasmid-mediated environmental stress tolerance of pathogenic bacteria. However, the possible cotransfer of both stress tolerance and antibiotic resistance-conferring traits cannot be overlooked, as they may have grave implications for consumer and patient safety considering both foodborne and nosocomial infections. Notably, Bojer *et al.* (2012) detected the cotransfer of both multidrug and heat resistance-conferring traits in a conjugative plasmid in *K. pneumoniae*.

Instead of focusing on particular genes, our aim in this study was to comprehensively approach the subject of finding novel genetic traits conferring heat resistance in *L. monocytogenes*. Furthermore, we wanted to take into account the possibility of plasmids acting as heat-resistance mediators. Consequently, WGS and comparison of the complete genomes of a heat-resistant and heat-sensitive strain were opted for. Indeed, whole-genome comparison revealed the most prominent difference between the heat-resistant AT3E and heat-sensitive AL4E *L. monocytogenes* strains to be a novel 58-kb plasmid. Furthermore, plasmid curing generated a significant decrease in heat resistance of the AT3E strain at 55 °C. pLM58 harboured a 2,115-bp ATP-dependent protease-encoding *clpL* gene. Introducing this gene to a naturally heat-sensitive *L. monocytogenes* 10403S resulted, in turn, in significantly improved survival under heat challenge. While chromosomal stress resistance-mediating Clp ATPases have been described in *L. monocytogenes* (Rouquette *et al.*, 1996, Nair *et al.*, 2000), this is the first study to identify a plasmid-borne Clp in *L. monocytogenes* conferring heat resistance. As the same *clpL* gene can be found in other *Listeria* spp. plasmids sequenced to date, heat resistance may prove to be even more prevalent within plasmid-harbouring *L. monocytogenes* strains.

As plate mating resulted in no apparent transconjugants, it seems that pLM58 is not self-transmissible. This was supported by the absence of the known type IV secretion system-encoding genes (Smillie *et al.*, 2010, Goessweiner-Mohr *et al.*, 2014) in pLM58. Whether pLM58 is mobilisable needs to be further investigated. However, due to the mosaic nature of *L. monocytogenes* plasmids (Canchaya *et al.*, 2010, Schmitz-Esser *et al.*, 2015), the possibility of a conjugative plasmid harbouring a heat resistance-mediating *clpL* gene in *L. monocytogenes* still remains. As WGS methods become increasingly common and affordable and as complete sequences of new *Listeria* spp. plasmids become available in open databases, future comparative genomic analyses will undoubtedly enlighten the contents and potential heat resistance-mediating genes harboured by other, possibly conjugative plasmids.

Neither plasmid curing nor introducing the *clpL* into a heat-sensitive *L. monocytogenes* strain resulted in alterations in the maximum growth

temperature or kinetic growth parameters at 42 °C. This may indicate different mechanisms for resistance to thermal kill than the ability to grow at high temperatures. Indeed, similar indications have been found in *K. pneumoniae*, as mutation in the heat-resistance mediating gene did not affect the maximum growth temperature (Bojer *et al.*, 2010). Consequently, in experimental designs it may also be important to differentiate between the ability to grow under a certain stress from the ability to resist and survive a harsher stress condition.

In this study, we were able to pinpoint a causative gene carried on a plasmid by comparing two complete genomes and combining the genomic information with traditional molecular experiments. However, a larger sample size would greatly facilitate the prospects of finding completely novel mechanisms conferring stress resistance: a large enough sample size – from hundreds to preferably even thousands – would allow genome-wide association studies (GWAS) methods to be employed that could identify not only complete plasmids or genes but also smaller variants, such as SNPs and indels, associated with certain phenotypes (Lees *et al.*, 2016, Kachroo *et al.*, 2019). It should be noted, however, that even in the case of a proven statistical association, traditional molecular methods are still needed to ascertain true causality (Read and Massey, 2014, Kremer *et al.*, 2017).

The findings in this study raise concern, amongst others, due to the potential cotransfer of plasmid-borne stress resistance-mediating genes together with those conferring resistance to virulence and antibiotics (Bojer *et al.*, 2010, Bojer *et al.*, 2012, Gullberg *et al.*, 2014, Fox *et al.*, 2016). This may provide bacteria with opportunities to populate different niches and to infect new hosts.

6.3 Characteristics of benzalkonium chloride resistance in *L. monocytogenes* (IV)

In this study, 392 *L. monocytogenes* isolates of Finnish and Swiss origin were analysed with respect to BC resistance. Altogether, 11.5% of the strains proved to be BC^r, corresponding with the lower estimates of previous findings in isolates from food-processing environments and foodstuffs (Aase *et al.*, 2000, Mullapudi *et al.*, 2008, Jiang *et al.*, 2016). It should be noted, however, that the definition of BC resistance has differed markedly between studies (Aase *et al.*, 2000, Mereghetti *et al.*, 2000, Xu *et al.*, 2014) due to the often-employed method of using as a threshold the MIC that prevents the growth of the majority of the test strains themselves. However, with large and variable enough sample material, the relative cut-off approach should provide a sufficient basis for selecting the clearly resistant strains. Essentially, these strains would also be the ones most likely to be problematic in food processing premises with regard to the successful disinfection of processing equipment and surfaces. Most worryingly, it has been suggested

that adaptation to BC may also augment antibiotic resistance within bacteria, as was shown within *Pseudomonas* spp. isolates (Tandukar *et al.*, 2013, Kim *et al.*, 2018). In *L. monocytogenes*, adaptation to BC not only enhanced tolerance to subsequent exposure to BC, but also resistance to antimicrobial agents such as cefotaxime, cephalothin and ciprofloxacin (Yu *et al.*, 2018). These outcomes suggest that the use of disinfectants should be well planned and apportioned to avoid the excess use of possibly multi-resistance-selecting agents at futile sites.

No clear correlation between BC resistance and the isolation country or isolation source was found in this study. Considering the serotypes, most of the BC^r strains were of serotype 1/2c, while in total, serotype 1/2a included the highest number of BC^r strains, which may reflect the overrepresentation of 1/2a strains in this study. However, a wide range of BC resistance among different serotypes has previously been reported: Mullanpudi *et al.* (2008) found BC^r *L. monocytogenes* strains to be significantly more predominant in serotype 1/2a and 1/2b than within 4b. A parallel distribution of BC^r strains had previously been reported by Mereghetti *et al.* (2000), while others detected no such correlation (Romanova *et al.*, 2002, Soumet *et al.*, 2005, Jiang *et al.*, 2016) and Xu *et al.* (2014), indeed, found an opposite BC resistance pattern among serotypes. It should be noted that the discrepancies might often be explained by the relatively restricted sample sizes or uneven representation of different serotypes among the study strains.

Screening of the known BC resistance-conferring genes *bcrABC* (Elhanafi *et al.*, 2010), *emrE* (Gilmour *et al.*, 2010, Kovacevic *et al.*, 2016) and *qacH* (Müller *et al.*, 2013) revealed fourteen strains lacking these genes altogether, indicating novel mechanisms for BC resistance in *L. monocytogenes*. Furthermore, reserpine screening indicated that in eleven of these strains, at least partially an efflux pump-dependent mechanism, other than those encoded by the known BC resistance-mediating genes, is putatively present. In the three strains both lacking the BC resistance-mediating genes and displaying independence of efflux pumps, a completely novel mechanism of BC resistance is suggested. Indeed, upregulation of peptidoglycan synthesis pathways in the presence of a QAC, benzethonium chloride, was seen in *L. monocytogenes* (Fox *et al.*, 2011), suggesting a potential role of cell-wall modifications in increased tolerance to BC (McDonnell and Russell, 1999). In future research, association study methods would be a noteworthy tool in screening for the potentially completely novel genetic variants putatively conferring BC resistance in *L. monocytogenes*.

7 CONCLUSIONS

1. HKs extensively contribute to stress tolerance in *L. monocytogenes*. YycG and LisK are needed upon an abrupt decline in temperature and cold acclimation, respectively, while LiaS plays a universal role in the tolerance of *L. monocytogenes* to high temperature, acid, alkali, osmotic, ethanol and oxidative stresses. Furthermore, the complete TCSs AgrCA and VirRS play roles under osmotic and ethanol stresses, respectively, and complete YycFG and LisRK under cold conditions. The present studies also revealed indications of potential cross-phosphorylation between noncognate TCS partners of *L. monocytogenes* under ambient stresses.
2. A novel plasmid, pLM58, confers heat resistance in *L. monocytogenes*. Furthermore, the plasmid-borne ATP-dependent protease ClpL contributes to the survival of *L. monocytogenes* at high temperatures, adding to the concern of resistance-mediating genes efficiently spreading by horizontal gene transfer among bacterial populations.
3. Emerging BC resistance is a potential concern in *L. monocytogenes* strains of Finnish and Swiss origin. BC resistance in strains lacking the so far known BC resistance-mediating genes indicates the existence of novel BC resistance mechanisms.

REFERENCES

- Aase, B., Sundheim, G., Langsrud, S. & Rørvik, L. M. (2000) Occurrence of and a possible mechanism for resistance to a quaternary ammonium compound in *Listeria monocytogenes*. *Int J Food Microbiol* 62: 57-63.
- Abajy, M. Y., Kopeć, J., Schiwon, K., Burzynski, M., Döring, M., Bohn, C. & Grohmann, E. (2007) A type IV-secretion-like system is required for conjugative DNA transport of broad-host-range plasmid pIP501 in gram-positive bacteria. *J Bacteriol* 189: 2487-2496.
- Abee, T., Koomen, J., Metselaar, K. I., Zwietering, M. H. & den Besten, H. M. W. (2016) Impact of pathogen population heterogeneity and stress-resistant variants on food safety. *Annu Rev Food Sci Technol* 7: 439-456.
- Abram, F., Starr, E., Karatzas, K. A. G., Matlawska-Wasowska, K., Boyd, A., Wiedmann, M., Boor, K. J., Connally, D. & O'Byrne, C. P. (2008) Identification of components of the Sigma B regulon in *Listeria monocytogenes* that contribute to acid and salt tolerance. *Appl Environ Microbiol* 74: 6848-6858.
- Aguilar, P. S., Hernandez-Arriaga, A. M., Cybulski, L. E., Erazo, A. C. & de Mendoza, D. (2001) Molecular basis of thermosensing: a two-component signal transduction thermometer in *Bacillus subtilis*. *EMBO J* 20: 1681-1691.
- Alikhan, N.-F., Petty, N. K., Ben Zakour, N. L. & Beatson, S. A. (2011) BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* 12: 402.
- Allam, M., Tau, N., Smouse, S. L., Mtshali, P. S., Mnyameni, F., Khumalo, Z. T. H., Ismail, A., Govender, N., Thomas, J. & Smith, A. M. (2018) Whole-genome sequences of *Listeria monocytogenes* sequence type 6 isolates associated with a large foodborne outbreak in South Africa, 2017 to 2018. *Genome Announc* 6: e00538-18.
- Almeida, G., Magalhães, R., Carneiro, L., Santos, I., Silva, J., Ferreira, V., Hogg, T. & Teixeira, P. (2013) Foci of contamination of *Listeria monocytogenes* in different cheese processing plants. *Int J Food Microbiol* 167: 303-309.
- Altamirano-Silva, P., Meza-Torres, J., Castillo-Zeledón, A., Ruiz-Villalobos, N., Zuñiga-Pereira, A. M., Chacón-Díaz, C., Moreno, E., Guzmán-Verri, C. & Chaves-Olarte, E. (2018) *Brucella abortus* senses the intracellular environment through the BvrR/BvrS two-component system, which allows *B. abortus* to adapt to its replicative niche. *Infect Immun* 86: e00713-17.
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389-3402.
- Amezaga, M. R., Davidson, I., Mclaggan, D., Verheul, A.,

- Abee, T. & Booth, I. R. (1995) The role of peptide metabolism in the growth of *Listeria monocytogenes* ATCC 23074 at high osmolarity. *Microbiology* 141: 41-49.
- Angelidis, A. S. & Smith, G. M. (2003) Role of the glycine betaine and carnitine transporters in adaptation of *Listeria monocytogenes* to chill stress in defined medium. *Appl Environ Microbiol* 69: 7492-7498.
- Angelo, K. M., Conrad, A. R., Saupe, A., Dragoo, H., West, N., Sorenson, A., Barnes, A., Doyle, M., Beal, J., Jackson, K. A., Stroika, S., Tarr, C., Kucerova, Z., Lance, S., Gould, L. H., Wise, M. & Jackson, B. R. (2017) Multistate outbreak of *Listeria monocytogenes* infections linked to whole apples used in commercially produced, prepackaged caramel apples: United States, 2014-2015. *Epidemiol Infect* 145: 848-856.
- Annous, B. A., Becker, L. A., Bayles, D. O., Labeda, D. P. & Wilkinson, B. J. (1997) Critical role of anteiso-C₁₅:0 fatty acid in the growth of *Listeria monocytogenes* at low temperatures. *Appl Environ Microbiol* 63: 3887-3894.
- Appleby, J. L., Parkinson, J. S. & Bourret, R. B. (1996) Signal transduction via the multi-step phosphorelay: not necessarily a road less traveled. *Cell* 86: 845-848.
- Archambaud, C., Nahori, M.-A., Pizarro-Cerdá, J., Cossart, P. & Dussurget, O. (2006) Control of *Listeria* superoxide dismutase by phosphorylation. *J Biol Chem* 281: 31812-31822.
- Arnaud, M., Chastanet, A. & Débarbouillé, M. (2004) New vector for efficient allelic replacement in naturally nontransformable, low-GC-content, gram-positive bacteria. *Appl Environ Microbiol* 70: 6887-6891.
- Arndt, D., Grant, J. R., Marcu, A., Sajed, T., Pon, A., Liang, Y. & Wishart, D. S. (2016) PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44: W16-W21.
- Aubin, G. G., Boutoille, D., Bourcier, R., Caillon, J., Lepelletier, D., Bémer, P. & Corvec, S. (2016) Unusual case of spondylodiscitis due to *Listeria monocytogenes*. *J Bone Jt Infect* 1: 7-9.
- Audurier, A., Chatelain, R., Chalons, F. & Piéchaud, M. (1979) [Bacteriophage typing of 823 "*Listeria monocytogenes*" strains isolated in France from 1958 to 1978 (author's transl)]. *Ann Microbiol (Paris)* 130B: 179-189.
- Aureli, P., Fiorucci, G. C., Caroli, D., Marchiaro, G., Novara, O., Leone, L. & Salmaso, S. (2000) An outbreak of febrile gastroenteritis associated with corn contaminated by *Listeria monocytogenes*. *N Engl J Med* 342: 1236-1241.
- Autio, T., Hielm, S., Miettinen, M., Sjöberg, A.-M., Aarnisalo, K., Björkroth, J., Mattila-Sandholm, T. & Korkeala, H. (1999) Sources of *Listeria monocytogenes* contamination in a cold-smoked rainbow trout processing plant detected by pulsed-field gel electrophoresis typing. *Appl Environ Microbiol* 65: 150-155.
- Autio, T., Markkula, A., Hellström, S., Niskanen, T., Lundén, J. & Korkeala, H. (2004) Prevalence and genetic

- diversity of *Listeria monocytogenes* in the tonsils of pigs. *J Food Prot* 67: 805-808.
- Autio, T., Säteri, T., Fredriksson-Ahomaa, M., Rahkio, M., Lundén, J. & Korkeala, H. (2000) *Listeria monocytogenes* contamination pattern in pig slaughterhouses. *J Food Prot* 63: 1438-1442.
- Autret, N., Raynaud, C., Dubail, I., Berche, P. & Charbit, A. (2003) Identification of the *agr* locus of *Listeria monocytogenes*: role in bacterial virulence. *Infect Immun* 71: 4463-4471.
- Aziz, R. K., Bartels, D., Best, A. A., Dejongh, M., Disz, T., Edwards, R. A., Formis, K., Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman, A. L., Overbeek, R. A., McNeil, L. K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G. D., Reich, C., Stevens, R., Vassieva, O., Vonstein, V., Wilke, A. & Zagnitko, O. (2008) The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9: 75.
- Bachiri, T., Lalaoui, R., Bakour, S., Allouache, M., Belkebla, N., Rolain, J. M. & Touati, A. (2017) First report of the plasmid-mediated colistin resistance gene *mcr-1* in *Escherichia coli* ST405 isolated from wildlife in Bejaia, Algeria. *Microb Drug Resist* 24: 890-895.
- Bae, D., Liu, C., Zhang, T., Jones, M., Peterson, S. N. & Wang, C. (2012) Global gene expression of *Listeria monocytogenes* to salt stress. *J Food Prot* 75: 906-912.
- Bakardjiev, A. I., Theriot, J. A. & Portnoy, D. A. (2006) *Listeria monocytogenes* traffics from maternal organs to the placenta and back. *PLoS Pathog* 2: e66.
- Baranyi, J. & Roberts, T. A. (1994) A dynamic approach to predicting bacterial growth in food. *Int J Food Microbiol* 23: 277-294.
- Barbosa, W. B., Cabedo, L., Wederquist, H. J., Sofos, J. N. & Schmidt, G. R. (1994) Growth variation among species and strains of *Listeria* in culture broth. *J Food Prot* 57: 765-769.
- Barbosa, W. B., Sofos, J. N., Schmidt, G. R. & Smith, G. C. (1995) Growth potential of individual strains of *Listeria monocytogenes* in fresh vacuum-packaged refrigerated ground top rounds of beef. *J Food Prot* 58: 398-403.
- Barker, C. & Park, S. F. (2001) Sensitization of *Listeria monocytogenes* to low pH, organic acids, and osmotic stress by ethanol. *Appl Environ Microbiol* 67: 1594-1600.
- Bayles, D. O., Annous, B. A. & Wilkinson, B. J. (1996) Cold stress proteins induced in *Listeria monocytogenes* in response to temperature downshock and growth at low temperatures. *Appl Environ Microbiol* 62: 1116-1119.
- Bayles, D. O. & Wilkinson, B. J. (2000) Osmoprotectants and cryoprotectants for *Listeria monocytogenes*. *Lett Appl Microbiol* 30: 23-27.
- Beales, N. (2004) Adaptation of microorganisms to cold temperatures, weak acid preservatives, low pH, and osmotic stress: a review. *Compr Rev Food Sci Food Saf* 3: 1-20.
- Beauregard, K. E., Lee, K.-D., Collier, R. J. & Swanson, J. A. (1997) pH-dependent

- perforation of macrophage phagosomes by listeriolysin O from *Listeria monocytogenes*. *J Exp Med* 186: 1159-1163.
- Bécavin, C., Bouchier, C., Lechat, P., Archambaud, C., Creno, S., Gouin, E., Wu, Z., Kühbacher, A., Brisse, S., Pucciarelli, M. G., García-del Portillo, F., Hain, T., Portnoy, D. A., Chakraborty, T., Lecuit, M., Pizarro-Cerdá, J., Moszer, I., Bierne, H. & Cossart, P. (2014) Comparison of widely used *Listeria monocytogenes* strains EGD, 10403S, and EGD-e highlights genomic differences underlying variations in pathogenicity. *MBio* 5: e00969-14.
- Becker, L. A., Çetin, M. S., Hutkins, R. W. & Benson, A. K. (1998) Identification of the gene encoding the alternative sigma factor σ^B from *Listeria monocytogenes* and its role in osmotolerance. *J Bacteriol* 180: 4547-4554.
- Becker, L. A., Evans, S. N., Hutkins, R. W. & Benson, A. K. (2000) Role of σ^B in adaptation of *Listeria monocytogenes* to growth at low temperature. *J Bacteriol* 182: 7083-7087.
- Beckerman, K. P., Rogers, H. W., Corbett, J. A., Schreiber, R. D., McDaniel, M. L. & Unanue, E. R. (1993) Release of nitric oxide during the T cell-independent pathway of macrophage activation. Its role in resistance to *Listeria monocytogenes*. *J Immunol* 150: 888-895.
- Begot, C., Lebert, I. & Lebert, A. (1997) Variability of the response of 66 *Listeria monocytogenes* and *Listeria innocua* strains to different growth conditions. *Food Microbiol* 14: 403-412.
- Beier, D. & Gross, R. (2006) Regulation of bacterial virulence by two-component systems. *Curr Opin Microbiol* 9: 143-152.
- Benson, A. K. & Haldenwang, W. G. (1993) The σ^B -dependent promoter of the *Bacillus subtilis* *sigB* operon is induced by heat shock. *J Bacteriol* 175: 1929-1935.
- Berche, P., Gaillard, J.-L. & Richard, S. (1988) Invasiveness and intracellular growth of *Listeria monocytogenes*. *Infection* 16: S145-S148.
- Bergholz, T. M., Bowen, B., Wiedmann, M. & Boor, K. J. (2012) *Listeria monocytogenes* shows temperature-dependent and -independent responses to salt stress, including responses that induce cross-protection against other stresses. *Appl Environ Microbiol* 78: 2602-2612.
- Bergholz, T. M., den Bakker, H. C., Fortes, E. D., Boor, K. J. & Wiedmann, M. (2010) Salt stress phenotypes in *Listeria monocytogenes* vary by genetic lineage and temperature. *Foodborne Pathog Dis* 7: 1537-1549.
- Berrang, M. E., Meinersmann, R. J., Northcutt, J. K. & Smith, D. P. (2002) Molecular characterization of *Listeria monocytogenes* isolated from a poultry further processing facility and from fully cooked product. *J Food Prot* 65: 1574-1579.
- Bertsch, D., Rau, J., Eugster, M. R., Haug, M. C., Lawson, P. A., Lacroix, C. & Meile, L. (2013) *Listeria fleischmannii* sp. nov., isolated from cheese. *Int J Syst Evol Microbiol* 63: 526-532.
- Bērziņš, A., Hellström, S., Siliņš, I. & Korkeala, H. (2010) Contamination patterns of

- Listeria monocytogenes* in cold-smoked pork processing. *J Food Prot* 73: 2103-2109.
- Betriu, C., Fuentemilla, S., Méndez, R., Picazo, J. J. & García-Sánchez, J. (2001) Endophthalmitis caused by *Listeria monocytogenes*. *J Clin Microbiol* 39: 2742-2744.
- Beumer, R. R., Te Giffel, M. C., Cox, L. J., Rombouts, F. M. & Abee, T. (1994) Effect of exogenous proline, betaine, and carnitine on growth of *Listeria monocytogenes* in a minimal medium. *Appl Environ Microbiol* 60: 1359-1363.
- Bibb, W. F., Gellin, B. G., Weaver, R., Schwartz, B., Plikaytis, B. D., Reeves, M. W., Pinner, R. W. & Broome, C. V. (1990) Analysis of clinical and food-borne isolates of *Listeria monocytogenes* in the United States by multilocus enzyme electrophoresis and application of the method to epidemiologic investigations. *Appl Environ Microbiol* 56: 2133-2141.
- Bibb, W. F., Schwartz, B., Gellin, B. G., Plikaytis, B. D. & Weaver, R. E. (1989) Analysis of *Listeria monocytogenes* by multilocus enzyme electrophoresis and application of the method to epidemiologic investigations. *Int J Food Microbiol* 8: 233-239.
- Bielecki, P., Jensen, V., Schulze, W., Gödeke, J., Strehmel, J., Eckweiler, D., Nicolai, T., Bielecka, A., Wille, T., Gerlach, R. G. & Häussler, S. (2015) Cross talk between the response regulators PhoB and TctD allows for the integration of diverse environmental signals in *Pseudomonas aeruginosa*. *Nucleic Acids Res* 43: 6413-6425.
- Bishop, D. K. & Hinrichs, D. J. (1987) Adoptive transfer of immunity to *Listeria monocytogenes*. The influence of *in vitro* stimulation on lymphocyte subset requirements. *J Immunol* 139: 2005-2009.
- Blériot, C., Dupuis, T., Jouvion, G., Eberl, G., Disson, O. & Lecuit, M. (2015) Liver-resident macrophage necroptosis orchestrates type 1 microbicidal inflammation and type-2-mediated tissue repair during bacterial infection. *Immunity* 42: 145-158.
- Boerlin, P., Rocourt, J., Grimont, F., Grimont, P. A., Jacquet, C. & Piffaretti, J.-C. (1992) *Listeria ivanovii* subsp. *londoniensis* subsp. nov. *Int J Syst Evol Microbiol* 42: 69-73.
- Bojer, M. S., Hammerum, A. M., Jørgensen, S. L., Hansen, F., Olsen, S. S., Krogfelt, K. A. & Struve, C. (2012) Concurrent emergence of multidrug resistance and heat resistance by CTX-M-15-encoding conjugative plasmids in *Klebsiella pneumoniae*. *APMIS* 120: 699-705.
- Bojer, M. S., Struve, C., Ingmer, H., Hansen, D. S. & Krogfelt, K. A. (2010) Heat resistance mediated by a new plasmid encoded Clp ATPase, ClpK, as a possible novel mechanism for nosocomial persistence of *Klebsiella pneumoniae*. *PLoS One* 5: e15467.
- Borezee, E., Msadek, T., Durant, L. & Berche, P. (2000) Identification in *Listeria monocytogenes* of MecA, a homologue of the *Bacillus subtilis* competence

- regulatory protein. *J Bacteriol* 182: 5931-5934.
- Bowman, J. P., Hages, E., Nilsson, R. E., Kocharunchitt, C. & Ross, T. (2012) Investigation of the *Listeria monocytogenes* Scott A acid tolerance response and associated physiological and phenotypic features via whole proteome analysis. *J Proteome Res* 11: 2409-2426.
- Bradshaw, J. G., Peeler, J. T., Corwin, J. J., Hunt, J. M., Tierney, J. T., Larkin, E. P. & Twedt, R. M. (1985) Thermal resistance of *Listeria monocytogenes* in milk. *J Food Prot* 48: 743-745.
- Braun, U., Stehle, C. & Ehrensperger, F. (2002) Clinical findings and treatment of listeriosis in 67 sheep and goats. *Vet Rec* 150: 38-42.
- Brul, S., Bassett, J., Cook, P., Kathariou, S., McClure, P., Jasti, P. R. & Betts, R. (2012) 'Omics' technologies in quantitative microbial risk assessment. *Trends Food Sci Technol* 27: 12-24.
- Brøndsted, L., Kallipolitis, B. H., Ingmer, H. & Knöchel, S. (2003) *kdpE* and a putative RsbQ homologue contribute to growth of *Listeria monocytogenes* at high osmolarity and low temperature. *FEMS Microbiol Lett* 219: 233-239.
- Bucur, F. I., Grigore-Gurgu, L., Crauwels, P., Riedel, C. U. & Nicolau, A. I. (2018) Resistance of *Listeria monocytogenes* to stress conditions encountered in food and food processing environments. *Front Microbiol* 9: 2700.
- Cabiscol, E., Tamarit, J. & Ros, J. (2000) Oxidative stress in bacteria and protein damage by reactive oxygen species. *Int Microbiol* 3: 3-8.
- Cacace, G., Mazzeo, M. F., Sorrentino, A., Spada, V., Malorni, A. & Siciliano, R. A. (2010) Proteomics for the elucidation of cold adaptation mechanisms in *Listeria monocytogenes*. *J Proteomics* 73: 2021-2030.
- Cai, S. J. & Inouye, M. (2002) EnvZ-OmpR interaction and osmoregulation in *Escherichia coli*. *J Biol Chem* 277: 24155-24161.
- Camejo, A., Buchrieser, C., Couvé, E., Carvalho, F., Reis, O., Ferreira, P., Sousa, S., Cossart, P. & Cabanes, D. (2009) *In vivo* transcriptional profiling of *Listeria monocytogenes* and mutagenesis identify new virulence factors involved in infection. *PLoS Pathog* 5: e1000449.
- Campero, C. M., Odeón, A. C., Cipolla, A. L., Moore, D. P., Poso, M. A. & Odriozola, E. (2002) Demonstration of *Listeria monocytogenes* by immunohistochemistry in formalin-fixed brain tissues from natural cases of ovine and bovine encephalitis. *J Vet Med B Infect Dis Vet Public Health* 49: 379-383.
- Canchaya, C., Giubellini, V., Ventura, M., de los Reyes-Gavilán, C. G. & Margolles, A. (2010) Mosaic-like sequences containing transposon, phage, and plasmid elements among *Listeria monocytogenes* plasmids. *Appl Environ Microbiol* 76: 4851-4857.
- Cao, Q., Feng, F., Wang, H., Xu, X., Chen, H., Cai, X. & Wang, X. (2018) Haemophilus parasuis CpxRA two-component system confers bacterial tolerance to environmental stresses and

- macrolide resistance. *Microbiol Res* 206: 177-185.
- Carattoli, A., Aschbacher, R., March, A., Larcher, C., Livermore, D. M. & Woodford, N. (2010) Complete nucleotide sequence of the IncN plasmid pKOX105 encoding VIM-1, QnrS1 and SHV-12 proteins in Enterobacteriaceae from Bolzano, Italy compared with IncN plasmids encoding KPC enzymes in the USA. *J Antimicrob Chemother* 65: 2070-2075.
- Carrique-Mas, J. J., Hökeberg, I., Andersson, Y., Arneborn, M., Tham, W., Danielsson-Tham, M.-L., Osterman, B., Leffler, M., Steen, M., Eriksson, E., Hedin, G. & Giesecke, J. (2003) Febrile gastroenteritis after eating on-farm manufactured fresh cheese – an outbreak of listeriosis? *Epidemiol Infect* 130: 79-86.
- Cartwright, E. J., Jackson, K. A., Johnson, S. D., Graves, L. M., Silk, B. J. & Mahon, B. E. (2013) Listeriosis outbreaks and associated food vehicles, United States, 1998-2008. *Emerg Infect Dis* 19: 1-9.
- Casino, P., Rubio, V. & Marina, A. (2010) The mechanism of signal transduction by two-component systems. *Curr Opin Struct Biol* 20: 763-771.
- Castro, H., Jaakkonen, A., Hakkinen, M., Korkeala, H. & Lindström, M. (2018) Occurrence, persistence, and contamination routes of *Listeria monocytogenes* genotypes on three Finnish dairy cattle farms: a longitudinal study. *Appl Environ Microbiol* 84: e02000-17.
- Castro, H., Ruusunen, M. & Lindström, M. (2017) Occurrence and growth of *Listeria monocytogenes* in packaged raw milk. *Int J Food Microbiol* 261: 1-10.
- CDC (Centers for Disease Control and Prevention). (2013) Vital signs: *Listeria* illnesses, deaths, and outbreaks – United States, 2009-2011. *MMWR Morb Mortal Wkly Rep* 62: 448-452.
- Chan, Y. C., Boor, K. J. & Wiedmann, M. (2007a) σ^B -dependent and σ^B -independent mechanisms contribute to transcription of *Listeria monocytogenes* cold stress genes during cold shock and cold growth. *Appl Environ Microbiol* 73: 6019-6029.
- Chan, Y. C., Hu, Y., Chaturongakul, S., Files, K. D., Bowen, B. M., Boor, K. J. & Wiedmann, M. (2008) Contributions of two-component regulatory systems, alternative sigma factors, and negative regulators to *Listeria monocytogenes* cold adaptation and cold growth. *J Food Prot* 71: 420-425.
- Chan, Y. C., Raengpradub, S., Boor, K. J. & Wiedmann, M. (2007b) Microarray-based characterization of the *Listeria monocytogenes* cold regulon in log- and stationary-phase cells. *Appl Environ Microbiol* 73: 6484-6498.
- Chan, Y. C. & Wiedmann, M. (2009) Physiology and genetics of *Listeria monocytogenes* survival and growth at cold temperatures. *Crit Rev Food Sci Nutr* 49: 237-253.
- Chand, P. & Sadana, J. R. (1999) Outbreak of *Listeria ivanovii* abortion in sheep in India. *Vet Rec* 145: 83-84.

- Chang, C. & Stewart, R. C. (1998) The two-component system. Regulation of diverse signaling pathways in prokaryotes and eukaryotes. *Plant Physiol* 117: 723-731.
- Chaturongakul, S., Raengpradub, S., Palmer, M. E., Bergholz, T. M., Orsi, R. H., Hu, Y., Ollinger, J., Wiedmann, M. & Boor, K. J. (2011) Transcriptomic and phenotypic analyses identify coregulated, overlapping regulons among PrfA, CtsR, HrcA, and the alternative sigma factors σ^B , σ^C , σ^H , and σ^L in *Listeria monocytogenes*. *Appl Environ Microbiol* 77: 187-200.
- Chaturongakul, S., Raengpradub, S., Wiedmann, M. & Boor, K. J. (2008) Modulation of stress and virulence in *Listeria monocytogenes*. *Trends Microbiol* 16: 388-396.
- Chen, L., Keramati, L. & Helmann, J. D. (1995) Coordinate regulation of *Bacillus subtilis* peroxide stress genes by hydrogen peroxide and metal ions. *Proc Natl Acad Sci U S A* 92: 8190-8194.
- Chen, Y., Luo, Y., Curry, P., Timme, R., Melka, D., Doyle, M., Parish, M., Hammack, T. S., Allard, M. W., Brown, E. W. & Strain, E. A. (2017) Assessing the genome level diversity of *Listeria monocytogenes* from contaminated ice cream and environmental samples linked to a listeriosis outbreak in the United States. *PLoS One* 12: e0171389.
- Cheroutre-Vialette, M., Lebert, I., Hebraud, M., Labadie, J. C. & Lebert, A. (1998) Effects of pH or a_w stress on growth of *Listeria monocytogenes*. *Int J Food Microbiol* 42: 71-77.
- Christiansen, J. K., Larsen, M. H., Ingmer, H., Sogaard-Andersen, L. & Kallipolitis, B. H. (2004) The RNA-binding protein Hfq of *Listeria monocytogenes*: role in stress tolerance and virulence. *J Bacteriol* 186: 3355-3362.
- Cole, M. B., Jones, M. V. & Holyoak, C. (1990) The effect of pH, salt concentration and temperature on the survival and growth of *Listeria monocytogenes*. *J Appl Bacteriol* 69: 63-72.
- Collins, B., Guinane, C. M., Cotter, P. D., Hill, C. & Ross, R. P. (2012) Assessing the contributions of the LiaS histidine kinase to the innate resistance of *Listeria monocytogenes* to nisin, cephalosporins, and disinfectants. *Appl Environ Microbiol* 78: 2923-2929.
- Cordero, N., Maza, F., Navea-Perez, H., Aravena, A., Marquez-Fontt, B., Navarrete, P., Figueroa, G., González, M., Latorre, M. & Reyes-Jara, A. (2016) Different transcriptional responses from slow and fast growth rate strains of *Listeria monocytogenes* adapted to low temperature. *Front Microbiol* 7: 229.
- Cossart, P. (2011) Illuminating the landscape of host-pathogen interactions with the bacterium *Listeria monocytogenes*. *Proc Natl Acad Sci U S A* 108: 19484-19491.
- Cotter, P. D., Draper, L. A., Lawton, E. M., Daly, K. M., Groeger, D. S., Casey, P. G., Ross, R. P. & Hill, C. (2008) Listeriolysin S, a novel peptide haemolysin associated with a subset of lineage I *Listeria*

- monocytogenes*. *PLoS Pathog* 4: e1000144.
- Cotter, P. D., Emerson, N., Gahan, C. G. M. & Hill, C. (1999) Identification and disruption of *lisRK*, a genetic locus encoding a two-component signal transduction system involved in stress tolerance and virulence in *Listeria monocytogenes*. *J Bacteriol* 181: 6840-6843.
- Cotter, P. D., Gahan, C. G. M. & Hill, C. (2000) Analysis of the role of the *Listeria monocytogenes* F₀F₁-ATPase operon in the acid tolerance response. *Int J Food Microbiol* 60: 137-146.
- Cotter, P. D., Gahan, C. G. M. & Hill, C. (2001a) A glutamate decarboxylase system protects *Listeria monocytogenes* in gastric fluid. *Mol Microbiol* 40: 465-475.
- Cotter, P. D., Guinane, C. M. & Hill, C. (2002) The LisRK signal transduction system determines the sensitivity of *Listeria monocytogenes* to nisin and cephalosporins. *Antimicrob Chemother* 46: 2784-2790.
- Cotter, P. D. & Hill, C. (2003) Surviving the acid test: responses of gram-positive bacteria to low pH. *Microbiol Mol Biol Rev* 67: 429-453.
- Cotter, P. D., O'Reilly, K. & Hill, C. (2001b) Role of the glutamate decarboxylase acid resistance system in the survival of *Listeria monocytogenes* LO28 in low pH foods. *J Food Prot* 64: 1362-1368.
- Crim, S. M., Iwamoto, M., Huang, J. Y., Griffin, P. M., Gilliss, D., Cronquist, A. B., Cartter, M., Tobin-D'Angelo, M., Blythe, D., Smith, K., Lathrop, S., Zansky, S., Cieslak, P. R., Dunn, J., Holt, K. G., Lance, S., Tauxe, R., Henao, O. L. & CDC. (2014) Incidence and trends of infection with pathogens transmitted commonly through food – Foodborne Diseases Active Surveillance Network, 10 U.S. sites, 2006-2013. *MMWR Morb Mortal Wkly Rep* 63: 328-332.
- Csonka, L. N. (1989) Physiological and genetic responses of bacteria to osmotic stress. *Microbiol Rev* 53: 121-147.
- Cummins, A. J., Fielding, A. K. & McLauchlin, J. (1994) *Listeria ivanovii* infection in a patient with AIDS. *J Infect* 28: 89-91.
- Dahlsten, E., Zhang, Z., Somervuo, P., Minton, N. P., Lindström, M. & Korkeala, H. (2014) The cold-induced two-component system CBO0366/CBO0365 regulates metabolic pathways with novel roles in group I *Clostridium botulinum* ATCC 3502 cold tolerance. *Appl Environ Microbiol* 80: 306-319.
- Dalton, C. B., Austin, C. C., Sobel, J., Hayes, P. S., Bibb, W. F., Graves, L. M., Swaminathan, B., Proctor, M. E. & Griffin, P. M. (1997) An outbreak of gastroenteritis and fever due to *Listeria monocytogenes* in milk. *N Engl J Med* 336: 100-105.
- de las Heras, A., Cain, R. J., Bielecka, M. K. & Vázquez-Boland, J. A. (2011) Regulation of *Listeria* virulence: PrfA master and commander. *Curr Opin Microbiol* 14: 118-127.
- de Valk, H., Jacquet, C., Goulet, V., Vaillant, V., Perra, A., Simon, F., Desenclos, J. C. & Martin, P. (2005) Surveillance of listeria infections in Europe. *Euro Surveill* 10: 251-255.

- del Solar, G., Giraldo, R., Ruiz-Echevarría, M. J., Espinosa, M. & Díaz-Orejas, R. (1998) Replication and control of circular bacterial plasmids. *Microbiol Mol Biol Rev* 62: 434-464.
- den Bakker, H. C., Bowen, B. M., Rodriguez-Rivera, L. D. & Wiedmann, M. (2012) FSL J1-208, a virulent uncommon phylogenetic lineage IV *Listeria monocytogenes* strain with a small chromosome size and a putative virulence plasmid carrying internalin-like genes. *Appl Environ Microbiol* 78: 1876-1889.
- den Bakker, H. C., Manuel, C. S., Fortes, E. D., Wiedmann, M. & Nightingale, K. K. (2013) Genome sequencing identifies *Listeria fleischmannii* subsp. *coloradonensis* subsp. nov., isolated from a ranch. *Int J Syst Evol Microbiol* 63: 3257-3268.
- den Bakker, H. C., Warchocki, S., Wright, E. M., Allred, A. F., Ahlstrom, C., Manuel, C. S., Stasiewicz, M. J., Burrell, A., Roof, S., Strawn, L. K., Fortes, E., Nightingale, K. K., Kephart, D. & Wiedmann, M. (2014) *Listeria floridensis* sp. nov., *Listeria aquatica* sp. nov., *Listeria cornellensis* sp. nov., *Listeria riparia* sp. nov. and *Listeria grandensis* sp. nov., from agricultural and natural environments. *Int J Syst Evol Microbiol* 64: 1882-1889.
- den Besten, H. M. W., Aryani, D. C., Metselaar, K. I. & Zwietering, M. H. (2017) Microbial variability in growth and heat resistance of a pathogen and a spoiler: All variabilities are equal but some are more equal than others. *Int J Food Microbiol* 240: 24-31.
- Denyer, S. P. (1995) Mechanisms of action of antibacterial biocides. *Int Biodeterior Biodegradation* 36: 227-245.
- Derman, Y., Isokallio, M., Lindström, M. & Korkeala, H. (2013) The two-component system CBO2306/CBO2307 is important for cold adaptation of *Clostridium botulinum* ATCC 3502. *Int J Food Microbiol* 167: 87-91.
- Derré, I., Rapoport, G. & Msadek, T. (1999) CtsR, a novel regulator of stress and heat shock response, controls *clp* and molecular chaperone gene expression in Gram-positive bacteria. *Mol Microbiol* 31: 117-131.
- Dintner, S., Staron, A., Berchtold, E., Petri, T., Mascher, T. & Gebhard, S. (2011) Coevolution of ABC transporters and two-component regulatory systems as resistance modules against antimicrobial peptides in *Firmicutes* Bacteria. *J Bacteriol* 193: 3851-3862.
- Disson, O. & Lecuit, M. (2012) Targeting of the central nervous system by *Listeria monocytogenes*. *Virulence* 3: 213-221.
- Dombek, K. M. & Ingram, L. O. (1984) Effects of ethanol on the *Escherichia coli* plasma membrane. *J Bacteriol* 157: 233-239.
- Dons, L., Eriksson, E., Jin, Y., Rottenberg, M. E., Kristensson, K., Larsen, C. N., Bresciani, J. & Olsen, J. E. (2004) Role of flagellin and the two-component CheA/CheY system of *Listeria monocytogenes* in host cell invasion and virulence. *Infect Immun* 72: 3237-3244.

- Doumith, M., Buchrieser, C., Glaser, P., Jacquet, C. & Martin, P. (2004) Differentiation of the major *Listeria monocytogenes* serovars by multiplex PCR. *J Clin Microbiol* 42: 3819-3822.
- Doyle, M. E., Mazzotta, A. S., Wang, T., Wiseman, D. W. & Scott, V. N. (2001) Heat resistance of *Listeria monocytogenes*. *J Food Prot* 64: 410-429.
- Doyle, M. P., Glass, K. A., Beery, J. T., Garcia, G. A., Pollard, D. J. & Schultz, R. D. (1987) Survival of *Listeria monocytogenes* in milk during high-temperature, short-time pasteurization. *Appl Environ Microbiol* 53: 1433-1438.
- Dubrac, S., Bisicchia, P., Devine, K. M. & Msadek, T. (2008) A matter of life and death: cell wall homeostasis and the WalKR (YycGF) essential signal transduction pathway. *Mol Microbiol* 70: 1307-1322.
- Duché, O., Trémoulet, F., Glaser, P. & Labadie, J. (2002a) Salt stress proteins induced in *Listeria monocytogenes*. *Appl Environ Microbiol* 68: 1491-1498.
- Duché, O., Trémoulet, F., Namane, A., the European *Listeria* Genome Consortium & Labadie, J. (2002b) A proteomic analysis of the salt stress response of *Listeria monocytogenes*. *FEMS Microbiol Lett* 215: 183-188.
- Dumont, J. & Cotoni, L. (1921) Bacille semblable à celui du rouget du porc rencontré dans le L.C.R. d'un méningitique. *Ann Inst Pasteur* 35: 625-633.
- Dussurget, O., Dumas, E., Archambaud, C., Chafsey, I., Chambon, C., Hébraud, M. & Cossart, P. (2005) *Listeria monocytogenes* ferritin protects against multiple stresses and is required for virulence. *FEMS Microbiol Lett* 250: 253-261.
- Dutta, V., Elhanafi, D. & Kathariou, S. (2013) Conservation and distribution of the benzalkonium chloride resistance cassette *bcrABC* in *Listeria monocytogenes*. *Appl Environ Microbiol* 79: 6067-6074.
- Dykes, G. A., Geornaras, I., Papathanasopoulos, M. A. & von Holy, A. (1994) Plasmid profiles of *Listeria* species associated with poultry processing. *Food Microbiol* 11: 519-523.
- Dykes, G. A. & Moorhead, S. M. (2000) Survival of osmotic and acid stress by *Listeria monocytogenes* strains of clinical or meat origin. *Int J Food Microbiol* 56: 161-166.
- Ebe, Y., Hasegawa, G., Takatsuka, H., Umezu, H., Mitsuyama, M., Arakawa, M., Mukaida, N. & Naito, M. (1999) The role of Kupffer cells and regulation of neutrophil migration into the liver by macrophage inflammatory protein-2 in primary listeriosis in mice. *Pathol Int* 49: 519-532.
- ECDC (European Centre for Disease Prevention and Control). (2016) Annual epidemiological report 2016 – listeriosis. <https://ecdc.europa.eu/en/publications-data/listeriosis-annual-epidemiological-report-2016-2014-data> Accessed at 27 Mar 2019.
- Elhanafi, D., Dutta, V. & Kathariou, S. (2010) Genetic characterization of plasmid-associated benzalkonium chloride resistance determinants in a

- Listeria monocytogenes* strain from the 1998-1999 outbreak. *Appl Environ Microbiol* 76: 8231-8238.
- Ericsson, H., Danielsson-Tham, M.-L., Stålhandske, P., Tham, W. & Ursing, J. (1993) Subtyping of a frequent phagovar of *Listeria monocytogenes* in Sweden by use of restriction endonuclease analysis. *APMIS* 101: 971-974.
- Ericsson, H., Eklöv, A., Danielsson-Tham, M.-L., Loncarevic, S., Mentzing, L.-O., Persson, I., Unnerstad, H. & Tham, W. (1997) An outbreak of listeriosis suspected to have been caused by rainbow trout. *J Clin Microbiol* 35: 2904-2907.
- Evans, J. R., Allen, A. C., Stinson, D. A., Bortolussi, R. & Peddle, L. J. (1985) Perinatal listeriosis: report of an outbreak. *Pediatr Infect Dis* 4: 237-241.
- Fabret, C. & Hoch, J. A. (1998) A two-component signal transduction system essential for growth of *Bacillus subtilis*: implications for anti-infective therapy. *J Bacteriol* 180: 6375-6383.
- Faleiro, M. L., Andrew, P. W. & Power, D. (2003) Stress response of *Listeria monocytogenes* isolated from cheese and other foods. *Int J Food Microbiol* 84: 207-216.
- Farber, J. M., Daley, E., Coates, F., Beausoleil, N. & Fournier, J. (1991) Feeding trials of *Listeria monocytogenes* with a nonhuman primate model. *J Clin Microbiol* 29: 2606-2608.
- Farber, J. M. & Pagotto, F. (1992) The effect of acid shock on the heat resistance of *Listeria monocytogenes*. *Lett Appl Microbiol* 15: 197-201.
- Fernández, L., Gooderham, W. J., Bains, M., McPhee, J. B., Wiegand, I. & Hancock, R. E. W. (2010) Adaptive resistance to the "last hope" antibiotics polymyxin B and colistin in *Pseudomonas aeruginosa* is mediated by the novel two-component regulatory system ParR-ParS. *Antimicrob Agents Chemother* 54: 3372-3382.
- Ferreira, A., O'Byrne, C. P. & Boor, K. J. (2001) Role of σ^B in heat, ethanol, acid, and oxidative stress resistance and during carbon starvation in *Listeria monocytogenes*. *Appl Environ Microbiol* 67: 4454-4457.
- Fistrovici, E. & Collins-Thompson, D. L. (1990) Use of plasmid profiles and restriction endonuclease digest in environmental studies of *Listeria* spp. from raw milk. *Int J Food Microbiol* 10: 43-50.
- Flamm, R. K., Hinrichs, D. J. & Thomashow, M. F. (1984) Introduction of pAM β 1 into *Listeria monocytogenes* by conjugation and homology between native *L. monocytogenes* plasmids. *Infect Immun* 44: 157-161.
- Flanary, P. L., Allen, R. D., Dons, L. & Kathariou, S. (1999) Insertional inactivation of the *Listeria monocytogenes* *cheYA* operon abolishes response to oxygen gradients and reduces the number of flagella. *Can J Microbiol* 45: 646-652.
- Fleming, D. W., Cochi, S. L., MacDonald, K. L., Brondum, J., Hayes, P. S., Plikaytis, B. D., Holmes, M. B., Audurier, A., Broome, C. V. & Reingold, A. L. (1985) Pasteurized milk as a vehicle of infection in an

- outbreak of listeriosis. *N Engl J Med* 312: 404-407.
- Fox, E. M., Allnutt, T., Bradbury, M. I., Fanning, S. & Chandry, P. S. (2016) Comparative genomics of the *Listeria monocytogenes* ST204 subgroup. *Front Microbiol* 7: 2057.
- Fox, E. M., Leonard, N. & Jordan, K. (2011) Physiological and transcriptional characterization of persistent and nonpersistent *Listeria monocytogenes* isolates. *Appl Environ Microbiol* 77: 6559-6569.
- Francois, K., Devlieghere, F., Standaert, A. R., Geeraerd, A. H., Van Impe, J. F. & Debevere, J. (2006) Effect of environmental parameters (temperature, pH and a_w) on the individual cell lag phase and generation time of *Listeria monocytogenes*. *Int J Food Microbiol* 108: 326-335.
- Fraser, K. R., Sue, D., Wiedmann, M., Boor, K. & O'Byrne, C. P. (2003) Role of σ^B in regulating the compatible solute uptake systems of *Listeria monocytogenes*: osmotic induction of *opuC* is σ^B dependent. *Appl Environ Microbiol* 69: 2015-2022.
- Fritsch, F., Mauder, N., Williams, T., Weiser, J., Oberle, M. & Beier, D. (2011) The cell envelope stress response mediated by the LiaFSR_{Lm} three-component system of *Listeria monocytogenes* is controlled via the phosphatase activity of the bifunctional histidine kinase LiaS_{Lm}. *Microbiology* 157: 373-386.
- Frost, L. S., Lepiae, R., Summers, A. O. & Toussaint, A. (2005) Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol* 3: 722-732.
- Frye, D. M., Zweig, R., Sturgeon, J., Tormey, M., LeCavalier, M., Lee, I., Lawani, L. & Mascola, L. (2002) An outbreak of febrile gastroenteritis associated with delicatessen meat contaminated with *Listeria monocytogenes*. *Clin Infect Dis* 35: 943-949.
- Fuangthong, M., Herbig, A. F., Bsat, N. & Helmann, J. D. (2002) Regulation of the *Bacillus subtilis* *fur* and *perR* genes by PerR: not all members of the PerR regulon are peroxide inducible. *J Bacteriol* 184: 3276-3286.
- Galinski, E. A. (1995) Osmoadaptation in bacteria. *Adv Microb Physiol* 37: 272-328.
- Galsworthy, S. B., Girdler, S. & Koval, S. F. (1990) Chemotaxis in *Listeria monocytogenes*. *Acta Microbiol Hung* 37: 81-85.
- Gandhi, M. & Chikindas, M. L. (2007) *Listeria*: a foodborne pathogen that knows how to survive. *Int J Food Microbiol* 113: 1-15.
- Gao, R. & Stock, A. M. (2009) Biological insights from structures of two-component proteins. *Annu Rev Microbiol* 63: 133-54.
- Gardan, R., Duché, O., Leroy-Sétrin, S., the European *Listeria* Genome Consortium & Labadie, J. (2003) Role of *ctc* from *Listeria monocytogenes* in osmotolerance. *Appl Environ Microbiol* 69: 154-161.
- Garmyn, D., Augagneur, Y., Gal, L., Vivant, A.-L. & Piveteau, P. (2012) *Listeria monocytogenes* differential transcriptome analysis reveals temperature-dependent Agr regulation

- and suggests overlaps with other regulons. *PLoS One* 7: e43154.
- George, S. M., Lund, B. M. & Brocklehurst, T. F. (1988) The effect of pH and temperature on initiation of growth of *Listeria monocytogenes*. *Lett Appl Microbiol* 6: 153-156.
- Gianfranceschi, M., Gattuso, A., Tartaro, S. & Aureli, P. (2003) Incidence of *Listeria monocytogenes* in food and environmental samples in Italy between 1990 and 1999: serotype distribution in food, environmental and clinical samples. *Eur J Epidemiol* 18: 1001-1006.
- Gilmour, M. W., Graham, M., Van Domselaar, G., Tyler, S., Kent, H., Trout-Yakel, K. M., Larios, O., Allen, V., Lee, B. & Nadon, C. (2010) High-throughput genome sequencing of two *Listeria monocytogenes* clinical isolates during a large foodborne outbreak. *BMC Genomics* 11: 120.
- Gilot, P., Genicot, A. & André, P. (1996) Serotyping and esterase typing for analysis of *Listeria monocytogenes* populations recovered from foodstuffs and from human patients with listeriosis in Belgium. *J Clin Microbiol* 34: 1007-1010.
- Giotis, E. S., Blair, I. S. & McDowell, D. A. (2007a) Morphological changes in *Listeria monocytogenes* subjected to sublethal alkaline stress. *Int J Food Microbiol* 120: 250-258.
- Giotis, E. S., Julotok, M., Wilkinson, B. J., Blair, I. S. & McDowell, D. A. (2008a) Role of sigma B factor in the alkaline tolerance response of *Listeria monocytogenes* 10403S and cross-protection against subsequent ethanol and osmotic stress. *J Food Prot* 71: 1481-1485.
- Giotis, E. S., McDowell, D. A., Blair, I. S. & Wilkinson, B. J. (2007b) Role of branched-chain fatty acids in pH stress tolerance in *Listeria monocytogenes*. *Appl Environ Microbiol* 73: 997-1001.
- Giotis, E. S., Muthaiyan, A., Blair, I. S., Wilkinson, B. J. & McDowell, D. A. (2008b) Genomic and proteomic analysis of the Alkali-Tolerance Response (AlTR) in *Listeria monocytogenes* 10403S. *BMC Microbiol* 8: 102.
- Giotis, E. S., Muthaiyan, A., Natesan, S., Wilkinson, B. J., Blair, I. S. & McDowell, D. A. (2010) Transcriptome analysis of alkali shock and alkali adaptation in *Listeria monocytogenes* 10403S. *Foodborne Pathog Dis* 7: 1147-1157.
- Glaser, P., Frangeul, L., Buchrieser, C., Rusniok, C., Amend, A., Baquero, F., Berche, P., Bloeker, H., Brandt, P., Chakraborty, T., Charbit, A., Chetouani, F., Couvé, E., de Daruvar, A., Dehoux, P., Domann, E., Domínguez-Bernal, G., Duchaud, E., Durant, L., Dussurget, O., Entian, K.-D., Fsihi, H., Garcia-Del Portillo, F., Garrido, P., Gautier, L., Goebel, W., Gómez-López, N., Hain, T., Hauf, J., Jackson, D., Jones, L.-M., Kaerst, U., Kreft, J., Kuhn, M., Kunst, F., Kurapkat, G., Madueño, E., Maitournam, A., Vicente, J. M., Ng, E., Nedjari, H., Nordsiek, G., Novella, S., de Pablos, B., Pérez-Díaz, J.-C., Purcell, R., Rammel, B., Rose, M., Schlueter, T., Simoes, N., Tierrez, A., Vázquez-Boland, J.-A., Voss, H., Wehland, J. &

- Cossart, P. (2001) Comparative genomics of *Listeria* species. *Science* 294: 849-852.
- Godshall, C. E., Suh, G. & Lorber, B. (2013) Cutaneous listeriosis. *J Clin Microbiol* 51: 3591-3596.
- Goessweiner-Mohr, N., Arends, K., Keller, W. & Grohmann, E. (2014) Conjugation in Gram-positive bacteria. *Microbiol Spectr* 2: PLAS-0004-2013.
- Gold, R. S., Meagher, M. M., Hutkins, R. & Conway, T. (1992) Ethanol tolerance and carbohydrate metabolism in lactobacilli. *J Ind Microbiol* 10: 45-54.
- Gorski, L., Flaherty, D. & Duhé, J. M. (2008) Comparison of the stress response of *Listeria monocytogenes* strains with sprout colonization. *J Food Prot* 71: 1556-1562.
- Gottschalk, S., Bygebjerg-Hove, I., Bonde, M., Nielsen, P. K., Nguyen, T. H., Gravesen, A. & Kallipolitis, B. H. (2008) The two-component system CesRK controls the transcriptional induction of cell envelope-related genes in *Listeria monocytogenes* in response to cell wall-acting antibiotics. *J Bacteriol* 190: 4772-4776.
- Gouin, E., Mengaud, J. & Cossart, P. (1994) The virulence gene cluster of *Listeria monocytogenes* is also present in *Listeria ivanovii*, an animal pathogen, and *Listeria seeligeri*, a nonpathogenic species. *Infect Immun* 62: 3550-3553.
- Goulet, V., Hebert, M., Hedberg, C., Laurent, E., Vaillant, V., de Valk, H. & Desenclos, J. C. (2012) Incidence of listeriosis and related mortality among groups at risk of acquiring listeriosis. *Clin Infect Dis* 54: 652-660.
- Goulet, V., King, L. A., Vaillant, V. & de Valk, H. (2013) What is the incubation period for listeriosis? *BMC Infect Dis* 13: 11.
- Goulet, V. & Marchetti, P. (1996) Listeriosis in 225 non-pregnant patients in 1992: clinical aspects and outcome in relation to predisposing conditions. *Scand J Infect Dis* 28: 367-374.
- Gouliau, M. (2010) Two-component signaling circuit structure and properties. *Curr Opin Microbiol* 13: 184-189.
- Grad, Y. H. & Fortune, S. M. (2016) Biodiversity and hypervirulence of *Listeria monocytogenes*. *Nat Genet* 48: 229-230.
- Graves, L. M., Helsel, L. O., Steigerwalt, A. G., Morey, R. E., Daneshvar, M. I., Roof, S. E., Orsi, R. H., Fortes, E. D., Milillo, S. R., den Bakker, H. C., Wiedmann, M., Swaminathan, B. & Sauders, B. D. (2010) *Listeria marthii* sp. nov., isolated from the natural environment, Finger Lakes National Forest. *Int J Syst Evol Microbiol* 60: 1280-1288.
- Graves, L. M., Swaminathan, B., Reeves, M. W., Hunter, S. B., Weaver, R. E., Plikaytis, B. D. & Schuchat, A. (1994) Comparison of ribotyping and multilocus enzyme electrophoresis for subtyping of *Listeria monocytogenes* isolates. *J Clin Microbiol* 32: 2936-2943.
- Graves, L. M., Swaminathan, B., Reeves, M. W. & Wenger, J. (1991) Ribosomal DNA fingerprinting of *Listeria monocytogenes* using a digoxigenin-labeled DNA

- probe. *Eur J Epidemiol* 7: 77-82.
- Gray, M. L. (1963) Epidermiological aspects of listeriosis. *Am J Public Health Nations Health* 53: 554-563.
- Gray, M. L. & Killinger, A. H. (1966) *Listeria monocytogenes* and listeric infections. *Bacteriol Rev* 30: 309-382.
- Gregory, S. H. & Liu, C.-C. (2000) CD8⁺ T-cell-mediated response to *Listeria monocytogenes* taken up in the liver and replicating within hepatocytes. *Immunol Rev* 174: 112-122.
- Grubaugh, D., Regeimbal, J. M., Ghosh, P., Zhou, Y., Lauer, P., Dubensky, T. W., Jr., & Higgins, D. E. (2018) The VirAB ABC transporter is required for VirR regulation of *Listeria monocytogenes* virulence and resistance to nisin. *Infect Immun* 86: e00901-17.
- Guckes, K. R., Breland, E. J., Zhang, E. W., Hanks, S. C., Gill, N. K., Algood, H. M., Schmitz, J. E., Stratton, C. W. & Hadjifrangiskou, M. (2017) Signaling by two-component system noncognate partners promotes intrinsic tolerance to polymyxin B in uropathogenic *Escherichia coli*. *Sci Signal* 10: eaag1775.
- Gueriri, I., Cyncynatus, C., Dubrac, S., Toledo-Arana, A., Dussurget, O. & Msadek, T. (2008) The DegU orphan response regulator of *Listeria monocytogenes* autorepresses its own synthesis and is required for bacterial motility, virulence and biofilm formation. *Microbiology* 154: 2251-2264.
- Guglielmetti, E., Korhonen, J. M., Heikkinen, J., Morelli, L. & von Wright, A. (2009) Transfer of plasmid-mediated resistance to tetracycline in pathogenic bacteria from fish and aquaculture environments. *FEMS Microbiol Lett* 293: 28-34.
- Guillet, C., Join-Lambert, O., Le Monnier, A., Leclercq, A., Mechai, F., Mamzer-Bruneel, M.-F., Bielecka, M. K., Scortti, M., Disson, O., Berche, P., Vazquez-Boland, J., Lortholary, O. & Lecuit, M. (2010) Human listeriosis caused by *Listeria ivanovii*. *Emerg Infect Dis* 16: 136-138.
- Gullberg, E., Albrecht, L. M., Karlsson, C., Sandegren, L. & Andersson, D. I. (2014) Selection of a multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. *MBio* 5: e01918-14.
- Gutierrez, C., Abee, T. & Booth, I. R. (1995) Physiology of the osmotic stress response in microorganisms. *Int J Food Microbiol* 28: 233-244.
- Haase, J. K., Didelot, X., Lecuit, M., Korkeala, H., *L. monocytogenes* MLST Study Group & Achtman, M. (2014) The ubiquitous nature of *Listeria monocytogenes* clones: a large-scale Multilocus Sequence Typing study. *Environ Microbiol* 16: 405-416.
- Hadorn, K., Hächler, H., Schaffner, A. & Kayser, F. H. (1993) Genetic characterization of plasmid-encoded multiple antibiotic resistance in a strain of *Listeria monocytogenes* causing endocarditis. *Eur J Clin Microbiol Infect Dis* 12: 928-937.
- Haft, R. J. F., Keating, D. H., Schwaegler, T., Schwalbach, M. S., Vinokur, J., Tremaine, M., Peters, J. M.,

- Kotlajich, M. V., Pohlmann, E. L., Ong, I. M., Grass, J. A., Kiley, P. J. & Landick, R. (2014) Correcting direct effects of ethanol on translation and transcription machinery confers ethanol tolerance in bacteria. *Proc Natl Acad Sci USA* 111: E2576-E2585.
- Hanawa, T., Kai, M., Kamiya, S. & Yamamoto, T. (2000) Cloning, sequencing, and transcriptional analysis of the *dnaK* heat shock operon of *Listeria monocytogenes*. *Cell Stress Chaperones* 5: 21-29.
- Harter, E., Wagner, E. M., Zaiser, A., Halecker, S., Wagner, M. & Rychli, K. (2017) Stress survival islet 2, predominantly present in *Listeria monocytogenes* strains of sequence type 121, is involved in the alkaline and oxidative stress responses. *Appl Environ Microbiol* 83: e00827-17.
- Hayakawa, T., Tanaka, T., Sakaguchi, K., Ōtake, N. & Yonehara, H. (1979) A linear plasmid-like DNA in *Streptomyces* sp. producing lankacidin group antibiotics. *J Gen Appl Microbiol* 25: 255-260.
- Hébraud, M. & Guzzo, J. (2000) The main cold shock protein of *Listeria monocytogenes* belongs to the family of ferritin-like proteins. *FEMS Microbiol Lett* 190: 29-34.
- Hébraud, M. & Potier, P. (1999) Cold shock response and low temperature adaptation in psychrotrophic bacteria. *J Mol Microbiol Biotechnol* 1: 211-219.
- Helmann, J. D., Wu, M. F. W., Gaballa, A., Kobel, P. A., Morshedi, M. M., Fawcett, P. & Paddon, C. (2003) The global transcriptional response of *Bacillus subtilis* to peroxide stress is coordinated by three transcription factors. *J Bacteriol* 185: 243-253.
- Herbig, A. F. & Helmann, J. D. (2001) Roles of metal ions and hydrogen peroxide in modulating the interaction of the *Bacillus subtilis* PerR peroxide regulon repressor with operator DNA. *Mol Microbiol* 41: 849-859.
- Herrero, A., Mendoza, M. C., Rodicio, R. & Rodicio, M. R. (2008) Characterization of pUO-StVR2, a virulence-resistance plasmid evolved from the pSLT virulence plasmid of *Salmonella enterica* serovar Typhimurium. *Antimicrob Agents Chemother* 52: 4514-4517.
- Hill, C., O'Driscoll, B. & Booth, I. (1995) Acid adaptation and food poisoning microorganisms. *Int J Food Microbiol* 28: 245-254.
- Hinderink, K., Lindström, M. & Korkeala, H. (2009) Group I *Clostridium botulinum* strains show significant variation in growth at low and high temperatures. *J Food Prot* 72: 375-383.
- Hingston, P., Chen, J., Allen, K., Truelstrup Hansen, L. & Wang, S. (2017a) Strand specific RNA-sequencing and membrane lipid profiling reveals growth phase-dependent cold stress response mechanisms in *Listeria monocytogenes*. *PLoS One* 12: e0180123.
- Hingston, P., Chen, J., Dhillon, B. K., Laing, C., Bertelli, C., Gannon, V., Tasara, T., Allen, K., Brinkman, F. S. L., Truelstrup Hansen, L. & Wang, S. (2017b) Genotypes associated with *Listeria monocytogenes* isolates displaying impaired or enhanced tolerances to cold, salt, acid, or desiccation

- stress. *Front Microbiol* 8: 369.
- Hingston, P. A., Piercey, M. J. & Truelstrup Hansen, L. (2015) Genes associated with desiccation and osmotic stress in *Listeria monocytogenes* as revealed by insertional mutagenesis. *Appl Environ Microbiol* 81: 5350-5362.
- Hinnebusch, J. & Tilly, K. (1993) Linear plasmids and chromosomes in bacteria. *Mol Microbiol* 10: 917-922.
- Hoffman, A. D., Gall, K. L., Norton, D. M. & Wiedmann, M. (2003) *Listeria monocytogenes* contamination patterns for the smoked fish processing environment and for raw fish. *J Food Prot* 66: 52-60.
- Houari, A. & Di Martino, P. (2007) Effect of chlorhexidine and benzalkonium chloride on bacterial biofilm formation. *Lett Appl Microbiol* 45: 652-656.
- Hu, Y., Oliver, H. F., Raengpradub, S., Palmer, M. E., Orsi, R. H., Wiedmann, M. & Boor, K. J. (2007a) Transcriptomic and phenotypic analyses suggest a network between the transcriptional regulators HrcA and σ^B in *Listeria monocytogenes*. *Appl Environ Microbiol* 73: 7981-7991.
- Hu, Y., Raengpradub, S., Schwab, U., Loss, C., Orsi, R. H., Wiedmann, M. & Boor, K. J. (2007b) Phenotypic and transcriptomic analyses demonstrate interactions between the transcriptional regulators CtsR and sigma B in *Listeria monocytogenes*. *Appl Environ Microbiol* 73: 7967-7980.
- Hudson, J. A., Mott, S. J. & Penney, N. (1994) Growth of *Listeria monocytogenes*, *Aeromonas hydrophila*, and *Yersinia enterocolitica* on vacuum and saturated carbon dioxide controlled atmosphere-packaged sliced roast beef. *J Food Prot* 57: 204-208.
- Hülphers, G. (1911) Lefvernekros hos kanin orsakad af en ej förut beskrifven bakterie. *Sven Vet Tidskr* 2: 265-273.
- Huffer, S., Clark, M. E., Ning, J. C., Blanch, H. W. & Clark, D. S. (2011) Role of alcohols in growth, lipid composition, and membrane fluidity of yeasts, bacteria, and archaea. *Appl Environ Microbiol* 77: 6400-6408.
- Husu, J. R. (1990) Epidemiological studies on the occurrence of *Listeria monocytogenes* in the feces of dairy cattle. *Zentralbl Veterinarmed B* 37: 276-282.
- Husu, J. R., Sivelä, S. K. & Rauramaa, A. L. (1990a) Prevalence of *Listeria* species as related to chemical quality of farm-ensiled grass. *Grass Forage Sci* 45: 309-314.
- Husu, J. R., Seppänen, J. T., Sivelä, S. K. & Rauramaa, A. L. (1990b) Contamination of raw milk by *Listeria monocytogenes* on dairy farms. *Zentralbl Veterinarmed B* 37: 268-275.
- Imlay, J. A. (2002) How oxygen damages microbes: oxygen tolerance and obligate anaerobiosis. *Adv Microb Physiol* 46: 111-153.
- Imlay, J. A. (2015a) Diagnosing oxidative stress in bacteria: not as easy as you might think. *Curr Opin Microbiol* 24: 124-131.
- Imlay, J. A. (2015b) Transcription factors that defend bacteria against reactive oxygen species. *Annu Rev Microbiol* 69: 93-108.
- Ingram, L. O. (1976) Adaptation of membrane lipids to

- alcohols. *J Bacteriol* 125: 670-678.
- Ingram, L. O. (1990) Ethanol tolerance in bacteria. *Crit Rev Biotechnol* 9: 305-319.
- Ivanov, I. (1962) Untersuchungen über die Listeriose der Schafe in Bulgarien. *Monatsh Veterinarmed* 17: 729-736.
- Jacks, A., Pihlajasaari, A., Vahe, M., Myntti, A., Kaukoranta, S.-S., Elomaa, N., Salmenlinna, S., Rantala, L., Lahti, K., Huusko, S., Kuusi, M., Siitonen, A. & Rimhanen-Finne, R. (2016) Outbreak of hospital-acquired gastroenteritis and invasive infection caused by *Listeria monocytogenes*, Finland, 2012. *Epidemiol Infect* 144: 2732-2742.
- Jacob-Dubuisson, F., Mechaly, A., Betton, J.-M. & Antoine, R. (2018) Structural insights into the signalling mechanisms of two-component systems. *Nat Rev Microbiol* 16: 585-593.
- Jensen, V. B., Harty, J. T. & Jones, B. D. (1998) Interactions of the invasive pathogens *Salmonella typhimurium*, *Listeria monocytogenes*, and *Shigella flexneri* with M cells and murine Peyer's patches. *Infect Immun* 66: 3758-3766.
- Jiang, X., Yu, T., Liang, Y., Ji, S., Guo, X., Ma, J. & Zhou, L. (2016) Efflux pump-mediated benzalkonium chloride resistance in *Listeria monocytogenes* isolated from retail food. *Int J Food Microbiol* 217: 141-145.
- Jiang, X., Yu, T., Xu, Y., Wang, H., Korkeala, H. & Shi, L. (2019) MdrL, a major facilitator superfamily efflux pump of *Listeria monocytogenes* involved in tolerance to benzalkonium chloride. *Appl Microbiol Biotechnol* 103: 1339-1350.
- Jones, G. S., Bussell, K. M., Myers-Morales, T., Fieldhouse, A. M., Bou Ghanem, E. N. & D'Orazio, S. E. F. (2015) Intracellular *Listeria monocytogenes* comprises a minimal but vital fraction of the intestinal burden following foodborne infection. *Infect Immun* 83: 3146-3156.
- Junttila, J. R., Niemelä, S. I. & Hirn, J. (1988) Minimum growth temperatures of *Listeria monocytogenes* and non-haemolytic listeria. *J Appl Bacteriol* 65: 321-327.
- Jydegaard-Axelsen, A.-M., Aaes-Jørgensen, A., Koch, A. G., Jensen, J. S. & Knøchel, S. (2005) Changes in growth, rRNA content, and cell morphology of *Listeria monocytogenes* induced by CO₂ up- and downshift. *Int J Food Microbiol* 98: 145-155.
- Jørgensen, S. B., Søråas, A., Arnesen, L. S., Leegaard, T., Sundsfjord, A. & Jenum, P. A. (2017) First environmental sample containing plasmid-mediated colistin-resistant ESBL-producing *Escherichia coli* detected in Norway. *APMIS* 125: 822-825.
- Kachroo, P., Eraso, J. M., Beres, S. B., Olsen, R. J., Zhu, L., Nasser, W., Bernard, P. E., Cantu, C. C., Saavedra, M. O., Arredondo, M. J., Strobe, B., Do, H., Kumaraswami, M., Vuopio, J., Gröndahl-Yli-Hannuksela, K., Kristinsson, K. G., Gottfredsson, M., Pesonen, M., Pensar, J., Davenport, E. R., Clark, A. G., Corander, J., Caugant, D. A., Gaini, S., Magnussen, M.

- D., Kubiak, S. L., Nguyen, H. A. T., Long, S. W., Porter, A. R., Deleo, F. R. & Musser, J. M. (2019) Integrated analysis of population genomics, transcriptomics and virulence provides novel insights into *Streptococcus pyogenes* pathogenesis. *Nat Genet* 51: 548-559.
- Kado, C. I. (1998) Origin and evolution of plasmids. *Antonie Van Leeuwenhoek* 73: 117-126.
- Kahm, M., Hasenbrink, G., Lichtenberg-Fraté, H., Ludwig, J. & Kschischo, M. (2010) grofit: fitting biological growth curves with R. *J Stat Softw* 33: 1-21.
- Kallipolitis, B. H. & Ingmer, H. (2001) *Listeria monocytogenes* response regulators important for stress tolerance and pathogenesis. *FEMS Microbiol Lett* 204: 111-115.
- Kallipolitis, B. H., Ingmer, H., Gahan, C. G., Hill, C. & Søgaard-Andersen, L. (2003) CesRK, a two-component signal transduction system in *Listeria monocytogenes*, responds to the presence of cell wall-acting antibiotics and affects β -lactam resistance. *Antimicrob Agents Chemother* 47: 3421-3429.
- Katharios-Lanwermyer, S., Rakic-Martinez, M., Elhanafi, D., Ratani, S., Tiedje, J. M. & Kathariou, S. (2012) Coselection of cadmium and benzalkonium chloride resistance in conjugative transfers from nonpathogenic *Listeria* spp. to other *Listeriae*. *Appl Environ Microbiol* 78: 7549-7556.
- Kathariou, S. (2002) *Listeria monocytogenes* virulence and pathogenicity, a food safety perspective. *J Food Prot* 65: 1811-1829.
- Katzav, M., Hyvönen, P., Muje, P., Rantala, L. & von Wright, A. (2006) Pulsed-field gel electrophoresis typing of *Listeria monocytogenes* isolated in two Finnish fish farms. *J Food Prot* 69: 1443-1447.
- Kazmierczak, M. J., Mithoe, S. C., Boor, K. J. & Wiedmann, M. (2003) *Listeria monocytogenes* σ^B regulates stress response and virulence functions. *J Bacteriol* 185: 5722-5734.
- Keim, P., Price, L. B., Klevytska, A. M., Smith, K. L., Schupp, J. M., Okinaka, R., Jackson, P. J. & Hugh-Jones, M. E. (2000) Multiple-locus variable-number tandem repeat analysis reveals genetic relationships within *Bacillus anthracis*. *J Bacteriol* 182: 2928-2936.
- Kempf, B. & Bremer, E. (1998) Uptake and synthesis of compatible solutes as microbial stress responses to high-osmolality environments. *Arch Microbiol* 170: 319-330.
- Keto-Timonen, R. O., Autio, T. J. & Korkeala, H. J. (2003) An improved amplified fragment length polymorphism (AFLP) protocol for discrimination of *Listeria* isolates. *Syst Appl Microbiol* 26: 236-244.
- Keto-Timonen, R., Tolvanen, R., Lundén, J. & Korkeala, H. (2007) An 8-year surveillance of the diversity and persistence of *Listeria monocytogenes* in a chilled food processing plant analyzed by amplified fragment length polymorphism. *J Food Prot* 70: 1866-1873.

- Kim, M., Weigand, M. R., Oh, S., Hatt, J. K., Krishnan, R., Tezel, U., Pavlostathis, S. G. & Konstantinidis, K. T. (2018) Widely used benzalkonium chloride disinfectants can promote antibiotic resistance. *Appl Environ Microbiol* 84: e01201-18.
- Kirby, J. R. (2009) Chemotaxis-like regulatory systems: unique roles in diverse bacteria. *Annu Rev Microbiol* 63: 45-59.
- Knudsen, G. M., Olsen, J. E. & Dons, L. (2004) Characterization of DegU, a response regulator in *Listeria monocytogenes*, involved in regulation of motility and contributes to virulence. *FEMS Microbiol Lett* 240: 171-179.
- Ko, R., Smith, L. T. & Smith, G. M. (1994) Glycine betaine confers enhanced osmotolerance and cryotolerance on *Listeria monocytogenes*. *J Bacteriol* 176: 426-431.
- Kolstad, J., Rørvik, L. M. & Granum, P. E. (1991) Characterization of plasmids from *Listeria* sp. *Int J Food Microbiol* 12: 123-131.
- Koopmans, M. M., Brouwer, M. C., Bijlsma, M. W., Bovenkerk, S., Keijzers, W., van der Ende, A. & van de Beek, D. (2013) *Listeria monocytogenes* sequence type 6 and increased rate of unfavorable outcome in meningitis: epidemiologic cohort study. *Clin Infect Dis* 57: 247-253.
- Korsak, D., Chmielowska, C., Szuplewska, M. & Bartosik, D. (2019) Prevalence of plasmid-borne benzalkonium chloride resistance cassette *bcrABC* and cadmium resistance *cadA* genes in nonpathogenic *Listeria* spp. isolated from food and food-processing environments. *Int J Food Microbiol* 290: 247-253.
- Kovacevic, J., Ziegler, J., Walecka-Zacharska, E., Reimer, A., Kitts, D. D. & Gilmour, M. W. (2016) Tolerance of *Listeria monocytogenes* to quaternary ammonium sanitizers is mediated by a novel efflux pump encoded by *emrE*. *Appl Environ Microbiol* 82: 939-953.
- Krell, T., Lacal, J., Busch, A., Silva-Jiménez, H., Guazzaroni, M.-E. & Ramos, J. L. (2010) Bacterial sensor kinases: diversity in the recognition of environmental signals. *Annu Rev Microbiol* 64: 539-559.
- Kremer, P. H. C., Lees, J. A., Koopmans, M. M., Ferwerda, B., Arends, A. W. M., Feller, M. M., Schipper, K., Valls Seron, M., van der Ende, A., Brouwer, M. C., van de Beek, D. & Bentley, S. D. (2017) Benzalkonium tolerance genes and outcome in *Listeria monocytogenes* meningitis. *Clin Microbiol Infect* 23: 265.e1-265.e7.
- Kroll, R. G. & Patchett, R. A. (1992) Induced acid tolerance in *Listeria monocytogenes*. *Lett Appl Microbiol* 14: 224-227.
- Krulwich, T. A., Sachs, G. & Padan, E. (2011) Molecular aspects of bacterial pH sensing and homeostasis. *Nat Rev Microbiol* 9: 330-343.
- Kuenne, C., Voget, S., Pischmarov, J., Oehm, S., Goesmann, A., Daniel, R., Hain, T. & Chakraborty, T. (2010) Comparative analysis of plasmids in the genus *Listeria*. *PloS One* 5: e12511.
- Kwong, J. C., Mercoulia, K., Tomita, T., Easton, M., Li, H. Y., Bulach, D. M.,

- Stinear, T. P., Seemann, T. & Howden, B. P. (2016) Prospective whole-genome sequencing enhances national surveillance of *Listeria monocytogenes*. *J Clin Microbiol* 54: 333-342.
- Lang Halter, E., Neuhaus, K. & Scherer, S. (2013) *Listeria weihenstephanensis* sp. nov., isolated from the water plant *Lemna trisulca* taken from a freshwater pond. *Int J Syst Evol Microbiol* 63: 641-647.
- Larsen, H. E. & Seeliger, H. P. R. (1966) A mannitol fermenting *Listeria*: *Listeria grayi* sp. n. In *Proceedings of the third international symposium on listeriosis*, 13-16 July 1966, Bilthoven, The Netherlands, p. 35.
- Larsen, M. H., Kallipolitis, B. H., Christiansen, J. K., Olsen, J. E. & Ingmer, H. (2006) The response regulator ResD modulates virulence gene expression in response to carbohydrates in *Listeria monocytogenes*. *Mol Microbiol* 61: 1622-1635.
- Laub, M. T. & Goulian, M. (2007) Specificity in two-component signal transduction pathways. *Annu Rev Genet* 41: 121-145.
- Lauer, P., Chow, M. Y. N., Loessner, M. J., Portnoy, D. A. & Calendar, R. (2002) Construction, characterization, and use of two *Listeria monocytogenes* site-specific phage integration vectors. *J Bacteriol* 184: 4177-4186.
- Lawrence, L. M., Harvey, J. & Gilmour, A. (1993) Development of a random amplification of polymorphic DNA typing method for *Listeria monocytogenes*. *Appl Environ Microbiol* 59: 3117-3119.
- Lebrun, M., Audurier, A. & Cossart, P. (1994) Plasmid-borne cadmium resistance genes in *Listeria monocytogenes* are similar to *cadA* and *cadC* of *Staphylococcus aureus* and are induced by cadmium. *J Bacteriol* 176: 3040-3048.
- Lebrun, M., Loulergue, J., Chaslus-Dancla, E. & Audurier, A. (1992) Plasmids in *Listeria monocytogenes* in relation to cadmium resistance. *Appl Environ Microbiol* 58: 3183-3186.
- Leclercq, A., Clermont, D., Bizet, C., Grimont, P. A. D., Le Flèche-Matéos, A., Roche, S. M., Buchrieser, C., Cadet-Daniel, V., Le Monnier, A., Lecuit, M. & Allerberger, F. (2010) *Listeria rocourtiae* sp. nov. *Int J Syst Evol Microbiol* 60: 2210-2214.
- Lecuit, M. (2007) Human listeriosis and animal models. *Microbes Infect* 9: 1216-1225.
- Lee, J.-W. & Helmann, J. D. (2006) The PerR transcription factor senses H₂O₂ by metal-catalysed histidine oxidation. *Nature* 440: 363-367.
- Lees, J. A., Vehkala, M., Välimäki, N., Harris, S. R., Chewapreecha, C., Croucher, N. J., Marttinen, P., Davies, M. R., Steer, A. C., Tong, S. Y. C., Honkela, A., Parkhill, J., Bentley, S. D. & Corander, J. (2016) Sequence element enrichment analysis to determine the genetic basis of bacterial phenotypes. *Nat Commun* 7: 12797.
- Lessing, M. P. A., Curtis, G. D. W. & Bowler, I. C. J. (1994) *Listeria ivanovii* infection. *J Infect* 29: 230-231.
- Lianou, A. & Koutsoumanis, K. P. (2013) Strain variability of

- the behavior of foodborne bacterial pathogens: a review. *Int J Food Microbiol* 167: 310-321.
- Lianou, A., Stopforth, J. D., Yoon, Y., Wiedmann, M. & Sofos, J. N. (2006) Growth and stress resistance variation in culture broth among *Listeria monocytogenes* strains of various serotypes and origins. *J Food Prot* 69: 2640-2647.
- Linder, P. & Jankowsky, E. (2011) From unwinding to clamping – the DEAD box RNA helicase family. *Nat Rev Mol Cell Biol* 12: 505-516.
- Lindstedt, B.-A., Tham, W., Danielsson-Tham, M.-L., Vardund, T., Helmersson, S. & Kapperud, G. (2008) Multiple-locus variable-number tandem-repeats analysis of *Listeria monocytogenes* using multicolour capillary electrophoresis and comparison with pulsed-field gel electrophoresis typing. *J Microbiol Methods* 72: 141-148.
- Lindström, M., Dahlsten, E., Söderholm, H., Selby, K., Somervuo, P., Heap, J. T., Minton, N. P. & Korkeala, H. (2012) Involvement of two-component system CBO0366/CBO0365 in the cold shock response and growth of group I (proteolytic) *Clostridium botulinum* ATCC 3502 at low temperatures. *Appl Environ Microbiol* 78: 5466-5470.
- Linnan, M. J., Mascola, L., Lou, X. D., Goulet, V., May, S., Salminen, C., Hird, D. W., Yonekura, M. L., Hayes, P., Weaver, R., Audurier, A., Plikaytis, B. D., Fannin, S. L., Kleks, A. & Broome, C. V. (1988) Epidemic listeriosis associated with Mexican-style cheese. *N Engl J Med* 319: 823-828.
- Liu, D., Lawrence, M. L., Wiedmann, M., Gorski, L., Mandrell, R. E., Ainsworth, A. J. & Austin, F. W. (2006) *Listeria monocytogenes* subgroups IIIA, IIIB, and IIIC delineate genetically distinct populations with varied pathogenic potential. *J Clin Microbiol* 44: 4229-4233.
- Liu, S., Graham, J. E., Bigelow, L., Morse, P. D., II & Wilkinson, B. J. (2002) Identification of *Listeria monocytogenes* genes expressed in response to growth at low temperature. *Appl Environ Microbiol* 68: 1697-1705.
- Liu, Y., Du, F.-L., Xiang, T.-X., Wan, L.-G., Wei, D.-D., Cao, X.-W. & Zhang, W. (2019) High prevalence of plasmid-mediated quinolone resistance determinants among serotype K1 hypervirulent *Klebsiella pneumoniae* isolates in China. *Microb Drug Resist* Online ahead of print: Jan 7 2019, <https://doi-org.libproxy.helsinki.fi/10.1089/mdr.2018.0173> Accessed at 27 Mar 2019
- Liu, Y.-Y., Wang, Y., Walsh, T. R., Yi, L.-X., Zhang, R., Spencer, J., Doi, Y., Tian, G., Dong, B., Huang, X., Yu, L.-F., Gu, D., Ren, H., Chen, X., Lv, L., He, D., Zhou, H., Liang, Z., Liu, J.-H. & Shen, J. (2016) Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 16: 161-168.
- Loepfe, C., Raimann, E., Stephan, R. & Tasara, T. (2010) Reduced host cell

- invasiveness and oxidative stress tolerance in double and triple *csp* gene family deletion mutants of *Listeria monocytogenes*. *Foodborne Pathog Dis* 7: 775-783.
- Lois, A. F., Weinstein, M., Ditta, G. S. & Helinski, D. R. (1993) Autophosphorylation and phosphatase activities of the oxygen-sensing protein FixL of *Rhizobium meliloti* are coordinately regulated by oxygen. *J Biol Chem* 268: 4370-4375.
- Lopez-Valladares, G., Danielsson-Tham, M.-L. & Tham, W. (2018) Implicated food products for listeriosis and changes in serovars of *Listeria monocytogenes* affecting humans in recent decades. *Foodborne Pathog Dis* 15: 387-397.
- López-Prieto, M. D., Aller García, A. I., Alcaraz García, S. & López Cepero, J. (2000) Liver abscess due to *Listeria monocytogenes*. *Clin Microbiol Infect* 6: 226-227.
- Lou, Y. & Yousef, A. E. (1997) Adaptation to sublethal environmental stresses protects *Listeria monocytogenes* against lethal preservation factors. *Appl Environ Microbiol* 63: 1252-1255.
- Lukinmaa, S., Miettinen, M., Nakari, U.-M., Korkeala, H. & Siitonen, A. (2003) *Listeria monocytogenes* isolates from invasive infections: variation of sero- and genotypes during an 11-year period in Finland. *J Clin Microbiol* 41: 1694-1700.
- Lundén, J., Autio, T., Markkula, A., Hellström, S. & Korkeala, H. (2003a) Adaptive and cross-adaptive responses of persistent and non-persistent *Listeria monocytogenes* strains to disinfectants. *Int J Food Microbiol* 82: 265-272.
- Lundén, J. M., Autio, T. J., Sjöberg, A. M. & Korkeala, H. J. (2003b) Persistent and nonpersistent *Listeria monocytogenes* contamination in meat and poultry processing plants. *J Food Prot* 66: 2062-2069.
- Lundén, J., Tolvanen, R. & Korkeala, H. (2008) Acid and heat tolerance of persistent and nonpersistent *Listeria monocytogenes* food plant strains. *Lett Appl Microbiol* 46: 276-280.
- Lyautey, E., Hartmann, A., Pagotto, F., Tyler, K., Lapen, D. R., Wilkes, G., Piveteau, P., Rieu, A., Robertson, W. J., Medeiros, D. T., Edge, T. A., Gannon, V. & Topp, E. (2007a) Characteristics and frequency of detection of fecal *Listeria monocytogenes* shed by livestock, wildlife, and humans. *Can J Microbiol* 53: 1158-1167.
- Lyautey, E., Lapen, D. R., Wilkes, G., McCleary, K., Pagotto, F., Tyler, K., Hartmann, A., Piveteau, P., Rieu, A., Robertson, W. J., Medeiros, D. T., Edge, T. A., Gannon, V. & Topp, E. (2007b) Distribution and characteristics of *Listeria monocytogenes* isolates from surface waters of the South Nation River watershed, Ontario, Canada. *Appl Environ Microbiol* 73: 5401-5410.
- Lyrstis, M., Bryant, A. E., Sloan, J., Awad, M. M., Nisbet, I. T., Stevens, D. L. & Rood, J. I. (1994) Identification and molecular analysis of a locus that regulates extracellular toxin production in *Clostridium perfringens*. *Mol Microbiol* 12: 761-777.

- Lyytikäinen, O., Autio, T., Maijala, R., Ruutu, P., Honkanen-Buzalski, T., Miettinen, M., Hatakka, M., Mikkola, J., Anttila, V.-J., Johansson, T., Rantala, L., Aalto, T., Korkeala, H. & Siitonen, A. (2000) An outbreak of *Listeria monocytogenes* serotype 3a infections from butter in Finland. *J Infect Dis* 181: 1838-1841.
- Löfmark, S., Jernberg, C., Billström, H., Andersson, D. I. & Edlund, C. (2008) Restored fitness leads to long-term persistence of resistant *Bacteroides* strains in the human intestine. *Anaerobe* 14: 157-160.
- Ma, L., Zhang, G. & Doyle, M. P. (2011) Green fluorescent protein labeling of *Listeria*, *Salmonella*, and *Escherichia coli* O157:H7 for safety-related studies. *PloS One* 6: e18083.
- MacGowan, A. P., Bowker, K., McLauchlin, J., Bennett, P. M. & Reeves, D. S. (1994) The occurrence and seasonal changes in the isolation of *Listeria* spp. in shop bought food stuffs, human faeces, sewage and soil from urban sources. *Int J Food Microbiol* 21: 325-334.
- Mackey, B. M., Pritchett, C., Norris, A. & Mead, G. C. (1990) Heat resistance of *Listeria*: strain differences and effects of meat type and curing salts. *Lett Appl Microbiol* 10: 251-255.
- Maijala, R., Lyytikäinen, O., Johansson, T., Autio, T., Aalto, T., Haavisto, L. & Honkanen-Buzalski, T. (2001) Exposure of *Listeria monocytogenes* within an epidemic caused by butter in Finland. *Int J Food Microbiol* 70: 97-109.
- Mandin, P., Fsihi, H., Dussurget, O., Vergassola, M., Milohanic, E., Toledo-Arana, A., Lasa, I., Johansson, J. & Cossart, P. (2005) VirR, a response regulator critical for *Listeria monocytogenes* virulence. *Mol Microbiol* 57: 1367-1380.
- Margolles, A. & de los Reyes-Gavilán, C. G. (1998) Characterization of plasmids from *Listeria monocytogenes* and *Listeria innocua* strains isolated from short-ripened cheeses. *Int J Food Microbiol* 39: 231-236.
- Markkula, A., Autio, T., Lundén, J. & Korkeala, H. (2005) Raw and processed fish show identical *Listeria monocytogenes* genotypes with pulsed-field gel electrophoresis. *J Food Prot* 68: 1228-1231.
- Markkula, A., Lindström, M., Johansson, P., Björkroth, J. & Korkeala, H. (2012a) Roles of four putative DEAD-box RNA helicase genes in growth of *Listeria monocytogenes* EGD-e under heat, pH, osmotic, ethanol, and oxidative stress conditions. *Appl Environ Microbiol* 78: 6875-6882.
- Markkula, A., Mattila, M., Lindström, M. & Korkeala, H. (2012b) Genes encoding putative DEAD-box RNA helicases in *Listeria monocytogenes* EGD-e are needed for growth and motility at 3°C. *Environ Microbiol* 14: 2223-2232.
- Martin, P. K., Li, T., Sun, D., Biek, D. P. & Schmid, M. B. (1999) Role in cell permeability of an essential two-component system in *Staphylococcus aureus*. *J Bacteriol* 181: 3666-3673.
- Mascher, G., Derman, Y., Kirk, D. G., Palonen, E., Lindström,

- M. & Korkeala, H. (2014) The CLO3403/CLO3404 two-component system of *Clostridium botulinum* E1 Beluga is important for cold shock response and growth at low temperatures. *Appl Environ Microbiol* 80: 399-407.
- Mastronicolis, S. K., German, J. B., Megoulas, N., Petrou, E., Foka, P. & Smith, G. M. (1998) Influence of cold shock on the fatty-acid composition of different lipid classes of the food-borne pathogen *Listeria monocytogenes*. *Food Microbiol* 15: 299-306.
- Mattila, M., Lindström, M., Somervuo, P., Markkula, A. & Korkeala, H. (2011) Role of *flhA* and *motA* in growth of *Listeria monocytogenes* at low temperatures. *Int J Food Microbiol* 148: 177-183.
- Mauder, N., Williams, T., Fritsch, F., Kuhn, M. & Beier, D. (2008) Response regulator DegU of *Listeria monocytogenes* controls temperature-responsive flagellar gene expression in its unphosphorylated state. *J Bacteriol* 190: 4777-4781.
- Maury, M. M., Tsai, Y.-H., Charlier, C., Touchon, M., Chenal-Francisque, V., Leclercq, A., Criscuolo, A., Gaultier, C., Roussel, S., Brisabois, A., Disson, O., Rocha, E. P. C., Brisse, S. & Lecuit, M. (2016) Uncovering *Listeria monocytogenes* hypervirulence by harnessing its biodiversity. *Nat Genet* 48: 308-313.
- Mazurier, S.-I. & Wernars, K. (1992) Typing of *Listeria* strains by random amplification of polymorphic DNA. *Res Microbiol* 143: 499-505.
- McDonnell, G. & Russell, A. D. (1999) Antiseptics and disinfectants: activity, action, and resistance. *Clin Microbiol Rev* 12: 147-179.
- McLauchlin, J. (1990) Distribution of serovars of *Listeria monocytogenes* isolated from different categories of patients with listeriosis. *Eur J Clin Microbiol Infect Dis* 9: 210-213.
- McLauchlin, J. (1996) The relationship between *Listeria* and listeriosis. *Food Control* 7: 187-193.
- McLauchlin, J., Hampton, M. D., Shah, S., Threlfall, E. J., Wieneke, A. A. & Curtis, G. D. W. (1997) Subtyping of *Listeria monocytogenes* on the basis of plasmid profiles and arsenic and cadmium susceptibility. *J Appl Microbiol* 83: 381-388.
- McLauchlin, J. & Low, J. C. (1994) Primary cutaneous listeriosis in adults: an occupational disease of veterinarians and farmers. *Vet Rec* 135: 615-617.
- McLauchlin, J. & Rees, C. E. D. (2015) Genus *Listeria*. In *Bergey's manual of systematics of archaea and bacteria*. Whitman, W. B., de Vos, P., Dedysh, S., Hedlund, B., Kämpfer, P., Rainey, F., Trujillo, M. E., Bowman, J. P., Brown, D. R., Glöckner, F. O., Oren, A., Paster, B. J., Wade, W., Ward, N., Busse, H.-J., Reysenbach, A.-L. (eds). 3rd edn. New Jersey, USA, John Wiley & Sons, Inc., pp. 1-29.
- Mead, P. S., Slutsker, L., Dietz, V., McCaig, L. F., Bresee, J. S., Shapiro, C., Griffin, P. M. & Tauxe, R. V. (1999) Food-related illness and death in the United States. *Emerg Infect Dis* 5: 607-625.
- Mereghetti, L., Quentin, R., Marquet-Van Der Mee, N. & Audurier, A. (2000) Low

- sensitivity of *Listeria monocytogenes* to quaternary ammonium compounds. *Appl Environ Microbiol* 66: 5083-5086.
- Metselaar, K. I., den Besten, H. M. W., Abee, T., Moezelaar, R. & Zwietering, M. H. (2013) Isolation and quantification of highly acid resistant variants of *Listeria monocytogenes*. *Int J Food Microbiol* 166: 508-514.
- Miettinen, M. K., Björkroth, K. J. & Korkeala, H. J. (1999a) Characterization of *Listeria monocytogenes* from an ice cream plant by serotyping and pulsed-field gel electrophoresis. *Int J Food Microbiol* 46: 187-192.
- Miettinen, M. K., Palmu, L., Björkroth, K. J. & Korkeala, H. (2001) Prevalence of *Listeria monocytogenes* in broilers at the abattoir, processing plant, and retail level. *J Food Prot* 64: 994-999.
- Miettinen, M. K., Siitonen, A., Heiskanen, P., Haajanen, H., Björkroth, K. J. & Korkeala, H. J. (1999b) Molecular epidemiology of an outbreak of febrile gastroenteritis caused by *Listeria monocytogenes* in cold-smoked rainbow trout. *J Clin Microbiol* 37: 2358-2360.
- Miladi, H., Elabed, H., Slama, R. B., Rhim, A. & Bakhrouf, A. (2017) Molecular analysis of the role of osmolyte transporters *opuCA* and *betL* in *Listeria monocytogenes* after cold and freezing stress. *Arch Microbiol* 199: 259-265.
- Mishra, S. & Imlay, J. (2012) Why do bacteria use so many enzymes to scavenge hydrogen peroxide? *Arch Biochem Biophys* 525: 145-160.
- Mitrophanov, A. Y. & Groisman, E. A. (2008) Signal integration in bacterial two-component regulatory systems. *Genes Dev* 22: 2601-2611.
- Mizuno, T. (1998) His-Asp phosphotransfer signal transduction. *J Biochem* 123: 555-563.
- Mizuno, T., Kaneko, T. & Tabata, S. (1996) Compilation of all genes encoding bacterial two-component signal transducers in the genome of the cyanobacterium, *Synechocystis* sp. strain PCC 6803. *DNA Res* 3: 407-414.
- Mohedano, M. L., Overweg, K., de la Fuente, A., Reuter, M., Altabe, S., Mulholland, F., de Mendoza, D., López, P. & Wells, J. M. (2005) Evidence that the essential response regulator YycF in *Streptococcus pneumoniae* modulates expression of fatty acid biosynthesis genes and alters membrane composition. *J Bacteriol* 187: 2357-2367.
- Moura, A., Criscuolo, A., Pouseele, H., Maury, M. M., Leclercq, A., Tarr, C., Björkman, J. T., Dallman, T., Reimer, A., Enouf, V., Larssonneur, E., Carleton, H., Bracq-Dieye, H., Katz, L. S., Jones, L., Touchon, M., Tourdjman, M., Walker, M., Stroika, S., Cantinelli, T., Chenal-Francisque, V., Kucerova, Z., Rocha, E. P. C., Nadon, C., Grant, K., Nielsen, E. M., Pot, B., Gerner-Smidt, P., Lecuit, M. & Brisse, S. (2016) Whole genome-based population biology and epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* 2: 16185.
- Müller, A., Rychli, K., Muhterem-Uyar, M., Zaiser, A., Stessl, B., Guinane, C. M., Cotter, P. D., Wagner, M. &

- Schmitz-Esser, S. (2013) Tn6188 - a novel transposon in *Listeria monocytogenes* responsible for tolerance to benzalkonium chloride. *PLoS One* 8: e76835.
- Mullapudi, S., Siletzky, R. M. & Kathariou, S. (2008) Heavy-metal and benzalkonium chloride resistance of *Listeria monocytogenes* isolates from the environment of turkey-processing plants. *Appl Environ Microbiol* 74: 1464-1468.
- Murphy, M., Corcoran, D., Buckley, J. F., O'Mahony, M., Whyte, P. & Fanning, S. (2007) Development and application of Multiple-Locus Variable number of tandem repeat Analysis (MLVA) to subtype a collection of *Listeria monocytogenes*. *Int J Food Microbiol* 115: 187-194.
- Murray, E. G. D., Webb, R. A. & Swann, M. B. R. (1926) A disease of rabbits characterized by large mononuclear leucocytosis, caused by a hitherto undescribed bacillus: *Bacterium monocytogenes* (n. sp.). *J Pathol Bacteriol* 29: 407-439.
- Mylonakis, E., Hohmann, E. L. & Calderwood, S. B. (1998) Central nervous system infection with *Listeria monocytogenes*. 33 years' experience at a general hospital and review of 776 episodes from the literature. *Medicine (Baltimore)* 77: 313-336.
- Mylonakis, E., Paliou, M., Hohmann, E. L., Calderwood, S. B. & Wing, E. J. (2002) Listeriosis during pregnancy: a case series and review of 222 cases. *Medicine (Baltimore)* 81: 260-269.
- Nadon, C. A., Woodward, D. L., Young, C., Rodgers, F. G. & Wiedmann, M. (2001) Correlations between molecular subtyping and serotyping of *Listeria monocytogenes*. *J Clin Microbiol* 39: 2704-2707.
- Nair, S., Derré, I., Msadek, T., Gaillot, O. & Berche, P. (2000) CtsR controls class III heat shock gene expression in the human pathogen *Listeria monocytogenes*. *Mol Microbiol* 35: 800-811.
- Nakashima, K., Sugiura, A., Momoi, H. & Mizuno, T. (1992) Phosphotransfer signal transduction between two regulatory factors involved in the osmoregulated *kdp* operon in *Escherichia coli*. *Mol Microbiol* 6: 1777-1784.
- Narberhaus, F., Waldminghaus, T. & Chowdhury, S. (2006) RNA thermometers. *FEMS Microbiol Rev* 30: 3-16.
- Nelson, K. E., Fouts, D. E., Mongodin, E. F., Ravel, J., DeBoy, R. T., Kolonay, J. F., Rasko, D. A., Angiuoli, S. V., Gill, S. R., Paulsen, I. T., Peterson, J., White, O., Nelson, W. C., Niernman, W., Beanan, M. J., Brinkac, L. M., Daugherty, S. C., Dodson, R. J., Durkin, A. S., Madupu, R., Haft, D. H., Selengut, J., Van Aken, S., Khouri, H., Fedorova, N., Forberger, H., Tran, B., Kathariou, S., Wonderling, L. D., Uhlich, G. A., Bayles, D. O., Luchansky, J. B. & Fraser, C. M. (2004) Whole genome comparisons of serotype 4b and 1/2a strains of the food-borne pathogen *Listeria monocytogenes* reveal new insights into the core genome components of this species. *Nucleic Acids Res* 32: 2386-2395.

- NicAogáin, K. & O'Byrne, C. P. (2016) The role of stress and stress adaptations in determining the fate of the bacterial pathogen. *Front Microbiol* 7: 1865.
- Nielsen, P. K., Andersen, A. Z., Mols, M., van der Veen, S., Abee, T. & Kallipolitis, B. H. (2012) Genome-wide transcriptional profiling of the cell envelope stress response and the role of LisRK and CesRK in *Listeria monocytogenes*. *Microbiology* 158: 963-974.
- Nightingale, K. K., Fortes, E. D., Ho, A. J., Schukken, Y. H., Grohn, Y. T. & Wiedmann, M. (2005a) Evaluation of farm management practices as risk factors for clinical listeriosis and fecal shedding of *Listeria monocytogenes* in ruminants. *J Am Vet Med Assoc* 227: 1808-1814.
- Nightingale, K. K., Schukken, Y. H., Nightingale, C. R., Fortes, E. D., Ho, A. J., Her, Z., Grohn, Y. T., McDonough, P. L. & Wiedmann, M. (2004) Ecology and transmission of *Listeria monocytogenes* infecting ruminants and in the farm environment. *Appl Environ Microbiol* 70: 4458-4467.
- Nightingale, K. K., Windham, K. & Wiedmann, M. (2005b) Evolution and molecular phylogeny of *Listeria monocytogenes* isolated from human and animal listeriosis cases and foods. *J Bacteriol* 187: 5537-5551.
- Nikitas, G., Deschamps, C., Disson, O., Niault, T., Cossart, P. & Lecuit, M. (2011) Transcytosis of *Listeria monocytogenes* across the intestinal barrier upon specific targeting of goblet cell accessible E-cadherin. *J Exp Med* 208: 2263-2277.
- Nilsson, R. E., Ross, T. & Bowman, J. P. (2011) Variability in biofilm production by *Listeria monocytogenes* correlated to strain origin and growth conditions. *Int J Food Microbiol* 150: 14-24.
- Nocera, D., Bannerman, E., Rocourt, J., Jatton-Ogay, K. & Bille, J. (1990) Characterization by DNA restriction endonuclease analysis of *Listeria monocytogenes* strains related to the Swiss epidemic of listeriosis. *J Clin Microbiol* 28: 2259-2263.
- Nolan, D. A., Chamblin, D. C. & Troller, J. A. (1992) Minimal water activity levels for growth and survival of *Listeria monocytogenes* and *Listeria innocua*. *Int J Food Microbiol* 16: 323-335.
- Norton, D. M., McCamey, M. A., Gall, K. L., Scarlett, J. M., Boor, K. J. & Wiedmann, M. (2001) Molecular studies on the ecology of *Listeria monocytogenes* in the smoked fish processing industry. *Appl Environ Microbiol* 67: 198-205.
- Nyfeldt, A. (1929) Étiologie de la mononucleose infectieuse. *C R Soc Biol* 101: 590-591.
- O'Driscoll, B., Gahan, C. G. & Hill, C. (1996) Adaptive acid tolerance response in *Listeria monocytogenes*: isolation of an acid-tolerant mutant which demonstrates increased virulence. *Appl Environ Microbiol* 62: 1693-1698.
- Oh, D.-H. & Marshall, D. L. (1993) Antimicrobial activity of ethanol, glycerol monolaurate or lactic acid against *Listeria monocytogenes*. *Int J Food Microbiol* 20: 239-246.
- Ohtani, K., Hirakawa, H., Tashiro, K., Yoshizawa, S., Kuhara, S.

- & Shimizu, T. (2010) Identification of a two-component VirR/VirS regulon in *Clostridium perfringens*. *Anaerobe* 16: 258-264.
- Ojeniyi, B., Høiby, N. & Rosdahl, V. T. (1991) Genome fingerprinting as a typing method used on polyagglutinable *Pseudomonas aeruginosa* isolates from cystic fibrosis patients. *APMIS* 99: 492-498.
- Ojeniyi, B., Wegener, H. C., Jensen, N. E. & Bisgaard, M. (1996) *Listeria monocytogenes* in poultry and poultry products: epidemiological investigations in seven Danish abattoirs. *J Appl Bacteriol* 80: 395-401.
- Olsen, K. N., Larsen, M. H., Gahan, C. G. M., Kallipolitis, B., Wolf, X. A., Rea, R., Hill, C. & Ingmer, H. (2005) The Dps-like protein Fri of *Listeria monocytogenes* promotes stress tolerance and intracellular multiplication in macrophage-like cells. *Microbiology* 151: 925-933.
- Ooi, S. T. & Lorber, B. (2005) Gastroenteritis due to *Listeria monocytogenes*. *Clin Infect Dis* 40: 1327-1332.
- Orsi, R. H., den Bakker, H. C. & Wiedmann, M. (2011) *Listeria monocytogenes* lineages: genomics, evolution, ecology, and phenotypic characteristics. *Int J Med Microbiol* 301: 79-96.
- Orsi, R. H., Sun, Q. & Wiedmann, M. (2008) Genome-wide analyses reveal lineage specific contributions of positive selection and recombination to the evolution of *Listeria monocytogenes*. *BMC Evol Biol* 8: 233.
- Overbeek, R., Begley, T., Butler, R. M., Choudhuri, J. V., Chuang, H.-Y., Cohoon, M., de Crécy-Lagard, V., Diaz, N., Disz, T., Edwards, R., Fonstein, M., Frank, E. D., Gerdes, S., Glass, E. M., Goesmann, A., Hanson, A., Iwata-Reuyl, D., Jensen, R., Jamshidi, N., Krause, L., Kubal, M., Larsen, N., Linke, B., McHardy, A. C., Meyer, F., Newweger, H., Olsen, G., Olson, R., Osterman, A., Portnoy, V., Pusch, G. D., Rodionov, D. A., Rückert, C., Steiner, J., Stevens, R., Thiele, I., Vassieva, O., Ye, Y., Zagnitko, O. & Vonstein, V. (2005) The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Res* 33: 5691-5702.
- Palonen, E., Lindström, M., Karttunen, R., Somervuo, P. & Korkeala, H. (2011) Expression of signal transduction system encoding genes of *Yersinia pseudotuberculosis* IP32953 at 28°C and 3°C. *PLoS One* 6: e25063.
- Parihar, V. S., Lopez-Valladares, G., Danielsson-Tham, M.-L., Peiris, I., Helmersson, S., Unemo, M., Andersson, B., Arneborn, M., Bannerman, E., Barbuddhe, S., Bille, J., Hajdu, L., Jacquet, C., Johansson, C., Löfdahl, M., Möllerberg, G., Ringberg, H., Rocourt, J., Tjernberg, I., Rosing, J., Henriques-Normark, B. & Tham, W. (2008) Characterization of human invasive isolates of *Listeria monocytogenes* in Sweden 1986-2007. *Foodborne Pathog Dis* 5: 755-761.

- Patchett, R. A., Kelly, A. F. & Kroll, R. G. (1992) Effect of sodium chloride on the intracellular solute pools of *Listeria monocytogenes*. *Appl Environ Microbiol* 58: 3959-3963.
- Pauly, T. M. & Tham, W. A. (2003) Survival of *Listeria monocytogenes* in wilted and additive-treated grass silage. *Acta Vet Scand* 44: 73-86.
- Pérez-Díaz, J. C., Vicente, M. F. & Baquero, F. (1982) Plasmids in *Listeria*. *Plasmid* 8: 112-118.
- Périchon, B., Bogaerts, P., Lambert, T., Frangeul, L., Courvalin, P. & Galimand, M. (2008) Sequence of conjugative plasmid pIP1206 mediating resistance to aminoglycosides by 16S rRNA methylation and to hydrophilic fluorquinolones by efflux. *Antimicrob Agents Chemother* 52: 2581-2592.
- Perrin, M., Bemer, M. & Delamare, C. (2003) Fatal case of *Listeria innocua* bacteremia. *J Clin Microbiol* 41: 5308-5309.
- Peterkin, P. I., Gardiner, M.-A., Malik, N. & Idziak, E. S. (1992) Plasmids in *Listeria monocytogenes* and other *Listeria* species. *Can J Microbiol* 38: 161-164.
- Pfaffl, M. W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29: e45.
- Phan-Thanh, L. & Gormon, T. (1995) Analysis of heat and cold shock proteins in *Listeria* by two-dimensional electrophoresis. *Electrophoresis* 16: 444-450.
- Phan-Thanh, L. & Mahouin, F. (1999) A proteomic approach to study the acid response in *Listeria monocytogenes*. *Electrophoresis* 20: 2214-2224.
- Phan-Thanh, L., Mahouin, F. & Aligé, S. (2000) Acid responses of *Listeria monocytogenes*. *Int J Food Microbiol* 55: 121-126.
- Piffaretti, J.-C., Kressebuch, H., Aeschbacher, M., Bille, J., Bannerman, E., Musser, J. M., Selander, R. K. & Rocourt, J. (1989) Genetic characterization of clones of the bacterium *Listeria monocytogenes* causing epidemic disease. *Proc Natl Acad Sci U S A* 86: 3818-3822.
- Pinheiro, J., Lisboa, J., Pombinho, R., Carvalho, F., Carreaux, A., Brito, C., Pöntinen, A., Korkeala, H., dos Santos, N. M. S., Morais-Cabral, J. H., Sousa, S. & Cabanes, D. (2018) MouR controls the expression of the *Listeria monocytogenes* Agr system and mediates virulence. *Nucleic Acids Res* 46: 9338-9352.
- Pinner, R. W., Schuchat, A., Swaminathan, B., Hayes, P. S., Deaver, K. A., Weaver, R. E., Plikaytis, B. D., Reeves, M., Broome, C. V., Wenger, J. D., Ajello, G., Przybyszewski, V., Malcolm, G., Ransom, R., Graves, L. M., Egal, M., Pierce, R., Anderson, G., Stone, E., Krauss, K., Reingold, A., Castillon, M., Mascola, L., Harvey, C., Stull, T., Stephens, D., Farley, M., Archer, P., Strack, J., Istre, G., Rados, M., Taylor, J. & Lefkowitz, L. (1992) Role of foods in sporadic listeriosis II. Microbiologic and epidemiologic investigation. *JAMA* 267: 2046-2050.
- Pirie, J. H. H. (1927) A new disease of veld rodents, "Tiger River

- Disease". *Pub S Afr Inst Med Res* 3: 163-186.
- Pirie, J. H. H. (1940) *Listeria*: change of name for a genus bacteria. *Nature* 145: 264.
- Pitcher, D. G., Saunders, N. A. & Owen, R. J. (1989) Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. *Lett Appl Microbiol* 8: 151-156.
- Pittman, J. R., Buntyn, J. O., Posadas, G., Nanduri, B., Pendarvis, K. & Donaldson, J. R. (2014) Proteomic analysis of cross protection provided between cold and osmotic stress in *Listeria monocytogenes*. *J Proteome Res* 13: 1896-1904.
- Pizarro-Cerdá, J., Kühbacher, A. & Cossart, P. (2012) Entry of *Listeria monocytogenes* in mammalian epithelial cells: an updated view. *Cold Spring Harb Perspect Med* 2: a010009.
- Poirel, L., Villa, L., Bertini, A., Pitout, J. D., Nordmann, P. & Carattoli, A. (2007) Expanded-spectrum β -lactamase and plasmid-mediated quinolone resistance. *Emerg Infect Dis* 13: 803-805.
- Pontello, M., Guaita, A., Sala, G., Cipolla, M., Gattuso, A., Sonnessa, M. & Gianfranceschi, M. V. (2012) *Listeria monocytogenes* serotypes in human infections (Italy, 2000-2010). *Ann Ist Super Sanita* 48: 146-150.
- Pope, C., Kim, S.-K., Marzo, A., Williams, K., Jiang, J., Shen, H. & Lefrançois, L. (2001) Organ-specific regulation of the CD8 T cell response to *Listeria monocytogenes* infection. *J Immunol* 166: 3402-3409.
- Potel, J. (1953) Ätiologie der granulomatosis infantiseptica. *Wiss Z Martin-Luther-Univ Halle-Wittenberg* 3: 341-349.
- Poupel, O., Moyat, M., Groizeleau, J., Antunes, L. C. S., Gribaldo, S., Msadek, T. & Dubrac, S. (2016) Transcriptional analysis and subcellular protein localization reveal specific features of the essential WalKR system in *Staphylococcus aureus*. *PLoS One* 11: e0151449.
- Poyart-Salmeron, C., Carlier, C., Trieu-Cuot, P., Courtieu, A.-L. & Courvalin, P. (1990) Transferable plasmid-mediated antibiotic resistance in *Listeria monocytogenes*. *Lancet* 335: 1422-1426.
- Prieto, M., Martínez, C., Aguerre, L., Rocca, M. F., Cipolla, L. & Callejo, R. (2016) Antibiotic susceptibility of *Listeria monocytogenes* in Argentina. *Enferm Infecc Microbiol Clin* 34: 91-95.
- Püttmann, M., Ade, N. & Hof, H. (1993) Dependence of fatty acid composition of *Listeria* spp. on growth temperature. *Res Microbiol* 144: 279-283.
- Quereda, J. J., Dussurget, O., Nahori, M.-A., Ghozlane, A., Volant, S., Dillies, M.-A., Regnault, B., Kennedy, S., Mondot, S., Villoing, B., Cossart, P. & Pizarro-Cerdá, J. (2016) Bacteriocin from epidemic *Listeria* strains alters the host intestinal microbiota to favor infection. *Proc Natl Acad Sci U S A* 113: 5706-5711.
- Radoshevich, L. & Cossart, P. (2018) *Listeria monocytogenes*: towards a complete picture of its physiology and pathogenesis. *Nat Rev Microbiol* 16: 32-46.
- Raengpradub, S., Wiedmann, M. & Boor, K. J. (2008) Comparative analysis of the σ^B -dependent stress

- responses in *Listeria monocytogenes* and *Listeria innocua* strains exposed to selected stress conditions. *Appl Environ Microbiol* 74: 158-171.
- Ragon, M., Wirth, T., Hollandt, F., Lavenir, R., Lecuit, M., Le Monnier, A. & Brisse, S. (2008) A new perspective on *Listeria monocytogenes* evolution. *PLoS Pathog* 4: e1000146.
- Raines, L. J., Moss, C. W., Farshtchi, D. & Pittman, B. (1968) Fatty acids of *Listeria monocytogenes*. *J Bacteriol* 96: 2175-2177.
- Rakic-Martinez, M., Drevets, D. A., Dutta, V., Katic, V. & Kathariou, S. (2011) *Listeria monocytogenes* strains selected on ciprofloxacin or the disinfectant benzalkonium chloride exhibit reduced susceptibility to ciprofloxacin, gentamicin, benzalkonium chloride, and other toxic compounds. *Appl Environ Microbiol* 77: 8714-8721.
- Ramage, C. P., Low, J. C., McLauchlin, J. & Donachie, W. (1999) Characterisation of *Listeria ivanovii* isolates from the UK using pulsed-field gel electrophoresis. *FEMS Microbiol Lett* 170: 349-353.
- Ranjbar, R. & Farahani, O. (2017) The prevalence of plasmid-mediated quinolone resistance genes in *Escherichia coli* isolated from hospital wastewater sources in Tehran, Iran. *Iran J Public Health* 46: 1285-1291.
- Rankin, D. J., Rocha, E. P. C. & Brown, S. P. (2011) What traits are carried on mobile genetic elements, and why? *Heredity* 106: 1-10.
- Rapose, A., Lick, S. D. & Ismail, N. (2008) *Listeria grayi* bacteremia in a heart transplant recipient. *Transpl Infect Dis* 10: 434-436.
- Rasmussen, O. F., Skouboe, P., Dons, L., Rossen, L. & Olsen, J. E. (1995) *Listeria monocytogenes* exists in at least three evolutionary lines: evidence from flagellin, invasive associated protein and listeriolysin O genes. *Microbiology* 141: 2053-2061.
- Rea, R., Hill, C. & Gahan, C. G. M. (2005) *Listeria monocytogenes* PerR mutants display a small-colony phenotype, increased sensitivity to hydrogen peroxide, and significantly reduced murine virulence. *Appl Environ Microbiol* 71: 8314-8322.
- Read, T. D. & Massey, R. C. (2014) Characterizing the genetic basis of bacterial phenotypes using genome-wide association studies: a new direction for bacteriology. *Genome Med* 6: 109.
- Regan, E. J., Harrison, G. A. J., Butler, S., McLauchlin, J., Thomas, M. & Mitchell, S. (2005) Primary cutaneous listeriosis in a veterinarian. *Vet Rec* 157: 207.
- Reij, M. W., Den Aantrekker, E. D. & ILSI Europe Risk Analysis in Microbiology Task Force. (2004) Recontamination as a source of pathogens in processed foods. *Int J Food Microbiol* 91: 1-11.
- Renterghem, B. V., Huysman, F., Rygole, R. & Verstraete, W. (1991) Detection and prevalence of *Listeria monocytogenes* in the agricultural ecosystem. *J Appl Bacteriol* 71:211-217.
- Revez, J., Espinosa, L., Albiger, B., Leitmeyer, K. C., Struelens, M. J. & ECDC national microbiology focal points

- and experts group (2017) Survey on the use of whole-genome sequencing for infectious diseases surveillance: rapid expansion of European national capacities, 2015-2016. *Front Public Health* 5: 347.
- Ribeiro, V. B. & Destro, M. T. (2014) *Listeria monocytogenes* serotype 1/2b and 4b isolates from human clinical cases and foods show differences in tolerance to refrigeration and salt stress. *J Food Prot* 77: 1519-1526.
- Ribeiro, V. B., Mujahid, S., Orsi, R. H., Bergholz, T. M., Wiedmann, M., Boor, K. J. & Destro, M. T. (2014) Contributions of σ^B and PrfA to *Listeria monocytogenes* salt stress under food relevant conditions. *Int J Food Microbiol* 177: 98-108.
- Ribet, D. & Cossart, P. (2015) How bacterial pathogens colonize their hosts and invade deeper tissues. *Microbes Infect* 17: 173-183.
- Riedel, C. U., Monk, I. R., Casey, P. G., Waidmann, M. S., Gahan, C. G. M. & Hill, C. (2009) AgrD-dependent quorum sensing affects biofilm formation, invasion, virulence and global gene expression profiles in *Listeria monocytogenes*. *Mol Microbiol* 71: 1177-1189.
- Rieu, A., Weidmann, S., Garmyn, D., Piveteau, P. & Guzzo, J. (2007) agr system of *Listeria monocytogenes* EGD-e: role in adherence and differential expression pattern. *Appl Environ Microbiol* 73: 6125-6133.
- Ringus, D. L., Ivy, R. A., Wiedmann, M. & Boor, K. J. (2012) Salt stress-induced transcription of σ^B - and CtsR-regulated genes in persistent and non-persistent *Listeria monocytogenes* strains from food processing plants. *Foodborne Pathog Dis* 9: 198-206.
- Roberts, A., Nightingale, K., Jeffers, G., Fortes, E., Kongo, J. M. & Wiedmann, M. (2006) Genetic and phenotypic characterization of *Listeria monocytogenes* lineage III. *Microbiology* 152: 685-693.
- Rocourt, J., Audurier, A., Courtieu, A. L., Durst, J., Ortel, S., Schrettenbrunner, A. & Taylor, A. G. (1985) A multi-centre study on the phage typing of *Listeria monocytogenes*. *Zentralbl Bakteriol Mikrobiol Hyg A* 259: 489-497.
- Rocourt, J., Boerlin, P., Grimont, F., Jacquet, C. & Piffaretti, J.-C. (1992) Assignment of *Listeria grayi* and *Listeria murrayi* to a single species, *Listeria grayi*, with a revised description of *Listeria grayi*. *Int J Syst Bacteriol* 42: 171-174.
- Rocourt, J. & Grimont, P. A. D. (1983) *Listeria welshimeri* sp. nov. and *Listeria seeligeri* sp. nov. *Int J Syst Evol Microbiol* 33: 866-869.
- Romanova, N., Favrin, S. & Griffiths, M. W. (2002) Sensitivity of *Listeria monocytogenes* to sanitizers used in the meat processing industry. *Appl Environ Microbiol* 68: 6405-6409.
- Romanova, N. A., Wolffs, P. F. G., Brovko, L. Y. & Griffiths, M. W. (2006) Role of efflux pumps in adaptation and resistance of *Listeria monocytogenes* to benzalkonium chloride. *Appl Environ Microbiol* 72: 3498-3503.

- Rouquette, C., Ripio, M.-T., Pellegrini, E., Bolla, J.-M., Tascon, R. I., Vázquez-Boland, J.-A. & Berche, P. (1996) Identification of a ClpC ATPase required for stress tolerance and in vivo survival of *Listeria monocytogenes*. *Mol Microbiol* 21: 977-987.
- Russell, A. D. (2003) Lethal effects of heat on bacterial physiology and structure. *Sci Prog* 86: 115-137.
- Russell, N. J., Evans, R. I., ter Steeg, P. F., Hellemons, J., Verheul, A. & Abee, T. (1995) Membranes as a target for stress adaptation. *Int J Food Microbiol* 28: 255-261.
- Ruusunen, M., Salonen, M., Pulkkinen, H., Huuskonen, M., Hellström, S., Revez, J., Hänninen, M.-L., Fredriksson-Ahomaa, M. & Lindström, M. (2013) Pathogenic bacteria in Finnish bulk tank milk. *Foodborne Pathog Dis* 10: 99-106.
- Ryan, S., Begley, M., Gahan, C. G. M. & Hill, C. (2009) Molecular characterization of the arginine deiminase system in *Listeria monocytogenes*: regulation and role in acid tolerance. *Environ Microbiol* 11: 432-445.
- Ryan, S., Begley, M., Hill, C. & Gahan, C. G. M. (2010) A five-gene stress survival islet (SSI-1) that contributes to the growth of *Listeria monocytogenes* in suboptimal conditions. *J Appl Microbiol* 109: 984-995.
- Ryser, E. T., Arimi, S. M. & Donnelly, C. W. (1997) Effects of pH on distribution of *Listeria* ribotypes in corn, hay, and grass silage. *Appl Environ Microbiol* 63: 3695-3697.
- Rørvik, L. M., Aase, B., Alvestad, T. & Caugant, D. A. (2003) Molecular epidemiological survey of *Listeria monocytogenes* in broilers and poultry products. *J Appl Microbiol* 94: 633-640.
- Rørvik, L. M., Caugant, D. A. & Yndestad, M. (1995) Contamination pattern of *Listeria monocytogenes* and other *Listeria* spp. in a salmon slaughterhouse and smoked salmon processing plant. *Int J Food Microbiol* 25: 19-27.
- Salamina, G., Dalle Donne, E., Niccolini, A., Poda, G., Cesaroni, D., Bucci, M., Fini, R., Maldini, M., Schuchat, A., Swaminathan, B., Bibb, W., Rocourt, J., Binkin, N. & Salmaso, S. (1996) A foodborne outbreak of gastroenteritis involving *Listeria monocytogenes*. *Epidemiol Infect* 117: 429-436.
- Salcedo, C., Arreaza, L., Alcalá, B., de la Fuente, L. & Vázquez, J. A. (2003) Development of a multilocus sequence typing method for analysis of *Listeria monocytogenes* clones. *J Clin Microbiol* 41: 757-762.
- Salipante, S. J., SenGupta, D. J., Cummings, L. A., Land, T. A., Hoogestraat, D. R. & Cookson, B. T. (2015) Application of whole-genome sequencing for bacterial strain typing in molecular epidemiology. *J Clin Microbiol* 53: 1072-1079.
- Sampathkumar, B., Khachatourians, G. G. & Korber, D. R. (2003) High pH during trisodium phosphate treatment causes membrane damage and destruction of *Salmonella enterica* serovar Enteritidis.

- Appl Environ Microbiol* 69: 122-129.
- Sapuan, S., Kortsalioudaki, C., Anthony, M., Chang, J., Embleton, N. D., Geethanath, R. M., Gray, J., Greenough, A., Lal, M. K., Luck, S., Pattanayak, S., Reynolds, P., Russell, A. B., Scorrer, T., Turner, M., Heath, P. T. & Vergnano, S. (2017) Neonatal listeriosis in the UK 2004-2014. *J Infect* 74: 236-242.
- Schjørring, S., Gillesberg Lassen, S., Jensen, T., Moura, A., Kjeldgaard, J. S., Müller, L., Thielke, S., Leclercq, A., Maury, M. M., Tourdjman, M., Donguy, M.-P., Lecuit, M., Ethelberg, S. & Nielsen, E. M. (2017) Cross-border outbreak of listeriosis caused by cold-smoked salmon, revealed by integrated surveillance and whole genome sequencing (WGS), Denmark and France, 2015 to 2017. *Euro Surveill* 22: 17-00762.
- Schlech, W. F., III. (2000) Foodborne listeriosis. *Clin Infect Dis* 31: 770-775.
- Schlech, W. F., III, Lavigne, P. M., Bortolussi, R. A., Allen, A. C., Haldane, E. V., Wort, A. J., Hightower, A. W., Johnson, S. E., King, S. H., Nicholls, E. S. & Broome, C. V. (1983) Epidemic listeriosis – evidence for transmission by food. *N Engl J Med* 308: 203-206.
- Schmid, B., Klumpp, J., Raimann, E., Loessner, M. J., Stephan, R. & Tasara, T. (2009) Role of cold shock proteins in growth of *Listeria monocytogenes* under cold and osmotic stress conditions. *Appl Environ Microbiol* 75: 1621-1627.
- Schmitz-Esser, S., Müller, A., Stessl, B. & Wagner, M. (2015) Genomes of sequence type 121 *Listeria monocytogenes* strains harbor highly conserved plasmids and prophages. *Front Microbiol* 6: 380.
- Schröder, G. & Lanka, E. (2005) The mating pair formation system of conjugative plasmids – a versatile secretion machinery for transfer of proteins and DNA. *Plasmid* 54: 1-25.
- Schwartz, B., Ciesielski, C. A., Broome, C. V., Gaventa, S., Brown, G. R., Gellin, B. G., Hightower, A. W., Mascola, L. & the Listeriosis Study Group. (1988) Association of sporadic listeriosis with consumption of uncooked hot dogs and undercooked chicken. *Lancet* 332: 779-782.
- Schweizer, C., Bischoff, P., Bender, J., Kola, A., Gastmeier, P., Hummel, M., Klefisch, F.-R., Schoenrath, F., Frühauf, A. & Pfeifer, Y. (2019) Plasmid-mediated transmission of KPC-2 carbapenemase in *Enterobacteriaceae* in critically ill patients. *Front Microbiol* 10: 276.
- Seeliger, H. P. (1984) Modern taxonomy of the *Listeria* group relationship to its pathogenicity. *Clin Invest Med* 7: 217-221.
- Seeliger, H. P. R. & Höhne, K. (1979) Chapter II Serotyping of *Listeria monocytogenes* and related species. *Methods Microbiol* 13: 31-49.
- Seeliger, H. P. R., Rocourt, J., Schrettenbrunner, A., Grimont, P. A. D. & Jones, D. (1984) *Listeria ivanovii* sp. nov. *Int J Syst Bacteriol* 34: 336-337.
- Seydlová, G., Halada, P., Fišer, R., Toman, O., Ulrych, A. & Svobodová, J. (2012) DnaK and GroEL chaperones are recruited to the *Bacillus subtilis* membrane after

- short-term ethanol stress. *J Appl Microbiol* 112: 765-774.
- Shabala, L., Budde, B., Ross, T., Siegmundfeldt, H. & McMeekin, T. (2002) Responses of *Listeria monocytogenes* to acid stress and glucose availability monitored by measurements of intracellular pH and viable counts. *Int J Food Microbiol* 75: 89-97.
- Shen, Q., Jangam, P. M., Soni, K. A., Nannapaneni, R., Schilling, W. & Silva, J. L. (2014) Low, medium, and high heat tolerant strains of *Listeria monocytogenes* and increased heat stress resistance after exposure to sublethal heat. *J Food Prot* 77: 1298-1307.
- Shi, C., Velázquez, P., Hohl, T. M., Leiner, I., Dustin, M. L. & Pamer, E. G. (2010) Monocyte trafficking to hepatic sites of bacterial infection is chemokine independent and directed by focal intercellular adhesion molecule-1 expression. *J Immunol* 184: 6266-6274.
- Shimizu, T., Shima, K., Yoshino, K., Yonezawa, K., Shimizu, T. & Hayashi, H. (2002) Proteome and transcriptome analysis of the virulence genes regulated by the VirR/VirS system in *Clostridium perfringens*. *J Bacteriol* 184: 2587-2594.
- Silk, B. J., Date, K. A., Jackson, K. A., Pouillot, R., Holt, K. G., Graves, L. M., Ong, K. L., Hurd, S., Meyer, R., Marcus, R., Shiferaw, B., Norton, D. M., Medus, C., Zansky, S. M., Cronquist, A. B., Henao, O. L., Jones, T. F., Vugia, D. J., Farley, M. M. & Mahon, B. E. (2012) Invasive listeriosis in the Foodborne Diseases Active Surveillance Network (FoodNet), 2004-2009: further targeted prevention needed for higher-risk groups. *Clin Infect Dis* 54: S396-S404.
- Silveira, M. G., Baumgärtner, M., Rombouts, F. M. & Abee, T. (2004) Effect of adaptation to ethanol on cytoplasmic and membrane protein profiles of *Oenococcus oeni*. *Appl Environ Microbiol* 70: 2748-2755.
- Sim, J., Hood, D., Finnie, L., Wilson, M., Graham, C., Brett, M. & Hudson, J. A. (2002) Series of incidents of *Listeria monocytogenes* non-invasive febrile gastroenteritis involving ready-to-eat meats. *Lett Appl Microbiol* 35: 409-413.
- Simon, C. (1956) Möglichkeiten zur Anreicherung von *Listeria monocytogenes* in flüssigen Vorkulturen. *Med Microbiol Immunol* 143: 159-172.
- Siryaporn, A. & Goulian, M. (2010) Characterizing cross-talk *in vivo*: avoiding pitfalls and overinterpretation. *Methods Enzymol* 471: 1-16.
- Skandamis, P. N., Yoon, Y., Stopforth, J. D., Kendall, P. A. & Sofos, J. N. (2008) Heat and acid tolerance of *Listeria monocytogenes* after exposure to single and multiple sublethal stresses. *Food Microbiol* 25: 294-303.
- Skovgaard, N. & Nørrung, B. (1989) The incidence of *Listeria* spp. in faeces of Danish pigs and in minced pork meat. *Int J Food Microbiol* 8: 59-63.
- Slater, F. R., Bailey, M. J., Tett, A. J. & Turner, S. L. (2008) Progress towards understanding the fate of plasmids in bacterial communities. *FEMS Microbiol Ecol* 66: 3-13.

- Sleator, R. D., Gahan, C. G. M. & Hill, C. (2001) Identification and disruption of the *proBA* locus in *Listeria monocytogenes*: role of proline biosynthesis in salt tolerance and murine infection. *Appl Environ Microbiol* 67: 2571-2577.
- Sleator, R. D. & Hill, C. (2005) A novel role for the LisRK two-component regulatory system in listerial osmotolerance. *Clin Microbiol Infect* 11: 599-601.
- Slettemeås, J. S., Sunde, M., Ulstad, C. R., Norström, M., Wester, A. L. & Urdahl, A. M. (2019) Occurrence and characterization of quinolone resistant *Escherichia coli* from Norwegian turkey meat and complete sequence of an IncX1 plasmid encoding *qnrS1*. *PLoS One* 14: e0212936.
- Small, P. L. C. & Waterman, S. R. (1998) Acid stress, anaerobiosis and *gadCB*: lessons from *Lactococcus lactis* and *Escherichia coli*. *Trends Microbiol* 6: 214-216.
- Smillie, C., Garcillán-Barcia, M. P., Francia, M. V., Rocha, E. P. C. & de la Cruz, F. (2010) Mobility of plasmids. *Microbiol Mol Biol Rev* 74: 434-452.
- Snapir, Y. M., Vaisbein, E. & Nassar, F. (2006) Low virulence but potentially fatal outcome – *Listeria ivanovii*. *Eur J Intern Med* 17: 286-287.
- Solovyev, V. & Salamov, A. (2011) Automatic annotation of microbial genomes and metagenomic sequences. In *Metagenomics and its applications in agriculture, biomedicine and environmental studies*. Li, R. W. (ed). New York, Nova Science Publishers, pp. 61-78.
- Soni, K. A., Nannapaneni, R. & Tasara, T. (2011) The contribution of transcriptomic and proteomic analysis in elucidating stress adaptation responses of *Listeria monocytogenes*. *Foodborne Pathog Dis* 8: 843-852.
- Soufi, B., Krug, K., Harst, A. & Macek, B. (2015) Characterization of the *E. coli* proteome and its modifications during growth and ethanol stress. *Front Microbiol* 6: 103.
- Soumet, C., Ragimbeau, C. & Maris, P. (2005) Screening of benzalkonium chloride resistance in *Listeria monocytogenes* strains isolated during cold smoked fish production. *Lett Appl Microbiol* 41: 291-296.
- Stack, H. M., Sleator, R. D., Bowers, M., Hill, C. & Gahan, C. G. M. (2005) Role for HtrA in stress induction and virulence potential in *Listeria monocytogenes*. *Appl Environ Microbiol* 71: 4241-4247.
- Stock, A. M., Robinson, V. L. & Goudreau, P. N. (2000) Two-component signal transduction. *Annu Rev Biochem* 69: 183-215.
- Sue, D., Fink, D., Wiedmann, M. & Boor, K. J. (2004) σ^B -dependent gene induction and expression in *Listeria monocytogenes* during osmotic and acid stress conditions simulating the intestinal environment. *Microbiology* 150: 3843-3855.
- Suo, Y., Gao, S., Baranzoni, G. M., Xie, Y. & Liu, Y. (2018) Comparative transcriptome RNA-Seq analysis of *Listeria monocytogenes* with sodium lactate

- adaptation. *Food Control* 91: 193-201.
- Suokko, A., Savijoki, K., Malinen, E., Palva, A. & Varmanen, P. (2005) Characterization of a mobile *clpL* gene from *Lactobacillus rhamnosus*. *Appl Environ Microbiol* 71: 2061-2069.
- Swaminathan, B. & Gerner-Smidt, P. (2007) The epidemiology of human listeriosis. *Microbes Infect* 9: 1236-1243.
- Szurmant, H., Fukushima, T. & Hoch, J. A. (2007) The essential YycFG two-component system of *Bacillus subtilis*. *Methods Enzymol* 422: 396-417.
- Takada, H., Shiwa, Y., Takino, Y., Osaka, N., Ueda, S., Watanabe, S., Chibazakura, T., Su'etsugu, M., Utsumi, R. & Yoshikawa, H. (2018) Essentiality of WalRK for growth in *Bacillus subtilis* and its role during heat stress. *Microbiology* 164: 670-684.
- Tandukar, M., Oh, S., Tezel, U., Konstantinidis, K. T. & Pavlostathis, S. G. (2013) Long-term exposure to benzalkonium chloride disinfectants results in change of microbial community structure and increased antimicrobial resistance. *Environ Sci Technol* 47: 9730-9738.
- Taormina, P. J. & Beuchat, L. R. (2001) Survival and heat resistance of *Listeria monocytogenes* after exposure to alkali and chlorine. *Appl Environ Microbiol* 67: 2555-2563.
- Tasara, T. & Stephan, R. (2006) Cold stress tolerance of *Listeria monocytogenes*: a review of molecular adaptive mechanisms and food safety implications. *J Food Prot* 69: 1473-1484.
- Tasara, T. & Stephan, R. (2007) Evaluation of housekeeping genes in *Listeria monocytogenes* as potential internal control references for normalizing mRNA expression levels in stress adaptation models using real-time PCR. *FEMS Microbiol Lett* 269: 265-272.
- Tay, E., Rajan, M. & Tuft, S. (2008) *Listeria monocytogenes* sclerokeratitis: a case report and literature review. *Cornea* 27: 947-949.
- Thackray, P. D. & Moir, A. (2003) SigM, an extracytoplasmic function sigma factor of *Bacillus subtilis*, is activated in response to cell wall antibiotics, ethanol, heat, acid, and superoxide stress. *J Bacteriol* 185: 3491-3498.
- Thomas, C. M. & Nielsen, K. M. (2005) Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nat Rev Microbiol* 3: 711-721.
- Thomas, C. M. & Smith, C. A. (1987) Incompatibility group P plasmids: genetics, evolution, and use in genetic manipulation. *Annu Rev Microbiol* 41: 77-101.
- Tiwari, S., Jamal, S. B., Hassan, S. S., Carvalho, P. V. S. D., Almeida, S., Barh, D., Ghosh, P., Silva, A., Castro, T. L. P. & Azevedo, V. (2017) Two-component signal transduction systems of pathogenic bacteria as targets for antimicrobial therapy: an overview. *Front Microbiol* 8: 1878.
- Toledo-Arana, A., Dussurget, O., Nikitas, G., Sesto, N., Guet-Revillet, H., Balestrino, D., Loh, E., Gripenland, J., Tiensuu, T., Vaitkevicius, K., Barthelemy, M., Vergassola, M., Nahori, M.-A., Soubigou, G., Régnault,

- B., Coppée, J.-Y., Lecuit, M., Johansson, J. & Cossart, P. (2009) The *Listeria* transcriptional landscape from saprophytism to virulence. *Nature* 459: 950-956.
- Tompkin, R. (2002) Control of *Listeria monocytogenes* in the food-processing environment. *J Food Prot* 65: 709-725.
- Tsai, Y.-H. L., Maron, S. B., McGann, P., Nightingale, K. K., Wiedmann, M. & Orsi, R. H. (2011) Recombination and positive selection contributed to the evolution of *Listeria monocytogenes* lineages III and IV, two distinct and well supported uncommon *L. monocytogenes* lineages. *Infect Genet Evol* 11: 1881-1890.
- Unnerstad, H., Romell, A., Ericsson, H., Danielsson-Tham, M.-L. & Tham, W. (2000) *Listeria monocytogenes* in faeces from clinically healthy dairy cows in Sweden. *Acta Vet Scand* 41: 167-171.
- van Bokhorst-van de Veen, H., Abee, T., Tempelaars, M., Bron, P. A., Kleerebezem, M. & Marco, M. L. (2011) Short- and long-term adaptation to ethanol stress and its cross-protective consequences in *Lactobacillus plantarum*. *Appl Environ Microbiol* 77: 5247-5256.
- van der Veen, S., Abee, T., de Vos, W. M. & Wells-Bennik, M. H. J. (2009) Genome-wide screen for *Listeria monocytogenes* genes important for growth at high temperatures. *FEMS Microbiol Lett* 295: 195-203.
- van der Veen, S., Hain, T., Wouters, J. A., Hossain, H., de Vos, W. M., Abee, T., Chakraborty, T. & Wells-Bennik, M. H. J. (2007) The heat-shock response of *Listeria monocytogenes* comprises genes involved in heat shock, cell division, cell wall synthesis, and the SOS response. *Microbiology* 153: 3593-3607.
- van der Veen, S., Moezelaar, R., Abee, T. & Wells-Bennik, M. H. J. (2008) The growth limits of a large number of *Listeria monocytogenes* strains at combinations of stresses show serotype- and niche-specific traits. *J Appl Microbiol* 105: 1246-1258.
- Vázquez-Boland, J. A., Dominguez, L., Blanco, M., Rocourt, J., Fernández-Garayzábal, J. F., Gutiérrez, C. B., Tascón, R. I. & Rodríguez-Ferri, E. F. (1992) Epidemiologic investigation of a silage-associated epizootic of ovine listeric encephalitis, using a new *Listeria*-selective enumeration medium and phage typing. *Am J Vet Res* 53: 368-371.
- Vázquez-Boland, J. A., Domínguez-Bernal, G., González-Zorn, B., Kreft, J. & Goebel, W. (2001) Pathogenicity islands and virulence evolution in *Listeria*. *Microbes Infect* 3: 571-584.
- Vázquez-Boland, J. A., Kryptou, E. & Scortti, M. (2017) *Listeria* placental infection. *MBio* 8: e00949-17.
- Vázquez-Boland, J. A., Kuhn, M., Berche, P., Chakraborty, T., Domínguez-Bernal, G., Goebel, W., González-Zorn, B., Wehland, J. & Kreft, J. (2001) *Listeria* pathogenesis and molecular virulence determinants. *Clin Microbiol Rev* 14: 584-640.
- Venturini, C., Beatson, S. A., Djordjevic, S. P. & Walker, M. J. (2010) Multiple antibiotic resistance gene

- recruitment onto the enterohemorrhagic *Escherichia coli* virulence plasmid. *FASEB J* 24: 1160-1166.
- Vialette, M., Pinon, A., Chasseignaux, E. & Lange, M. (2003) Growth kinetics comparison of clinical and seafood *Listeria monocytogenes* isolates in acid and osmotic environment. *Int J Food Microbiol* 82: 121-131.
- Vines, A., Reeves, M. W., Hunter, S. & Swaminathan, B. (1992) Restriction fragment length polymorphism in four virulence-associated genes of *Listeria monocytogenes*. *Res Microbiol* 143: 281-294.
- Vivant, A. L., Garmyn, D., Gal, L., Hartmann, A. & Piveteau, P. (2015) Survival of *Listeria monocytogenes* in soil requires AgrA-mediated regulation. *Appl Environ Microbiol* 81: 5073-5084.
- Vivant, A. L., Garmyn, D., Gal, L. & Piveteau, P. (2014) The Agr communication system provides a benefit to the populations of *Listeria monocytogenes* in soil. *Front Cell Infect Microbiol* 4: 160.
- Vogel, B. F., Jørgensen, L. V., Ojeniyi, B., Huss, H. H. & Gram, L. (2001) Diversity of *Listeria monocytogenes* isolates from cold-smoked salmon produced in different smokehouses as assessed by Random Amplified Polymorphic DNA analyses. *Int J Food Microbiol* 65: 83-92.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., van de Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. & Zabeau, M. (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23: 4407-4414.
- Walker, S. J., Archer, P. & Banks, J. G. (1990) Growth of *Listeria monocytogenes* at refrigeration temperatures. *J Appl Bacteriol* 68: 157-162.
- Ward, T. J., Ducey, T. F., Usgaard, T., Dunn, K. A. & Bielawski, J. P. (2008) Multilocus genotyping assays for single nucleotide polymorphism-based subtyping of *Listeria monocytogenes* isolates. *Appl Environ Microbiol* 74: 7629-7642.
- Ward, T. J., Gorski, L., Borucki, M. K., Mandrell, R. E., Hutchins, J. & Pupedis, K. (2004) Intraspecific phylogeny and lineage group identification based on the *prfA* virulence gene cluster of *Listeria monocytogenes*. *J Bacteriol* 186: 4994-5002.
- Warrens, A. N., Jones, M. D. & Lechler, R. I. (1997) Splicing by overlap extension by PCR using asymmetric amplification: an improved technique for the generation of hybrid proteins of immunological interest. *Gene* 186: 29-35.
- Wawrzynow, A., Banecki, B. & Zylicz, M. (1996) The Clp ATPases define a novel class of molecular chaperones. *Mol Microbiol* 21: 895-899.
- Weis, J. & Seeliger, H. P. R. (1975) Incidence of *Listeria monocytogenes* in nature. *Appl Microbiol* 30: 29-32.
- Weiss, V., Kramer, G., Dünnebier, T. & Flotho, A. (2002) Mechanism of regulation of the bifunctional histidine kinase NtrB in *Escherichia coli*. *J Mol Microbiol Biotechnol* 4: 229-233.
- Weller, D., Andrus, A., Wiedmann, M. & den Bakker, H. C. (2015) *Listeria booriae* sp. nov. and *Listeria newyorkensis* sp. nov., from food processing

- environments in the USA. *Int J Syst Evol Microbiol* 65: 286-292.
- Welshimer, H. J. & Meredith, A. L. (1971) *Listeria murrayi*. sp. n.: a nitrate-reducing mannitol-fermenting *Listeria*. *Int J Syst Evol Microbiol* 21: 3-7.
- Wemekamp-Kamphuis, H. H., Sleator, R. D., Wouters, J. A., Hill, C. & Abee, T. (2004a) Molecular and physiological analysis of the role of osmolyte transporters BetL, Gbu, and OpuC in growth of *Listeria monocytogenes* at low temperatures. *Appl Environ Microbiol* 70: 2912-2918.
- Wemekamp-Kamphuis, H. H., Wouters, J. A., de Leeuw, P. P., Hain, T., Chakraborty, T. & Abee, T. (2004b) Identification of sigma factor σ^B -controlled genes and their impact on acid stress, high hydrostatic pressure, and freeze survival in *Listeria monocytogenes* EGD-e. *Appl Environ Microbiol* 70: 3457-3466.
- West, A. H. & Stock, A. M. (2001) Histidine kinases and response regulator proteins in two-component signaling systems. *Trends Biochem Sci* 26: 369-376.
- Wiedmann, M., Arvik, T. J., Hurley, R. J. & Boor, K. J. (1998) General stress transcription factor σ^B and its role in acid tolerance and virulence of *Listeria monocytogenes*. *J Bacteriol* 180: 3650-3656.
- Wiedmann, M., Bruce, J. L., Keating, C., Johnson, A. E., McDonough, P. L. & Batt, C. A. (1997) Ribotypes and virulence gene polymorphisms suggest three distinct *Listeria monocytogenes* lineages with differences in pathogenic potential. *Infect Immun* 65: 2707-2716.
- Wiedmann, M., Bruce, J. L., Knorr, R., Bodis, M., Cole, E. M., McDowell, C. I., McDonough, P. L. & Batt, C. A. (1996) Ribotype diversity of *Listeria monocytogenes* strains associated with outbreaks of listeriosis in ruminants. *J Clin Microbiol* 34: 1086-1090.
- Wilesmith, J. W. & Gitter, M. (1986) Epidemiology of ovine listeriosis in Great Britain. *Vet Rec* 119: 467-470.
- Wilkins, P. O., Bourgeois, R. & Murray, R. G. E. (1972) Psychrotrophic properties of *Listeria monocytogenes*. *Can J Microbiol* 18: 543-551.
- Williams, T., Bauer, S., Beier, D. & Kuhn, M. (2005a) Construction and characterization of *Listeria monocytogenes* mutants with in-frame deletions in the response regulator genes identified in the genome sequence. *Infect Immun* 73: 3152-3159.
- Williams, T., Joseph, B., Beier, D., Goebel, W. & Kuhn, M. (2005b) Response regulator DegU of *Listeria monocytogenes* regulates the expression of flagella-specific genes. *FEMS Microbiol Lett* 252: 287-298.
- Wilson, R. L., Brown, L. L., Kirkwood-Watts, D., Warren, T. K., Lund, S. A., King, D. S., Jones, K. F. & Hruby, D. E. (2006) *Listeria monocytogenes* 10403S HtrA is necessary for resistance to cellular stress and virulence. *Infect Immun* 74: 765-768.
- Wonderling, L. D., Wilkinson, B. J. & Bayles, D. O. (2004) The *htrA* (*degP*) gene of *Listeria monocytogenes* 10403S is

- essential for optimal growth under stress conditions. *Appl Environ Microbiol* 70: 1935-1943.
- Wright, E., Neethirajan, S., Warriner, K., Retterer, S. & Srijanto, B. (2014) Single cell swimming dynamics of *Listeria monocytogenes* using a nanoporous microfluidic platform. *Lab Chip* 14: 938-946.
- Wuichet, K., Cantwell, B. J. & Zhulin, I. B. (2010) Evolution and phyletic distribution of two-component signal transduction systems. *Curr Opin Microbiol* 13: 219-225.
- Xu, D., Li, Y., Zahid, M. S. H., Yamasaki, S., Shi, L., Li, J.-R. & Yan, H. (2014) Benzalkonium chloride and heavy-metal tolerance in *Listeria monocytogenes* from retail foods. *Int J Food Microbiol* 190: 24-30.
- Yates, C. M., Shaw, D. J., Roe, A. J., Woolhouse, M. E. J. & Amyes, S. G. B. (2006) Enhancement of bacterial competitive fitness by apramycin resistance plasmids from non-pathogenic *Escherichia coli*. *Biol Lett* 2: 463-465.
- Yu, T., Jiang, X., Zhang, Y., Ji, S., Gao, W. & Shi, L. (2018) Effect of benzalkonium chloride adaptation on sensitivity to antimicrobial agents and tolerance to environmental stresses in *Listeria monocytogenes*. *Front Microbiol* 9: 2906.
- Zetzmann, M., Sánchez-Kopper, A., Waidmann, M. S., Blombach, B. & Riedel, C. U. (2016) Identification of the *agr* peptide of *Listeria monocytogenes*. *Front Microbiol* 7: 989.
- Zhang, H., Hu, Y., Zhou, C., Yang, Z., Wu, L., Zhu, M., Bao, H., Zhou, Y., Pang, M., Wang, R. & Zhou, X. (2018) Stress resistance, motility and biofilm formation mediated by a 25kb plasmid pLMSZo8 in *Listeria monocytogenes*. *Food Control* 94: 345-352.
- Zhou, Y., Liang, Y., Lynch, K. H., Dennis, J. J. & Wishart, D. S. (2011) PHAST: a fast phage search tool. *Nucleic Acids Res* 39: W347-W352.
- Zilberstein, D., Agmon, V., Schuldiner, S. & Padan, E. (1984) *Escherichia coli* intracellular pH, membrane potential, and cell growth. *J Bacteriol* 158: 246-252.
- Zschiedrich, C. P., Keidel, V. & Szurmant, H. (2016) Molecular mechanisms of two-component signal transduction. *J Mol Biol* 428: 3752-3775.
- Zuber, U. & Schumann, W. (1994) CIRCE, a novel heat shock element involved in regulation of heat shock operon *dnaK* of *Bacillus subtilis*. *J Bacteriol* 176: 1359-1363.